

## Asymptotics In-Class Practice Problems

Recall the definitions we have introduced in class:

### Big-Oh

Let  $f, g: \mathbb{N} \rightarrow \mathbb{R}$  such that  $g(n) \geq 0$ .

We say that  $f(n) \in O(g(n))$  or  $f \in O(g)$  if  $\exists B, b$ , both positive, such that

$$0 \leq |f(n)| \leq Bg(n) \quad \forall n \geq b$$

### Big-Omega

Let  $f, g: \mathbb{N} \rightarrow \mathbb{R}^+$ . We say that  $f(n) \in \Omega(g(n))$  or  $f \in \Omega(g)$  if  $\exists A, a$ , both positive, such that

$$0 \leq Ag(n) \leq f(n) \quad \forall n \geq b$$

### Big-Theta

Let  $f, g: \mathbb{N} \rightarrow \mathbb{R}^+$ . We say that  $f(n) \in \Theta(g(n))$  or  $f \in \Theta(g)$  if  $\exists A, B, k$ , all positive, such that

$$Ag(n) \leq f(n) \leq Bg(n) \quad \forall n \geq k$$

### Practice Questions:

1. Is  $2^{n+1} \in O(2^n)$ ? What about  $2^{2n}$ ? Provide proof.
2. Suppose  $f \in O(g)$ . Is  $kf(n) \in O(g(n))$  for  $k \in \mathbb{R}, k \neq 0$ ? If so, prove it. If not, provide a counterexample.
3. For non-negative functions, is the relation big-Oh (provide proof):
  - a. Reflexive?
  - b. Symmetric? [cf Homework 4]
  - c. Transitive?
4. Show that for any  $k, d \geq 0$ ,  $n^k \in O(n^d) \Leftrightarrow d \geq k$
5. Let  $f(n) = n^2$  and  $g(n) = \frac{1}{2}n^2$ . Show that:  $2^{f(n)} \notin O(2^{g(n)})$
6. Let  $f, g: \mathbb{N} \rightarrow \mathbb{R}^+$ . Show that  $f \in O(g) \Leftrightarrow g \in \Omega(f)$
7. Let  $f, g: \mathbb{N} \rightarrow \mathbb{R}^+$ . Show that  $f \in \Theta(g) \Leftrightarrow f \in O(g) \text{ and } f \in \Omega(g)$
8. Using any previous results on this practice sheet or from class/HWs/labs, show the following. Do not use  $A, B, k, a, b$ , etc.:
  - a.  $\Omega$  is reflexive
  - b.  $\Omega$  is not symmetric
  - c.  $\Omega$  is transitive