

# Week 7 Homework

## Exercise

Find a recurrence relation for the number,  $b_n$  of length  $n$  binary strings with no two consecutive ones (No 11 in the string). What are the initial conditions? What is the closed form solution to this recurrence relationship?

## Exercise

Solve the following homogeneous recurrence relations:

- a.  $a_n = 5a_{n-1} - 6a_{n-2}$  with  $a_1 = -1$  and  $a_2 = 1$
- b.  $b_n = 6b_{n-1} - 9b_{n-2}$  with  $b_1 = 1$  and  $b_2 = 9$

## Exercise

Solve the following homogeneous recurrence relations:

- a.  $a_n = 6a_{n-1} - 8a_{n-2}$ ;  $a_0 = 1$ ,  $a_1 = 0$
- b.  $a_n = 2a_{n-1} + 8a_{n-2}$ ;  $a_0 = 4$ ,  $a_1 = 10$

## Exercise

Consider tiling a  $2 \times n$  board with  $2 \times 1$  and  $2 \times 2$  tiles.

- a. Let  $a_n$  count the number of tilings of where the  $2 \times 1$  can only be positioned vertically. Find a recursive relationship for  $a_n$ . Find initial conditions for  $a_n$ .
- b. Let  $b_n$  count the number of tilings of where the  $2 \times 1$  can only be positioned horizontally. Find a recursive relationship for  $b_n$  and find initial conditions for  $b_n$ .
- c. Let  $c_n$  count the number of tilings of where the  $2 \times 1$  can be positioned horizontally or vertically. Find a recursive relationship for  $c_n$  and find initial conditions for  $c_n$ .

**i** Exercise

Let  $s_n$  be the number of length- $n$  bit strings with no three consecutive 0s. Find  $s_1, s_2, s_3$  and find a recursive definition of this sequence.

**i** Exercise

Find a closed form solution to recurrence relation  $x_n = n$  for  $0 \leq n < m$  and  $x_n = x_{n-m} + 1$  for  $n \geq m$ .

**i** Exercise

What is dynamic programming? Describe the basic algorithm and explain its connection to recurrence relations.

**i** Exercise

What is the Josephus number? Define it and give its historical origins. What is its connection to recurrence relations?

**i** Exercise

A string that contains only 0's, 1's, and 2's is called a **ternary string**. Find a recurrence relation for the number of ternary strings that contain two consecutive 0s? What are the initial conditions? Explain your answer. What is the recursion and initial conditions for ternary strings that contain 000?

**i** Exercise

Each day Angela eats lunch at a deli, ordering one of the following: chicken salad, a tuna sandwich, or a turkey wrap. Find a recurrence relation for the number of ways for her to order lunch for the " $n$ " days if she never orders chicken salad three days in a row.

**i** Exercise

Find the number of non-negative integer solutions to  $x + y + z + w \leq 6$ . Generalize this to find the number of non-negative integer solutions to  $x + y + z + w \leq k$ .

**i** Exercise

Let  $F_{n+2} = F_{n+1} + F_n$  with  $F_0 = 0$  and  $F_1 = 1$ ,  $L_{n+2} = L_{n+1} + L_n$  with  $L_0 = 2$  and  $L_1 = 1$ . Find a relationship between  $L_n$ ,  $F_{n-1}$ , and  $F_{n+1}$ .

**i** Exercise

Find the first 5 terms of the Taylor expansion at  $x = 0$  of  $\sqrt{1-4x}$ . By examining the pattern of coefficients find  $a_n$  where  $\sqrt{1-4x} = \sum_{n=0}^{\infty} a_n x^n$ .

**i** Exercise

Use the Auxiliary Equation method to find a closes form solution to the following recurrence relations:

- a.  $c_n = 3c_{n-1} - 3c_{n-2} + c_{n-3}$  with  $c_0 = 1$ ,  $c_1 = 2$ , and  $c_2 = 4$
- b.  $x_n = 5x_{n-1} - 4x_{n-2} - 6x_{n-3}$  with  $x_0 = 0$ ,  $x_1 = 0$ , and  $x_2 = 1$

**i** Proof

Prove

$$\frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^n(n+1)!} 4^n = \frac{1}{n+1} \binom{2n}{n}.$$

**i** Proof

Prove  $\binom{3n}{3} = 3\binom{n}{3} + 6n\binom{n}{2} + n^2\binom{n}{1}$ .

**i** Proof

Using induction, prove that for the sequence defined by  $a_1 = 0$ ,  $a_2 = 1/2$ , and  $a_{n+2} = \frac{1}{2}(a_n + a_{n+1})$  for  $n \geq 1$  that

$$a_n = \frac{1}{3} \left( 1 - \left( -\frac{1}{2} \right)^{n-1} \right)$$

for every  $n \geq 1$ .

**i** Proof

Consider the sequence defined by  $a_n = \frac{1}{2}a_{n-1} + c$  where  $a_1 = c$ . Prove that  $a_n = c(2 - \frac{1}{2^{n-1}})$  using mathematical induction.

**i** Proof

Prove

$$\sum_{k=0}^n k \binom{n}{k} = n2^{n-1}$$

**i** Proof

Let  $T_n = \frac{n(n+1)}{2}$  be the triangular numbers and  $P_n = \frac{3n^2-n}{2}$  be the pentagonal numbers. Using the direct proof method, prove  $T_{5n-2} = T_{2n-2} + 6P_n + 3T_{n-1}$  for all  $n \geq 1$ .

**i** Proof

Prove

$$\sum_{i=0}^k (-1)^i \binom{n}{i} = (-1)^k \binom{n-1}{k}$$

**i** Proof

Prove

$$\sum_{k=0}^m k \binom{m}{k} \binom{n}{k} = n \binom{m+n-1}{n}$$

Hint: See Example 1.10 and 1.11 on page 9 of the course text.

**i** Proof

Consider the following modification of the Towers of Hanoi problem with three posts labeled A, B, and C with the added two rules 1) A disk may be moved only from a post and adjacent post, and 2) the disks must be moved from post A to post C. Justify a recursive description of this problem, and prove by induction that the number of moves need to move  $n$  disks is  $3^n - 1$ .

**i** Proof

Let  $F_{n+2} = F_{n+1} + F_n$  with  $F_0 = 0$  and  $F_1 = 1$ , the Fibonacci numbers, and  $L_{n+2} = L_{n+1} + L_n$  with  $L_0 = 2$  and  $L_1 = 1$ , the Lucas numbers. Prove

$$\sum_{k=0}^n 2^k L_k = 2^{n+1} F_{n+1}$$

for all  $n \geq 0$ .

**i** Proof

Let  $F_{n+2} = F_{n+1} + F_n$  with  $F_0 = 0$  and  $F_1 = 1$ , the Fibonacci numbers, and  $L_{n+2} = L_{n+1} + L_n$  with  $L_0 = 2$  and  $L_1 = 1$ , the Lucas numbers. Prove

$$\sum_{k=0}^n (-1)^k 2^{n-k} L_{k+1} = (-1)^n F_{n+1}$$

for all  $n \geq 0$ .

**i** Proof

Define  $x_1 = 1$  and  $x_2 = 2$  and  $x_{n+2} = x_{n+1} + x_n$  for  $n \geq 1$ . Prove that  $4^n x_n < 7^n$  for all positive integers  $n$ .

**i** Proof

Prove  $\binom{n^2}{2} = n\binom{n}{2} + n^2\binom{n}{2}$ .