DEPARTMENT OF SCIENCE AND TECHNOLOGY
FOOD AND NUTRITION RESEARCH INSTITUTE

DOST-FNRI

NATIONAL NUTRITION SUMMIT



Halfway Point to 2030: Key Findings of the 2023 National Nutrition Survey

DECEMBER 10, 2024

DUSIT THANI MANILA (MAYUREE 1)


2023 NNS REGIONAL ESTIMATES of SELECTED INDICATORS



HOUSEHOLD FOOD SECURITY



Prevalence of moderate to severe household food insecurity, by region: Philippines, 2023




95% Confidence interval of the national estimate

DOST-FNRI

Prevalence of moderate to severe household food insecurity, by region: Philippines, 2023


Based on FAO's Hunger Map
0% - 9.9%
10% - 24.9%
25% - 39.9%
40% - 59.9%
60% - 79.9%
80% - 100%

Philippines: 31.4%



DOST-FNRI

Prevalence of severe household food insecurity, by region: Philippines, 2023




DOST-FNRI

CHILDREN UNDER FIVE (0-59 MONTHS)




Prevalence of underweight among children 0-59 months old, by region: Philippines, 2023



95% Confidence interval of the national estimate

DOST-FNRI

Prevalence of underweight among children 0-59 months old, by region: ENNS 2018, 2019, 2021




DOST-FNRI

Prevalence of underweight among children 0-59 months old, by region: Philippines, 2023

Magnitude and Severity (WHO, 1995)*
Low: ≤10%
Low to Medium (using 95% CI)
Medium: 10-19%
Medium to High (using 95% CI)
High: 20-29%
High to Very High (using 95% CI)
Very High: ≥30%


**using 95% confidence interval estimate of underweight*

Philippines: 15.1%



DOST-FNRI


Prevalence of stunting among children 0-59 months old, by region: Philippines, 2023



95% Confidence interval of the national estimate

DOST-FNRI

Prevalence of stunting among children 0-59 months old, by region: ENNS 2018, 2019, 2021




DOST-FNRI

Prevalence of stunting among children 0-59 months old, by region: Philippines, 2023

Public Health Significance (de Onis et al., 2019)*
Very Low: <2.5%
Very Low to Low (using 95% CI)
Low: ≤10%
Low to Medium (using 95% CI)
Medium: 10-<20%
Medium to High (using 95% CI)
High: 20 - <30%
High to Very High (using 95% CI)
Very High ≥30%


**using 95% confidence interval estimate of stunting*

Philippines: 23.6%



DOST-FNRI


Prevalence of wasting among children 0-59 months old, by region: Philippines, 2023



95% Confidence interval of the national estimate

DOST-FNRI

Prevalence of wasting among children 0-59 months old, by region: ENNS 2018, 2019, 2021




DOST-FNRI

Prevalence of wasting among children 0-59 months old, by region: Philippines, 2023

Public Health Significance (de Onis et al., 2019)*
Very Low: <2.5%
Very Low to Low (using 95% CI)
Low: 2.5<5%
Low to Medium (using 95% CI)
Medium: 5-<10%
Medium to High (using 95% CI)
High: 10-15%
High to Very High (using 95% CI)
Very High ≥15%


**using 95% confidence interval estimate of wasting*



Philippines: 5.6%

DOST-FNRI


Prevalence of overweight-for-height among children 0-59 months old, by region: Philippines, 2023



95% Confidence interval of the national estimate

DOST-FNRI

Prevalence of overweight-for-height among children 0-59 months old, by region: ENNS 2018, 2019, 2021




DOST-FNRI

Prevalence of overweight-for-height among children 0-59 months old, by region: Philippines, 2023

Public Health Significance (de Onis et al., 2019)^
Very Low: <2.5%
Very Low to Low (using 95% CI)
Low: 2.5<5%
Low to Medium (using 95% CI)
Medium: 5-<10%
Medium to High (using 95% CI)
High: 10-15%
High to Very High (using 95% CI)
Very High ≥15%


**using 95% confidence interval estimate of overweight-for-height*

Philippines: 3.7%



DOST-FNRI

Prevalence of anemia among children 6 to 59 months old, by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of anemia among children 6 to 59 months, by region: Philippines, 2023

Public Health Significance (WHO, 2011)*
Low: ≤4.9%
Low to Mild (using 95% CI)
Mild: 5-19.9%
Mild to Moderate (using 95% CI)
Moderate: 20-39.9%
Moderate to Severe (using 95% CI)
Severe: ≥40.0%

**using 95% confidence interval estimate of anemia*

Philippines: 11.4%




DOST-FNRI

SCHOOL AGE CHILDREN (5-10 YEARS OLD)




Prevalence of underweight among school-age children 5-10 years old, by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of underweight among school age children 5-10 years old, by region: ENNS 2018, 2019, 2021



DOST-FNRI


Prevalence of stunting among school-age children 5-10 years old, by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of stunting among school age children 5-10 years old, by region: ENNS 2018, 2019, 2021



DOST-FNRI


Prevalence of wasting among school age children 5-10 years old, by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of wasting among school age children 5-10 years old, by region: ENNS 2018, 2019, 2021



DOST-FNRI


Prevalence of overweight/obesity among school-age children 5-10 years old, by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of overweight/obesity among school age children 5-10 years old, by region: ENNS 2018, 2019, 2021



DOST-FNRI

Prevalence of anemia among school age children 5-10 years old by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of anemia among school age children, by region: Philippines, 2023

Public Health Significance (WHO, 2011)
Low: ≤4.9%
Low to Mild (using 95% CI)
Mild: 5-19.9%
Mild to Moderate (using 95% CI)
Moderate: 20-39.9%
Moderate to Severe (using 95% CI)
Severe: ≥40.0%

**using 95% confidence interval estimate of anemia*

Philippines: 5.4%




DOST-FNRI

ADOLESCENTS (10.08-19.0 YEARS OLD)



DOST-FNRI

Prevalence of stunting among adolescents (10.08-19.0 years old)[1] by region: Philippines, 2023




95% Confidence interval of the national estimate

[1] *121-228 months*

DOST-FNRI


Prevalence of stunting among adolescents (10.08-19.0 years old)[1] by region: ENNS 2018, 2019, 2021



[1] *121-228 months*

DOST-FNRI

Prevalence of wasting among adolescents (10.08-19.0 years old)[1] by region: Philippines, 2023




95% Confidence interval of the national estimate

[1] 121-228 months

DOST-FNRI


Prevalence of wasting among adolescents (10.08-19.0 years old)[1] by region: ENNS 2018, 2019, 2021



[1] *121-228 months*

DOST-FNRI

Prevalence of overweight/obesity among adolescents (10.08-19.0 years old)[1] by region: Philippines, 2023




95% Confidence interval of the national estimate

[1] *121-228 months*

DOST-FNRI


Prevalence of overweight/obesity among adolescents (10.08-19.0 years old)[1] by region: ENNS 2018, 2019, 2021



[1] *121-228 months*

DOST-FNRI

Prevalence of anemia among adolescents (10.08-19.0 years old)[1], by region: Philippines, 2023



95% Confidence interval of the national estimate

[1] *121-228 months*


DOST-FNRI

Prevalence of anemia among adolescents, by region: Philippines, 2023

Public Health Significance (WHO, 2011)
Low: ≤4.9%
Low to Mild (using 95% CI)
Mild: 5-19.9%
Mild to Moderate (using 95% CI)
Moderate: 20-39.9%
Moderate to Severe (using 95% CI)
Severe: ≥40.0%

**using 95% confidence interval estimate of anemia*

Philippines: 5.8%




DOST-FNRI

ADULTS
(20-59 YEARS OLD)




Prevalence of chronic energy deficiency among adults 20-59 years old, by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of chronic energy deficiency among adults 20-59 years old, by region: ENNS 2018, 2019, 2021



DOST-FNRI

Prevalence of overweight & obesity* among adults 20-59 years old, by region: Philippines, 2023




95% Confidence interval of the national estimate

** based on Asia Pacific classification and cut-off points*

DOST-FNRI

Prevalence of anemia among adults 20-59 years old, by region: Philippines, 2023




95% Confidence interval of the national estimate

DOST-FNRI

Prevalence of anemia among adults 20-50 years old, by region: Philippines, 2023

Public Health Significance (WHO, 2011)*
Low: ≤4.9%
Low to Mild (using 95% CI)
Mild: 5-19.9%
Mild to Moderate (using 95% CI)
Moderate: 20-39.9%
Moderate to Severe (using 95% CI)
Severe: ≥40.0%

**using 95% confidence interval estimate of anemia*




Philippines: 8.0%

DOST-FNRI

OLDER PERSONS (60 YEARS OLD and ABOVE)




Prevalence of chronic energy deficiency among older persons 60 years old and above, by region: Philippines, 2023



95% Confidence interval of the national estimate

DOST-FNRI

Prevalence of chronic energy deficiency among older persons 60 years old and above, by region: ENNS 2018, 2019, 2021




DOST-FNRI

Prevalence of chronic energy deficiency among older persons 60 years old and above, by region: Philippines, 2023

Chronic Energy Deficiency (Magnitude and Severity, 1995)*
Low: 5-9%
Low to Medium (using 95% CI)
Medium: 10-19%
Medium to High (using 95% CI)
High: 20-39%
High to Very High (using 95% CI)
Very High: ≥40%


**using 95% confidence interval estimate of CED*

Philippines: 11.7%



DOST-FNRI

Prevalence of overweight and obesity* among older persons 60 years old and above, by region: Philippines, 2023




95% Confidence interval of the national estimate

**** based on Asia Pacific BMI Classification***

DOST-FNRI

Prevalence of anemia among older people 60 years old and above, by region: Philippines, 2023



95% Confidence interval of the national estimate


DOST-FNRI

Prevalence of anemia among older persons, by region: Philippines, 2023

Public Health Significance (WHO, 2011)
Low: ≤4.9%
Low to Mild (using 95% CI)
Mild: 5-19.9%
Mild to Moderate (using 95% CI)
Moderate: 20-39.9%
Moderate to Severe (using 95% CI)
Severe: ≥40.0%

**using 95% confidence interval estimate of anemia*

Philippines: 17.2%



DOST-FNRI

2023 NNS SUMMARY

Household Food Insecurity

Highest in BARMM, Eastern Visayas, SOCCSKSARGEN, Zamboanga Peninsula, Caraga

Lowest in Central Luzon, NCR, Ilocos Region, CAR, CALABARZON

Anemia Status of Children 6-59 months

Highest in BARMM, Bicol, SOCCSKSARGEN, Central Luzon

Lowest in Ilocos Eastern Visayas, Zamboanga Peninsula

Nutritional Status of Children Under Five

UNDERWEIGHT

Highest in MIMAROPA, Zamboanga Peninsula, BARMM
Lowest in Ilocos Region, Central Luzon, CAR

STUNTING

Highest in BARMM, SOCCSKSARGEN, Zamboanga Peninsula, MIMAROPA
Lowest in Cagayan Valley, Central Luzon, Ilocos Region

WASTING

Highest in MIMAROPA, Caraga, Bicol Region
Lowest in NCR, CAR, Ilocos Region

OVERWEIGHT-FOR-HEIGHT

Highest in Central Luzon, CAR, NCR
Lowest in Western Visayas, SOCCSKSARGEN, BARMM, Eastern Visayas

DOST-FNRI

2023 NNS SUMMARY

Nutritional Status of Children, 5-10 years old

UNDERWEIGHT

Highest in MIMAROPA, BARMM, Zamboanga Peninsula
Lowest in CALABARZON, Cagayan Valley, CAR

STUNTING

Highest in BARMM, Zamboanga Peninsula, Northern Mindanao
Lowest in CALABARZON, Ilocos, Cagayan Valley

WASTING

Highest in MIMAROPA, Western Visayas, Cagayan Valley
Lowest in Davao, SOCCSKSARGEN, CAR

OVERWEIGHT/OBESITY

Highest in Ilocos, Cagayan Valley, NCR
Lowest in Zamboanga Peninsula, Northern Mindanao, BARMM

Anemia Status

SCHOOL-AGE CHILDREN

Highest in BARMM, Cagayan Valley, NCR
Lowest in Eastern Visayas, Central Visayas, Northern Mindanao

Nutritional Status of Adolescents, 10.08-19 years old

STUNTING

Highest in BARMM, SOCCSKSARGEN, Zamboanga Peninsula
Lowest in Ilocos, NCR, Central Luzon

WASTING

Highest in MIMAROPA, Western Visayas, Central Luzon
Lowest in Northern Mindanao, Davao, CAR

OVERWEIGHT/OBESITY

Highest in Central Luzon, CAR, NCR
Lowest in Bicol, BARMM, Zamboanga Peninsula

Anemia Status

ADOLESCENTS

Highest in Davao, Western Visayas, Zamboanga Peninsula
Lowest in Central Visayas, Eastern Visayas, CAR

2023 NNS SUMMARY

Nutritional Status of Adults, 20-59 years old

CHRONIC ENERGY DEFICIENCY

Highest in BARMM, Zamboanga Peninsula, Bicol
Lowest in Central Visayas, Caraga, CAR

OVERWEIGHT and OBESITY

Highest in NCR, CAR, Central Luzon
Lowest in Western Visayas, Zamboanga Peninsula, BARMM

Older Persons, 60 years old and above

CHRONIC ENERGY DEFICIENCY

Highest in Western Visayas, Zamboanga Peninsula, Cagayan Valley
Lowest in CAR, NCR, CALABARZON

OVERWEIGHT and OBESITY

Highest in NCR, Central Luzon, CALABARZON
Lowest in MIMAROPA, BARMM, Western Visayas

Anemia Status

ADULTS

Highest in Central Luzon, Cagayan Valley, Ilocos
Lowest in Central Visayas, Northern Mindanao, CAR, Eastern Visayas

OLDER PERSONS

Highest in Western Visayas, Central Luzon, Cagayan Valley
Lowest in Northern Mindanao, Central Visayas, CAR

DOST-FNRI

THANK YOU!



For more information on food and nutrition, please contact the

Department of Science and Technology
Food and Nutrition Research Institute
DOST Compound, Gen. Santos Ave., Bicutan
Taguig City, Metro Manila, Philippines
Tel nos.: (632) 8837-8113/8114 local 301;
(632) 8839-1839
Telefax No.: (632) 8837-2934
E-mail: dostfnri47@fnri.dost.gov.ph
Website: http://www.fnri.dost.gov.ph