

MA 106

Tutorial 4 Solutions

D1 T5

GYANDEV GUPTA

March 31, 2021

IIT BOMBAY



QUESTION 4.3

QUESTION 4.4

QUESTION 4.5

QUESTION 4.6

QUESTION 4.7

QUESTION 4.8



QUESTION 4.3



QUESTION 4.3

Part(i)

We have the basis set $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ of $\mathbb{R}^{3 \times 1}$ and $F = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4)$ of $\mathbb{R}^{4 \times 1}$,

$$T(\mathbf{e}_1) = \begin{bmatrix} 1 & 0 & 1 & 1 \end{bmatrix}^T = 1\mathbf{e}_1 + 0\mathbf{e}_2 + 1\mathbf{e}_3 + 1\mathbf{e}_4$$

$$T(\mathbf{e}_2) = \begin{bmatrix} 1 & 1 & 0 & 1 \end{bmatrix}^T = 1\mathbf{e}_1 + 1\mathbf{e}_2 + 0\mathbf{e}_3 + 1\mathbf{e}_4$$

$$T(\mathbf{e}_3) = \begin{bmatrix} 0 & 1 & 1 & 1 \end{bmatrix}^T = 0\mathbf{e}_1 + 1\mathbf{e}_2 + 1\mathbf{e}_3 + 1\mathbf{e}_4$$

$$M_F^E(T) = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$



QUESTION 4.3

Part(ii). Try whether the set E' and set F' forms a basis set? Indeed yes they form

$$T(\mathbf{e}_1 + \mathbf{e}_2) = \begin{bmatrix} 2 & 1 & 1 & 2 \end{bmatrix}^T =$$
$$0(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3) + 0(\mathbf{e}_2 + \mathbf{e}_3 + \mathbf{e}_4) + 1(\mathbf{e}_3 + \mathbf{e}_4 + \mathbf{e}_1) + 1(\mathbf{e}_4 + \mathbf{e}_1 + \mathbf{e}_2)$$

$$T(\mathbf{e}_2 + \mathbf{e}_3) = \begin{bmatrix} 1 & 2 & 1 & 2 \end{bmatrix}^T =$$
$$0(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3) + 1(\mathbf{e}_2 + \mathbf{e}_3 + \mathbf{e}_4) + 0(\mathbf{e}_3 + \mathbf{e}_4 + \mathbf{e}_1) + 1(\mathbf{e}_4 + \mathbf{e}_1 + \mathbf{e}_2)$$

$$T(\mathbf{e}_3 + \mathbf{e}_1) = \begin{bmatrix} 1 & 1 & 2 & 2 \end{bmatrix}^T =$$
$$0(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3) + 1(\mathbf{e}_2 + \mathbf{e}_3 + \mathbf{e}_4) + 1(\mathbf{e}_3 + \mathbf{e}_4 + \mathbf{e}_1) + 0(\mathbf{e}_4 + \mathbf{e}_1 + \mathbf{e}_2)$$

$$M_{F'}^{E'}(T) = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$



QUESTION 4.4



QUESTION 4.4

Proposition

Let $\mathbf{A}, \mathbf{B} \in \mathbb{K}^{n \times n}$. Then $\mathbf{A} \sim \mathbf{B}$ if and only if there is an ordered basis E for $\mathbb{K}^{n \times 1}$ such that \mathbf{B} is the matrix of the linear transformation $T_{\mathbf{A}} : \mathbb{K}^{n \times 1} \rightarrow \mathbb{K}^{n \times 1}$ with respect to E .

In fact, $\mathbf{B} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$ if and only if the columns of \mathbf{P} form an ordered basis, say E , for $\mathbb{K}^{n \times 1}$ and $\mathbf{B} = \mathbf{M}_E^E(T_{\mathbf{A}})$.

Using the theorem we get $E = \{\mathbf{P}_1, \mathbf{P}_2, \mathbf{P}_3, \mathbf{P}_4\}$



QUESTION 4.5



QUESTION 4.5

For $|\mathbf{A} - \mu\mathbf{I}| = 0 = (\mu - \lambda)^3$ its true for all vector $\mathbf{x} = (x_1, x_2, x_3)$ and hence eigen space is \mathbb{R}^3

For $|\mathbf{B} - \mu\mathbf{I}| = 0 = (\mu - \lambda)^3$ and for corresponding eigen vector $\mathbf{x} = (x_1, x_2, x_3)$
Solve $(\mathbf{B} - \lambda\mathbf{I})\mathbf{x} = 0 \implies x_2 = 0$ and hence eigen space is \mathbb{R}^2

For $|\mathbf{C} - \mu\mathbf{I}| = 0 = (\mu - \lambda)^3$ and for corresponding eigen vector $\mathbf{x} = (x_1, x_2, x_3)$
Solve $(\mathbf{B} - \lambda\mathbf{I})\mathbf{x} = 0 \implies x_2 = 0, x_3 = 0$ and hence eigen space is \mathbb{R}



QUESTION 4.6



QUESTION 4.6

Check $|\mathbf{A} - 3\mathbf{I}| = 0$, we get $\det \begin{bmatrix} 0 & 0 & 0 \\ -2 & 1 & 2 \\ -2 & 1 & 2 \end{bmatrix} = 0$

$$\mathbf{Ax} = 3\mathbf{x}$$

$$\begin{bmatrix} 3 & 0 & 0 \\ -2 & 4 & 2 \\ -2 & 1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 3 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

We get $-2x_1 + x_2 + 2x_3 = 0$. So all eigen vectors of form $\mathbf{x} = x_1(1, 2, 0) + x_3(0, -2, 1)$ where $x_3, x_1 \in \mathbb{R}$



QUESTION 4.6

To prove $\begin{bmatrix} 0 & 1 & 1 \end{bmatrix}^T$ is an eigenvector of \mathbf{A}

$$\begin{bmatrix} 3 & 0 & 0 \\ -2 & 4 & 2 \\ -2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 6 \\ 6 \end{bmatrix} = 6 \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

We get the eigen value to be 6.



QUESTION 4.7



QUESTION 4.7

For $|\mathbf{A} - \mu\mathbf{I}| = 0$,

$$\det\left(\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} - \mu \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

$$\det\left(\begin{bmatrix} \cos \theta - \mu & -\sin \theta \\ \sin \theta & \cos \theta - \mu \end{bmatrix}\right) = 0$$

$$\mu^2 - 2\mu \cos \theta + 1 = 0 \implies \mu = \cos \theta \pm i \sin \theta$$

Let $\mathbf{x} = (x_1, x_2)$ where $x_1, x_2 \in \mathbb{C}$

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mu \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$



QUESTION 4.7

We get $\cos \theta x_1 - \sin \theta x_2 = (\cos \theta - i \sin \theta)x_1 \implies x_2 = ix_1$

We get $\mathbf{x} = x_1(1, i)$ where $x_1 \in \mathbb{C}$

For other eigen value $\cos \theta x_1 + \sin \theta x_2 = (\cos \theta + i \sin \theta)x_1 \implies x_2 = -ix_1$

We get $\mathbf{x} = x_1(1, -i)$ where $x_1 \in \mathbb{C}$

$\mathbf{P} := \begin{bmatrix} 1 & 1 \\ i & -i \end{bmatrix}$ and Check it $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{bmatrix} \cos \theta - i \sin \theta & 0 \\ 0 & \cos \theta + i \sin \theta \end{bmatrix}$



QUESTION 4.8



QUESTION 4.8

$\text{Rank}\mathbf{A} = 1$ and $\text{Nullity}\mathbf{A} = n - 1$

Eigen vector = $\begin{bmatrix} 1 & 1 & \dots & 1 \end{bmatrix}^T$ for eigen value = n

To find $|\mathbf{A} - \mu\mathbf{I}| = 0$, Swap all rows initially ($R_i \iff R_{n+1-i} \forall i=1$ to n) and perform $R_1 \mapsto \sum_{i=1}^n R_i$ and take $(n-\mu)$ common and then $R_k \mapsto R_k - R_1 \forall k=2$ to n and then expand via last column

we get $\mu^{n-1}(\mu - n) = 0 \implies \mu = 0$ GM is $n-1$, $\mu = n$ GM is 1

Now find eigen vectors corresponding to all eigen values $(\mathbf{A} - \mu\mathbf{I})\mathbf{x} = 0$ we get

For $\mu = 0$, $\mathbf{v} = \{ \mathbf{x} : \sum_{i=1}^n x_i = 0 \}$

For $\mu = n$ we get $\mathbf{v} = x_1(1, 1, 1, 1, \dots)^T \forall x_1 \in \mathbb{R}$



QUESTION 4.8

$$\mathbf{P} := \begin{bmatrix} -1 & -1 & -1 & . & . & -1 & 1 \\ 1 & 0 & 0 & . & . & 0 & 1 \\ 0 & 1 & 0 & . & . & 0 & 1 \\ 0 & 0 & 1 & . & . & 0 & 1 \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & . \\ 0 & 0 & 0 & . & . & 1 & 1 \end{bmatrix}$$

Perform $\mathbf{P}^{-1}\mathbf{A}\mathbf{P}$ to get to a diagonal matrix $(0, 0, 0, \dots, n)^T$



QUESTIONS?

Contact me via 190100051@iitb.ac.in

THANK YOU

