

MA 108

# Tutorial 4 Solutions

D1 T5

---

GYANDEV GUPTA

June 9, 2021

IIT BOMBAY



QUESTION 4.1

QUESTION 4.7

QUESTION 4.10

QUESTION 4.11

QUESTION 4.14

QUESTION 5.1

QUESTION 5.7



---

## QUESTION 4.1



## QUESTION 4.1.vi

You can find homogeneous solution by putting  $y = x^m$

$$m^2 - 2m = 0$$

$$m_1 = 0, m_2 = 2$$

$$y_{hs} = c_1 x^{m_1} + c_2 x^{m_2} = c_1 + c_2 x^2$$

Divide the equations by  $x$ , we get  $y'' - y'/x = (x+3)x^2 e^x$

So  $r(x) = (x+3)x^2 e^x$  now what we can do ??



## QUESTION 4.1.vi

By method of variation of parameter to find particular solution

$$y_{ps} = -y_1 \int \frac{y_2 r(x)}{W(y_1, y_2)} + y_2 \int \frac{y_1 r(x)}{W(y_1, y_2)}$$

where  $y_1 = 1$ ,  $y_2 = x^2$  and  $W(y_1, y_2) = 2x$

$$y_{ps} = \frac{x^2 e^x (x^2 - 1)}{2}$$



---

## QUESTION 4.7



## QUESTION 4.7

Observation: if  $y_1, y_2$  are two solutions of a non homogeneous linear equation, then  $y_1 - y_2$  is a solution of the homogeneous part.

Applying this here we get  $y_1 - y_2, y_2 - y_3$  are solutions of the homogeneous part. One can check that these two solutions are independent (by computing the Wronskian, for instance or otherwise).

Therefore, the general solution of the given DE may be written as

$$c_1(y_1 - y_2) + c_2(y_2 - y_3) + y_1$$
$$c_1(e^{x^2}(1 - x)) + c_2(2 - e^{x^2}) + 1 + e^{x^2}$$



---

## QUESTION 4.10





## QUESTION 4.10.i

For homogenous part ( complementary fn) :  $y_{hs} = c_1 e^{ix} + c_2 e^{-ix} + c_3 x e^{ix} + c_4 x e^{-ix}$

For particular solution:(Use of annihilator I have shown you can also solve via basic too)

The annihilator for  $r(x) = \sin x$  is  $A = D^2 + 1$

$$\implies AL(y) = (D^2 + 1)(D^4 + 2D^2 + 1) = (D^2 + 1)^3$$

$$\implies D = \pm i, \text{ repeated thrice}$$



## QUESTION 4.10.i

The general solution of  $AL(y) = 0$  is

$$c_1 e^{ix} + c_2 e^{-ix} + c_3 x e^{ix} + c_4 x e^{-ix} + c_5 x^2 e^{ix} + c_6 x^2 e^{-ix}$$

Here the first four terms are annihilated, so we have a solution of the form

$$c_5 x^2 e^{ix} + c_6 x^2 e^{-ix} \text{ or } ax^2 \sin x + bx^2 \cos x$$

Plugging this in the ODE, to get the values of  $a$  and  $b$  by comparing coefficients we get  $a = -1/8$  and  $b = 0$



---

## QUESTION 4.11



## QUESTION 4.11.ii

You can find homogeneous solution by putting  $y = x^m$

$$m^4 + 2m^3 + 3m^2 + 2m + 1 = 0$$

$$m^4 + m^2 + 1 + 2m^3 + 2m^2 + 2m = 0 = (m^2 + m + 1)^2$$

$$m_1 = w, m_2 = w, m_3 = \bar{w}, m_4 = \bar{w}$$

$$y_{hs} = c_1 x^{m_1} + c_2 \ln x x^{m_2} + c_3 x^{m_3} + c_4 \ln x x^{m_4} = c_1 x^w + c_2 \ln x x^w + c_3 x^{\bar{w}} + c_4 \ln x x^{\bar{w}}$$



## QUESTION 4.11.ii

Now take the particular soln to be of form  $y_{ps} = ax^3$

Put this in ODE we get  $a=1/121$

So

$$y = c_1x^w + c_2\ln xx^w + c_3x^{\bar{w}} + c_4\ln xx^{\bar{w}} + x^3/121$$

Thanks to Vaibhav for pointing out :)



---

## QUESTION 4.14



## QUESTION 4.14

One of the way to prove, we know it would be a second order homogeneous DE in general form (GDE) as  $y'' + p(x)y' + q(x)y = 0$

Lets have  $y_1 = x^2e^x$  and  $y_2 = x^3e^x$

The wronskian for this is  $W(y_1, y_2) = x^4e^{2x}$

So  $W = e^{-\int p dx}$  we get  $p(x) = -2-4/x$

Substituting  $y_1$  or  $y_2$  in GDE we get  $q(x) = 1 + 6/x^2$

Any constant coefficient homogeneous differential equation with  $x^2e^x$  as a solution must also have  $e^x$  and  $xe^x$  as solutions.

Hence it cannot have them as solution



---

## QUESTION 5.1





## QUESTION 5.1.iv

Assume  $a$  to be *real*<sup>1</sup>

Extra info : [ The result is true for complex  $a$  but the proof given does not go through. One must use Cauchy's integral formula to deform the path of the integral into the complex domain. The result however may be used even for complex  $a$  ].

If you don't know the bracket part you can skip it not needed

Put  $t(s + a) = u$  in the defining integral,

$$\mathcal{L}(t^2 e^{-at}) = \int_0^\infty t^2 e^{-t(s+a)} dt = \frac{1}{(s+a)^3} \int_0^\infty u^2 e^{-u} du = \frac{2}{(s+a)^3}$$



More generally for complex  $a$  and  $b$  the following holds:

$$\mathcal{L}(t^a e^{-bt}) = \frac{\Gamma(a+1)}{(s+b)^{a+1}}$$

,

---

## QUESTION 5.7



## QUESTION 5.7

$$\mathcal{L}(f) = \sum_{n=1}^{\infty} \int_{n-1}^n n e^{-st} dt = \frac{e^s - 1}{s} \sum_{n=1}^{\infty} n e^{-ns} = \frac{e^s}{s(e^s - 1)}$$



QUESTIONS?

Contact me via [190100051@iitb.ac.in](mailto:190100051@iitb.ac.in)

THANK YOU

