

Principles of Quantum Mechanics – R. Shankar

Notes by Gene

2026

Contents

1	Mathematical Introduction	2
1.1	The Delta function	2
1.2	The Derivative Operator	3
1.2.1	The Hermitian Operator $K = -iD$	3
1.2.2	Eigenvalues of K	4
1.2.3	Conjugate actions of X and K	5
2	Review of Classical Mechanics	6
2.1	Lagrangian Mechanics	6
2.1.1	Procedure	7
2.1.2	The Euler-Lagrange Equation	7
2.1.3	Conservation Laws from the Lagrangian	8
2.1.4	The Electromagnetic Lagrangian	9
2.1.5	The Two-Body Problem	10
2.2	Hamiltonian Mechanics	10
2.2.1	Derivation	10
2.2.2	The Electromagnetic Hamiltonian	11
2.2.3	Conservation Laws from the Hamiltonian	12
2.2.4	Canonical Transformations	12
2.2.5	Active Transformations	13
2.2.6	Symmetries	14
2.2.7	A Relation Between S and E	14
2.3	A Problem: Wave-Particle Duality	15
2.3.1	Plane Waves	15
2.3.2	Double-Slit Experiment	15
3	Postulates of Quantum Mechanics	16
3.1	The Postulates	16
3.1.1	Ensembles	17
3.1.2	Compatible Variables	18
3.1.3	Density Matrices	19
3.2	Example: Gaussian Wavefunction	19
3.3	Generalization to N Degrees of Freedom	21
3.4	The Schrödinger Equation	21
3.4.1	Hamiltonian Setup	21

3.4.2	Time-Independent Schrödinger Equation	22
3.4.3	Time-Dependent Schrödinger Equation	22
4	Simple Problems in 1D	23
4.1	The Free Particle	23
4.1.1	Allowed Energy States	23
4.1.2	Propagator	23
4.2	The Particle in a Box	24
4.2.1	Wavefunction	24
4.2.2	Allowed Energies	25

1 Mathematical Introduction

1.1 The Delta function

In discrete space with basis $|x_i\rangle$,

- Orthogonality: $\langle x_i | x_j \rangle = \delta_{ij}$
- Completeness: $\sum_{i=1}^n |x_i\rangle \langle x_i| = I$

Natural generalization to continuous functions.

- Orthogonality: $\langle x | x' \rangle = \delta(x - x')$
- Completeness: $\int_a^b |x'\rangle \langle x| dx = I$

δ is the Dirac delta function, defined as

$$\begin{aligned} \delta(x - x') &= 0, \quad x \neq x' \\ \int_a^b \delta(x - x') dx' &= 1, \quad a < x < b \end{aligned}$$

Sometimes we write $\delta(0) = \infty$, but this isn't well defined except in the context of an integral. The δ also works like a "sampling function":

$$\int \delta(x - x') f(x') dx' = f(x)$$

Derivative: The derivative of the delta function acts like an operator

$$\delta'(x - x') = \delta(x - x') \frac{d}{dx}$$

To see this, note that

$$\int \delta'(x - x') f(x') dx' = \frac{d}{dx} \int \delta(x - x') f(x') dx' = \frac{df(x)}{dx}$$

Fourier Series: We can also represent the delta function using a fourier series. Recall that

$$\text{Fourier transform: } f(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ikx} f(x) dx$$

$$\text{Inverse transform: } f(x') = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ikx'} f(k) dk$$

$$f(x') = \int_{-\infty}^{\infty} \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ik(x'-x)} dk \right) f(x) dx$$

Reasoning from the use of delta as a sampling function,

$$\delta(x' - x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ik(x'-x)} dk$$

1.2 The Derivative Operator

We can have operators in infinite dimensions, e.g.

$$\Omega |f\rangle = |\tilde{f}\rangle$$

The derivative is one such operator:

$$D |f\rangle = \left| \frac{df}{dx} \right\rangle$$

The elements of D in the x basis are

$$D_{xx'} \equiv \langle x | D | x' \rangle = \delta'(x - x') = \delta(x - x') \frac{d}{dx'}$$

To see this, recall the use of δ as a sampling function.

$$\begin{aligned} \langle x | D | f \rangle &= \langle x | \frac{df}{dx} \rangle = \frac{df(x)}{dx} \\ \int \langle x | D | x' \rangle \langle x' | f \rangle dx' &= \frac{df}{dx} \end{aligned}$$

1.2.1 The Hermitian Operator $K = -iD$

However, D is not Hermitian: $D_{xx'} \neq D_{x'x}^*$.

$$\begin{aligned} D_{xx'} &= \delta'(x - x') \\ D_{x'x}^* &= \delta'(x' - x)^* = -\delta'(x - x') \end{aligned}$$

Instead, we can make D Hermitian using a constant complex term: $K = -iD$. Then

$$\begin{aligned} K_{xx'} &= -i\delta'(x - x') \\ K_{x'x}^* &= [-i\delta'(x' - x)]^* = i\delta'(x' - x) = -i\delta(x - x') \end{aligned}$$

Warning: For operators in infinite dimensions, $K_{xx'} = K_{x'x}^*$ doesn't actually guarantee Hermiticity. To see this, note that a Hermitian operator must satisfy (for functions f, g)

$$\begin{aligned}\langle g|K|f\rangle &= \langle f|K|g\rangle^* \\ \int_a^b \int_a^b \langle g|x\rangle \langle x|K|x'\rangle \langle x'|f\rangle dx dx' &\stackrel{?}{=} \left(\int_a^b \int_a^b \langle f|x\rangle \langle x|K|x'\rangle \langle x'|g\rangle dx dx' \right)^* \\ \int_a^b g^*(x) \left[-i \frac{df(x)}{dx} \right] dx &\stackrel{?}{=} \left(\int_a^b f^*(x) \left[-i \frac{dg(x)}{dx} \right] dx \right)^* \\ -ig^*(x)f(x) \Big|_a^b + i \int_a^b \frac{dg^*(x)}{dx} f(x) dx &\neq i \int_a^b \frac{dg^*(x)}{dx} f(x) dx\end{aligned}$$

In order to be Hermitian, the **surface term** of K must be 0. K is Hermitian in the space of functions where this holds. $K_{xx'} = K_{x'x}^*$ is no longer a sufficient condition.

$$-ig^*(x)f(x) \Big|_a^b = 0$$

Examples where

- String clamped at one end, $f(0) = f(L) = 0$. Derivative is $K = -i \frac{d}{dx}$
- Periodic function, $f(\phi) = f(\phi + 2\pi)$. Derivative is $K = -i \frac{d}{d\phi}$

In QM, we study functions $-\infty < x < \infty$. There are two classes of such functions.

1. f vanishes as $x \rightarrow -\infty$ and $x \rightarrow \infty$. Here, $K = -i \frac{d}{dx}$ is Hermitian.
2. f doesn't vanish, like e^{ikx} . If K is Hermitian, it must satisfy

$$e^{ikx} e^{ik'x} \Big|_{-\infty}^{\infty} = 0$$

If $k = k'$, this holds. Otherwise, we define the limit as $|x| \rightarrow \infty$ to be the average over a large interval. Then if $k \neq k'$,

$$\lim_{x \rightarrow \infty} e^{ikx} e^{ik'x} = \lim_{L, \Delta \rightarrow \infty} \frac{1}{\Delta} \int_L^{L+\Delta} e^{i(k-k')x} dx = 0$$

Thus K is also Hermitian for class 2 functions.

1.2.2 Eigenvalues of K

For eigenvalue k ,

$$\begin{aligned}K|k\rangle &= k|k\rangle \\ \langle x|K|k\rangle &= k\langle x|k\rangle \\ \int \langle x|K|x'\rangle \langle x'|k\rangle dx' &= k\langle x|k\rangle\end{aligned}$$

Let $\langle x|k\rangle = \psi_k(x)$. Using the definition $K = -iD$,

$$-i \frac{d}{dx} \psi_k(x) = k \psi_k(x)$$

The solution is $\psi_k(x) = A e^{ikx}$. Thus, any real number k is an eigenvalue of K with eigenvector $A e^{ikx}$. Since A is a free parameter, set it to $1/\sqrt{2\pi}$ so that

$$|k\rangle \leftrightarrow \frac{1}{\sqrt{2\pi}} e^{ikx}$$

$$\langle k|k'\rangle = \int_{-\infty}^{\infty} \langle k|x\rangle \langle x|k'\rangle dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(k'-k)x} dx = \delta(k' - k)$$

Thus $\langle k|k'\rangle = 0$ if $k \neq k'$, and blows up if $k = k'$.

We require a real eigenvalue k because if k is imaginary, e^{ikx} blows up as $x \rightarrow \infty$. The choice of a real eigenvalue restricts us to the **physical Hilbert space**.

1.2.3 Conjugate actions of X and K

In finite dimensions, the eigenfunctions of a Hermitian operator form a complete basis. Assume this holds for infinite dimensions.

K is Hermitian. Expand in both X and K basis. Recall $|k\rangle = \frac{1}{\sqrt{2\pi}} e^{ikx}$.

$$f(k) = \langle k|f\rangle = \int_{-\infty}^{\infty} \langle k|x\rangle \langle x|f\rangle dx = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ikx} f(x) dx$$

$$f(x) = \langle x|f\rangle = \int_{-\infty}^{\infty} \langle x|k\rangle \langle k|f\rangle dk = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ikx} f(k) dk$$

The Fourier transform pass f from one complete basis to another.

Additionally, note the matrix elements of X and K in the X and K basis, respectively:

$$\langle x|X|x'\rangle = x' \delta(x - x')$$

$$\langle k|K|k'\rangle = k' \delta(k - k')$$

In finite dimensions, we had (in the X basis) $X|x\rangle = x|x\rangle$. To see the action of X in infinite dimensions, let $X|f\rangle = |\tilde{f}\rangle$.

$$\langle x|X|f\rangle = \int_{-\infty}^{\infty} \langle x|X|x'\rangle \langle x'|f\rangle dx' = x f(x)$$

$$\langle x|X|f\rangle = \langle x|\tilde{f}\rangle = \tilde{f}(x)$$

Thus $\tilde{f}(x) = x f(x)$, and the effect of X in infinite dimensions is $X|f(x)\rangle = |x f(x)\rangle$. That is, multiplying $f(x)$ by x .

An interesting phenomenon occurs if we calculate the matrix elements of X in the K basis. Recall $|k\rangle = \frac{1}{\sqrt{2\pi}} e^{ikx}$.

$$\begin{aligned}\langle k|X|k'\rangle &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle k|x\rangle \langle x|X|x'\rangle \langle x'|k'\rangle dx dx' \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x' \delta(x-x') \langle k|x\rangle \langle x'|k'\rangle dx dx' \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ikx} e^{ik'x} dx \\ &= i \frac{d}{dk} \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(k'-k)x} dx \right) = i \delta'(k-k')\end{aligned}$$

Thus, if g is a function with image $g(k)$ in the K basis,

$$X |g(k)\rangle = |i \frac{dg(k)}{dk}\rangle$$

To summarize,

$$\text{In } X \text{ basis } \begin{cases} X \equiv x \\ K \equiv -i \frac{d}{dx} \end{cases}$$

$$\text{In } K \text{ basis } \begin{cases} X \equiv i \frac{d}{dx} \\ K \equiv k \end{cases}$$

Operators like these are **conjugate** to each other.

Note that X and K do not commute.

$$\begin{aligned}XK|f\rangle &= -ix \frac{df(x)}{dx} \\ KX|f\rangle &= -i \frac{d}{dx} (xf(x)) = -ix \frac{df(x)}{dx} - if(x)\end{aligned}$$

Thus $[X, K]|f\rangle = if = iI|f\rangle$. Since f was arbitrary, we have $[X, K] = iI$.

In QM, X and K are the main operators we'll be seeing (X gives position, K gives derivative). However, they take different forms: $\hat{X} = X$ and $\hat{P} = \hbar K$.

2 Review of Classical Mechanics

2.1 Lagrangian Mechanics

Newtonian mechanics can find the path of a particle at an infinitesimal time slice:

$$m\ddot{x} = -\frac{dV}{dx}$$

Given $x(t_i)$ and $\dot{x}(t_i)$ at a point t_i , we can obtain the classical path's $x_{\text{cl}}(t_i + \Delta t)$ and $\dot{x}_{\text{cl}}(t_i + \Delta t)$ after some infinitesimal time Δt using the equation above. By continuing this process, we recreate the classical path x_{cl} of the particle.

$$\begin{aligned}x_{\text{cl}}(t_i + \Delta t) &= x(t_i) + \dot{x}(t_i)\Delta t \\ \dot{x}_{\text{cl}}(t_i + \Delta t) &= \dot{x}(t_i) + \ddot{x}(t_i)\Delta t\end{aligned}$$

Lagrangian mechanics seeks to find the classical path immediately. Given two points (x_i, t_i) and (x_f, t_f) , we can find the path taken by the particle between them. But what distinguishes the classical path from all other possible ones?

2.1.1 Procedure

1. Define the Lagrangian $L = T - V$. Assume time-independent potential, thus $L = L(x, \dot{x})$.
2. For each path from (x_i, t_i) to (x_f, t_f) , define the action as a function of the entire path:

$$S[x(t)] = \int_{t_i}^{t_f} L(x, \dot{x}) dt$$

3. **Principle of least action:** the classical path is that on which S is minimized.

NB: assumption of a time-independent potential doesn't always hold. For example, $F_B = qv \times B$. So $L = T - V$ isn't always the correct form.

2.1.2 The Euler-Lagrange Equation

For finite variables, the function was at a extremal if the 1st order change δf vanishes:

$$\begin{aligned}f(\mathbf{x}^0 + \eta) &= f(\mathbf{x}^0) + \sum_{i=1}^n \left. \frac{\partial f}{\partial x_i} \right|_{\mathbf{x}^0} \eta_i + O(\eta^2) \\ \delta f &= \sum_{i=1}^n \left. \frac{\partial f}{\partial x_i} \right|_{\mathbf{x}^0} \eta_i = 0\end{aligned}$$

Assume this holds for functions of infinite variables, like $S[x(t)]$. Let $x_{\text{cl}}(t)$ be the classical path, and $x_{\text{cl}}(t) + \eta(t)$ a nearby path. Endpoints must match, so $\eta(t_i) = \eta(t_f) = 0$. Now find the extremal where 1st order change δS vanishes:

$$\begin{aligned}S[x_{\text{cl}}(t) + \eta(t)] &= \int_{t_i}^{t_f} L(x_{\text{cl}}(t) + \eta(t), \dot{x}_{\text{cl}}(t) + \dot{\eta}(t)) dt \\ &= \int_{t_i}^{t_f} \left[L(x_{\text{cl}}(t), \dot{x}_{\text{cl}}(t)) + \left. \frac{\partial L}{\partial x(t)} \right|_{x_{\text{cl}}} \cdot \eta(t) + \left. \frac{\partial L}{\partial \dot{x}(t)} \right|_{x_{\text{cl}}} \cdot \dot{\eta}(t) + \dots \right] dt \\ &\approx S[x_{\text{cl}}(t)] + \delta S\end{aligned}$$

Find where $\delta S = 0$ for an arbitrary deviation η :

$$\begin{aligned}
 0 = \delta S &= \int_{t_i}^{t_f} \left[\left. \frac{\partial L}{\partial x(t)} \right|_{x_{cl}} \cdot \eta(t) + \left. \frac{\partial L}{\partial \dot{x}(t)} \right|_{x_{cl}} \cdot \dot{\eta}(t) \right] dt \\
 &= \int_{t_i}^{t_f} \left[\left. \frac{\partial L}{\partial x(t)} \right|_{x_{cl}} \cdot \eta(t) \right] dt + \left[\left. \frac{\partial L}{\partial \dot{x}(t)} \right|_{x_{cl}} \cdot \eta(t) \right]_{t_i}^{t_f} - \int_{t_i}^{t_f} \left[\left. \frac{d}{dt} \frac{\partial L}{\partial \dot{x}(t)} \right|_{x_{cl}} \cdot \eta(t) \right] dt \\
 &= \int_{t_i}^{t_f} \left[\left. \frac{\partial L}{\partial x(t)} - \frac{d}{dt} \frac{\partial L}{\partial \dot{x}(t)} \right|_{x_{cl}} \right] \cdot \eta(t) dt
 \end{aligned}$$

Thus we get the Euler-Lagrange equation, which must be satisfied by a classical path:

$$\left[\left. \frac{\partial L}{\partial x(t)} - \frac{d}{dt} \frac{\partial L}{\partial \dot{x}(t)} \right|_{x_{cl}} \right] = 0$$

2.1.3 Conservation Laws from the Lagrangian

A benefit of the Lagrangian is that unlike the Newtonian, coordinates don't have to be Cartesian. We can have an arbitrary q_i and $\dot{q}_i$.

Further note:

- Canonical momentum conjugate: $p_i = \partial L / \partial \dot{q}_i$
- Canonical force conjugate: $F_i = \partial L / \partial q_i$

$$F_i = \frac{d}{dt} p_i$$

E.g. if $q_i = \theta$, then p_i is angular momentum and F_i is torque.

Call q_i a **cyclic coordinate** if L depends on $\dot{q}_i$, but not q_i . Then p_i is conserved:

$$\frac{dp_i}{dt} = \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_i} \right) = \frac{\partial L}{\partial q_i} = 0$$

Conservation laws fall out naturally out of the Lagrangian, while it may be opaque in the Newtonian. For example, consider a Lagrangian in polar coordinates $L = L(\rho, \phi)$. The metric is $ds^2 = d\rho^2 + \rho d\phi^2$. Thus the infinitesimal velocity is

$$v = \frac{ds}{dt} = \sqrt{(\dot{\rho})^2 + (\rho \dot{\phi})^2}$$

The Lagrangian is

$$L = \frac{1}{2} m [\dot{\rho}^2 + \rho^2 \dot{\phi}^2] - V(\rho, \phi)$$

Applying Euler-Lagrange for ρ and ϕ respectively gives

$$\begin{aligned}
 \frac{d}{dt} (m\dot{\rho}) &= -\frac{\partial V}{\partial \rho} + m\rho\dot{\phi}^2 \quad \longrightarrow \quad m\ddot{\rho} = -\frac{\partial V}{\partial \rho} + m\rho(\dot{\phi})^2 \\
 \frac{d}{dt} (m\rho^2\dot{\phi}) &= -\frac{\partial V}{\partial \phi} \quad \longrightarrow \quad m\ddot{\phi} = -\frac{1}{\rho^2} \frac{\partial V}{\partial \phi} - \frac{1}{\rho} (2m\dot{\rho}\dot{\phi})
 \end{aligned}$$

Note how easily we can derive the centrifugal and coriolis forces. Also note that conservation laws for cyclic coordinates $V(\rho, \phi) = V(\rho)$ and $V(\rho, \phi) = V(\phi)$ fall out naturally, and the corresponding equations of motion are that of $\partial V/\partial \phi = 0$ and $\partial V/\partial \rho = 0$.

In sum:

- Equations of motion are simpler to find with Lagrangian
- Conservation laws are easy to see from Lagrangian
- Path integral formalism uses Lagrangian (later)

2.1.4 The Electromagnetic Lagrangian

Electromagnetic force:

$$\vec{F} = q \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right)$$

Note: this is Gaussian units, hence the $1/c$ factor for magnetic force.

The electromagnetic force is velocity-dependent, so $L = T - V$ doesn't work. Instead, we construct L to reduce to the expected equations. The EM lagrangian is

$$L_{\text{EM}} = \frac{1}{2} m \vec{v} \cdot \vec{v} - q\phi + \frac{q}{c} \vec{v} \cdot \vec{A}$$

ϕ and $\vec{A}$ are the scalar and vector potentials, related to E and B by

$$\begin{aligned} \vec{E} &= -\nabla\phi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t} \\ \vec{B} &= \nabla \times \vec{A} \end{aligned}$$

The Euler-Lagrange equation is

$$\begin{aligned} \frac{d}{dt} \left(m\dot{x}_i + \frac{q}{c} A_i \right) &= -q \frac{\partial \phi}{\partial x_i} + \frac{q}{c} \frac{\partial (\vec{v} \cdot \vec{A})}{\partial x_i}, \quad i = 1, 2, 3 \\ \frac{d}{dt} \left(m\vec{v} + \frac{q}{c} \vec{A} \right) &= -q \nabla \phi + \frac{q}{c} \nabla (\vec{v} \cdot \vec{A}) \\ \frac{d}{dt} (m\vec{v}) &= -q \nabla \phi + \frac{q}{c} \left[-\frac{d\vec{A}}{dt} + \nabla (\vec{v} \cdot \vec{A}) \right] \end{aligned} \tag{1}$$

By time-dependence of the magnetic vector potential $\vec{A}$,

$$\frac{d\vec{A}}{dt} = \frac{\partial \vec{A}}{\partial t} + \sum_i \frac{\partial \vec{A}}{\partial x_i} \frac{dx_i}{dt} = \frac{\partial \vec{A}}{\partial t} + (\vec{v} \cdot \nabla) \vec{A}$$

Thus equation (1) becomes

$$\frac{d}{dt} (m\vec{v}) = -q \nabla \phi - \frac{q}{c} \frac{\partial \vec{A}}{\partial t} + \frac{q}{c} \underbrace{\left[\nabla (\vec{v} \cdot \vec{A}) - (\vec{v} \cdot \nabla) \vec{A} \right]}_{\vec{v} \times (\nabla \times \vec{A})} \tag{2}$$

This reduces to $\vec{F} = q\vec{E} + \frac{q}{c}\vec{v} \times \vec{B}$.

The time-dependent EM field is usually not conservative, so there's no path-independent work function to use as the potential. Thus L_{EM} is made to fit the situation and reduce as expected. $L = T - V$ only works for conservative forces.

2.1.5 The Two-Body Problem

Assume 2 masses m_1, m_2 that exert equal and opposite forces on each other. We can reduce this to a one-body problem with Lagrangian mechanics.

The potential only depends on relative position, $V(\vec{r}_1, \vec{r}_2) = V(\vec{r}_1 - \vec{r}_2)$. Thus $6 \rightarrow 3$ cyclic coordinates in 3D.

The two-body Lagrangian is

$$L = \frac{1}{2}m_1|\dot{\vec{r}}_1|^2 + \frac{1}{2}m_2|\dot{\vec{r}}_2|^2 - V(r_1 - r_2) \quad (1)$$

Define $\vec{r}_{\text{CM}}$ and $\vec{r}$ such that

$$\begin{aligned} \vec{r}_{\text{CM}} &= \frac{m_1\vec{r}_1 + m_2\vec{r}_2}{m_1 + m_2}, \quad \vec{r} = \vec{r}_1 - \vec{r}_2 \\ \vec{r}_1 &= \vec{r}_{\text{CM}} + \frac{m_2\vec{r}}{m_1 + m_2} \\ \vec{r}_2 &= \vec{r}_{\text{CM}} - \frac{m_1\vec{r}}{m_1 + m_2} \end{aligned}$$

The two-body Lagrangian becomes

$$L = \frac{1}{2}(m_1 + m_2)|\dot{\vec{r}}_{\text{CM}}|^2 + \frac{1}{2}\frac{m_1m_2}{m_1 + m_2}|\dot{\vec{r}}|^2 - V(\vec{r}) \quad (2)$$

This can be thought of as the Lagrangian of two fictitious particles that don't interact: one "CM" particle with mass $M = m_1 + m_2$, and one "reduced mass" particle with mass $\mu = (m_1m_2)/(m_1 + m_2)$. Note that $L(r, \dot{r}, r_{\text{CM}}, \dot{r}_{\text{CM}}) = L(r, \dot{r}) + L(r_{\text{CM}}, \dot{r}_{\text{CM}})$.

By going to the CM frame, we can eliminate $\dot{r}_{\text{CM}}$ and just solve for the motion of μ . We've reduced the two-body problem to a one-body problem.

2.2 Hamiltonian Mechanics

2.2.1 Derivation

The Lagrangian uses $L(q, \dot{q})$. The Hamiltonian reverses the role of p and $\dot{q}$, using $H(q, p)$. This "role reversal" is achieved with a **Legendre transform**. Suppose $u(x) = df/dx$ and $x(u) = dg/du$. The relationship between f and g is a Legendre transform:

$$g(u) = x(u)u - f(x(u))$$

Define the Hamiltonian $H(q, p)$ as a Legendre transform of the Lagrangian $L(q, \dot{q})$:

$$H(q, p) = \sum_{i=1}^n p_i \dot{q}_i - L(q, \dot{q}) \quad , \quad p_i = \frac{\partial L}{\partial \dot{q}_i}$$

Hamilton's **canonical equations** can be found:

$$\dot{q}_i = \frac{\partial H}{\partial p_i} \quad , \quad \dot{p}_i = -\frac{\partial H}{\partial q_i}$$

Lagrangian	Hamiltonian
n coordinates q_i and n velocities $\dot{q}_i$	n coordinates q_i and n momenta p_i
State of system is point in n -dimensional configuration space	State of system is point in $2n$ -dimensional configuration space
n coordinates evolve according to n 2nd-order differential equations	$2n$ coordinates and momenta evolve according to $2n$ 1st-order differential equations
For given L , multiple trajectories can pass through a point in configuration space depending on $\dot{q}_i$.	For given H , only one trajectory passes through a point in configuration space.

For conservative forces with $T = \frac{1}{2}m\dot{x}_i^2$, we get

$$p_i = \frac{\partial L}{\partial \dot{x}_i} = m_i \dot{x}_i$$

$$H = \sum_i p_i \dot{x}_i - L(x, \dot{x}) = T + V$$

But this isn't always the right form. It's only for conservative forces.

2.2.2 The Electromagnetic Hamiltonian

If L_{EM} generates the correct equations of motion, then H_{EM} will as well. But it won't take the form $H = T + V$ since now the force isn't conservative.

Start from the electromagnetic Lagrangian:

$$L_{\text{EM}} = \frac{1}{2}m\vec{v} \cdot \vec{v} - q\phi + \frac{q}{c}\vec{v} \cdot \vec{A}$$

Now find the canonical momentum $\vec{p}$ and the Hamiltonian H :

$$\vec{p} = \frac{\partial L}{\partial \vec{v}} = m\vec{v} + \frac{q}{c}\vec{A}$$

$$H_{\text{EM}} = \vec{p} \cdot \vec{v} - L_{\text{EM}} = \frac{1}{2}m\vec{v} \cdot \vec{v} + q\phi = \frac{|\vec{p} - q\vec{A}/c|^2}{2m} + q\phi$$

2.2.3 Conservation Laws from the Hamiltonian

If q_i is a cyclic coordinate in the Hamiltonian, then

$$\dot{p}_i = -\frac{\partial H}{\partial q_i} = 0$$

The associated momentum is conserved. But other quantities are too, and this is often easier to see in the Hamiltonian than in the Lagrangian.

Let $\omega(p, q)$ be a state function with **no explicit dependence on t** . Define the **Poisson bracket**

$$\{\omega, \lambda\} = \sum_i \left(\frac{\partial \omega}{\partial q_i} \frac{\partial \lambda}{\partial p_i} - \frac{\partial \omega}{\partial p_i} \frac{\partial \lambda}{\partial q_i} \right)$$

The time derivative of ω is:

$$\begin{aligned} \frac{d\omega}{dt} &= \sum_i \left(\frac{\partial \omega}{\partial q_i} \frac{\partial q_i}{\partial t} + \frac{\partial \omega}{\partial p_i} \frac{\partial p_i}{\partial t} \right) \\ &= \sum_i \left(\frac{\partial \omega}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \omega}{\partial p_i} \frac{\partial H}{\partial q_i} \right) \\ &= \{\omega, H\} \end{aligned}$$

Any state function ω whose Poisson bracket with H vanishes is conserved. H itself is conserved if there's no explicit t dependence.

The Poisson bracket between q and p satisfy

$$\begin{aligned} \{q_i, q_j\} &= \{p_i, p_j\} = 0 \\ \{q_i, p_j\} &= \delta_{ij} \end{aligned}$$

From these, the canonical equations can be derived in terms of Poisson brackets:

$$\begin{aligned} \dot{q}_i &= \{q_i, H\} \\ \dot{p}_i &= \{p_i, H\} \end{aligned}$$

2.2.4 Canonical Transformations

Euler-Lagrange equations are invariant under a coordinate change $(q, \dot{q}) \rightarrow (\bar{q}, \dot{\bar{q}})$, which implies the invariance of Hamilton's equations under $(q, p) \rightarrow (\bar{q}, \bar{p})$.

General transformations of the form $q \rightarrow \bar{q}(q, p)$ and $p \rightarrow \bar{p}(q, p)$ are called **canonical transformations** if they obey the canonical equations. $(\bar{q}, \bar{p})$ are called **canonical coordinates**.

Theorem: If $(\bar{q}(q, p), \bar{p}(q, p))$ are canonical coordinates iff.

$$\begin{aligned} \{\bar{q}_j, \bar{q}_k\} &= 0 = \{\bar{p}_j, \bar{p}_k\} \\ \{\bar{q}_j, \bar{p}_k\} &= \delta_{jk} \end{aligned}$$

Proof: For state functions $\omega(q, p)$,

$$\dot{\omega} = \{\omega, H\} = \sum_i \left(\frac{\partial \omega}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \omega}{\partial p_i} \frac{\partial H}{\partial q_i} \right)$$

Thus for $\dot{q}_j$,

$$\dot{q}_j = \sum_i \left(\frac{\partial \bar{q}_j}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \bar{q}_j}{\partial p_i} \frac{\partial H}{\partial q_i} \right) \quad (1)$$

Using the chain rule, we can also get

$$\begin{cases} \frac{\partial H(q, p)}{\partial p_i} = \frac{\partial H(\bar{q}, \bar{p})}{\partial p_i} = \sum_k \left(\frac{\partial H}{\partial \bar{q}_k} \frac{\partial \bar{q}_k}{\partial p_i} + \frac{\partial H}{\partial \bar{p}_k} \frac{\partial \bar{p}_k}{\partial p_i} \right) \\ \frac{\partial H(q, p)}{\partial q_i} = \frac{\partial H(\bar{q}, \bar{p})}{\partial q_i} = \sum_k \left(\frac{\partial H}{\partial \bar{q}_k} \frac{\partial \bar{q}_k}{\partial q_i} + \frac{\partial H}{\partial \bar{p}_k} \frac{\partial \bar{p}_k}{\partial q_i} \right) \end{cases} \quad (2)$$

Plugging (2) into (1) and simplifying gives (following a similar procedure for $\dot{p}_j$),

$$\begin{cases} \dot{q}_j = \sum_k \left(\frac{\partial H}{\partial \bar{q}_k} \{\bar{q}_j, \bar{q}_k\} + \frac{\partial H}{\partial \bar{p}_k} \{\bar{q}_j, \bar{p}_k\} \right) \\ \dot{p}_j = \sum_k \left(\frac{\partial H}{\partial \bar{q}_k} \{\bar{p}_j, \bar{q}_k\} + \frac{\partial H}{\partial \bar{p}_k} \{\bar{p}_j, \bar{p}_k\} \right) \end{cases} \quad (3)$$

For (3) to reduce to the canonical equations, we must have $\{\bar{q}_j, \bar{q}_k\} = 0 = \{\bar{p}_j, \bar{p}_k\}$ and $\{\bar{q}_j, \bar{p}_k\} = \delta_{jk}$.

*Note that the Poisson bracket is independent of parametrization. Whether $\omega(q, p)$ or $\omega(\bar{q}, \bar{p})$,

$$\{\omega, \lambda\}_{q, p} = \{\omega, \lambda\}_{\bar{q}, \bar{p}}$$

2.2.5 Active Transformations

The transformation $q \rightarrow \bar{q}(q, p)$ and $p \rightarrow \bar{p}(q, p)$ is passive if (q, p) and $(\bar{q}, \bar{p})$ refer to the same point in phase space.

If the transformation $(q, p) \rightarrow (\bar{q}, \bar{p})$ preserves the range of variables, it's called **regular**.

- E.g. transformation from a Cartesian plane to a translated/rotated one is regular.
- But a transformation from a Cartesian plane to spherical coordinates isn't. Some variables are now bounded by 2π instead of ∞ .

A transformation that “shifts” the phase space so that (q, p) and $(\bar{q}, \bar{p})$ no longer refer to the same point is called **active**. Active transformations are a subset of regular ones.

We call ω invariant under the regular transformation $(q, p) \rightarrow (\bar{q}, \bar{p})$ if

$$\omega(q, p) = \omega(\bar{q}, \bar{p})$$

2.2.6 Symmetries

Two results:

1. If H is invariant under the following infinitesimal transformations, then $g(q, p)$ is conserved. We call g a **generator of the transformation**.

$$\begin{aligned} q_i &\rightarrow \bar{q}_i + \varepsilon \frac{\partial g}{\partial p_i} \equiv q_i + \delta q_i \\ p_i &\rightarrow \bar{p}_i - \varepsilon \frac{\partial g}{\partial q_i} \equiv p_i + \delta p_i \end{aligned}$$

2. If H is invariant under the regular, canonical, but not necessarily infinitesimal transformation $(q, p) \rightarrow (\bar{q}, \bar{p})$, and if (q, p) is a solution to the equations of motion, so is $(\bar{q}, \bar{p})$.

Proof of (1): Note that

$$\begin{aligned} \delta H &= \sum_i \frac{\partial H}{\partial q_i} (\delta q_i) + \frac{\partial H}{\partial p_i} (\delta p_i) \\ &= \sum_i \frac{\partial H}{\partial q_i} \left(\varepsilon \frac{\partial g}{\partial p_i} \right) + \frac{\partial H}{\partial p_i} \left(-\varepsilon \frac{\partial g}{\partial q_i} \right) \\ &= \varepsilon \{H, g\} \end{aligned}$$

For H to be invariant (to first order), $\delta H = 0$ and $\{H, g\} = 0$. Thus g is conserved.

In general for state variables ω ,

$$\delta \omega = \varepsilon \{\omega, g\}$$

2.2.7 A Relation Between S and E

Let x_{cl} be a classical path from (x_i, t_i) to (x_f, t_f) , $H(t_f)$ be the Hamiltonian at t_f , and S the classical action. A neat relation is

$$\frac{\partial S_{\text{cl}}(x_f, t_f; x_i, t_i)}{\partial t_f} = -H(t_f)$$

If E is conserved, this becomes

$$\frac{\partial S_{\text{cl}}(x_f, t_f; x_i, t_f)}{\partial t_f} = -E$$

Proof: Hold x_i, x_f fixed and vary with $\eta(t)$. For convenience, $x_i = t_i = 0$ and $\eta(0) = 0$.

$$x(t) = x_{\text{cl}}(t) + \eta(t)$$

The variation in action can be split into (1) the contribution from difference in path, (2) the contribution from the new “end” time $t = t_f + \Delta t$.

$$\begin{aligned}
 \delta S_{\text{cl}} &= \int_0^{t_f} \frac{\partial L}{\partial t} dt + L(t_f) \Delta t \\
 &= \int_0^{t_f} \left[\frac{\partial L}{\partial x} \eta(t) + \frac{\partial L}{\partial \dot{x}} \dot{\eta}(t) \right] dt + L(t_f) \Delta t \\
 &= \int_0^{t_f} \left[-\frac{d}{dt} \frac{\partial L}{\partial \dot{x}} + \frac{\partial L}{\partial x} \right] \eta(t) dt + \int_0^{t_f} \frac{d}{dt} \left[\frac{\partial L}{\partial \dot{x}} \eta(t) \right] dt + L(t_f) \Delta t \\
 &= 0 + \left[\frac{\partial L}{\partial \dot{x}} \eta(t) \right]_0^{t_f} + L(t_f) \Delta t
 \end{aligned}$$

Substitute $\eta(t_f) = -\dot{x}(t_f) \Delta t$ to get

$$\delta S = \left[-\frac{\partial L}{\partial \dot{x}} \dot{x} + L \right]_{t_f} \Delta t = -H(t_f) \Delta t$$

2.3 A Problem: Wave-Particle Duality

2.3.1 Plane Waves

An electromagnetic wave ψ that travels at c obey the wave equation

$$\nabla^2 \psi = \frac{1}{c^2} \frac{\partial^2 \psi}{\partial t^2}$$

A special case is the plane wave:

$$\psi(x, t) = A e^{i\phi} = A \exp \left(i \left(\frac{2\pi}{\lambda} x - \frac{2\pi}{T} t \right) \right)$$

- Wave number $k = 2\pi/\lambda$ (phase change per unit length at fixed t)
- Angular frequency $\omega = 2\pi/T$ (phase change per unit time at fixed x)
- Wave speed $v = \omega/T$

This can be extended to plane waves in 3D, using the **wave vector** $\vec{k}$.

$$\psi(\vec{r}, t) = A e^{i(\vec{k} \cdot \vec{r} - \omega t)} \quad , \quad \omega = |\vec{k}|v$$

2.3.2 Double-Slit Experiment

The wave prediction is $I_{1+2} \neq I_1 + I_2$, due to interference effects.

The classical particle prediction is $I_{1+2} = I_1 + I_2$.

By opening one slit and lowering the intensity, we find that light energy comes in bundles. Each bundle has energy and momentum that satisfies

$$E = \hbar\omega \quad , \quad p = \hbar k$$

Thus light seems to be made up of particles called photons. But if both slits are opened at I , we get a wave effect: $I_{1+2} \neq I_1 + I_2$. The difference can't be reconciled classically.

Born suggested that each photon is associated with a wave ψ called the **probability amplitude**. $|\psi(x)|^2$ is the probability of finding the particle at x . The wave is described by

$$\omega = \frac{E}{\hbar} \quad , \quad k = \frac{p}{\hbar}$$

The wave interferes with itself (after being split by the double-slit) to form the overall pattern according to $|\psi(x)|^2$.

De Broglie suggested that every particle of momentum p has a wave associated with it, not just massless photons. The wave number is

$$k = \frac{p}{\hbar}$$

3 Postulates of Quantum Mechanics

3.1 The Postulates

Classical:

1. State of particle is described by $x(t)$ and $p(t)$ in 2D phase space.
2. Every dynamical variable $\omega(x, p)$ is a function of x, p .
3. If a particle is in the state $(x; p)$, a perfect measurement of ω yields $\omega(x, p)$ without changing the state.
4. Time evolution described by

$$\dot{x} = \frac{\partial H}{\partial p} \quad , \quad \dot{p} = -\frac{\partial H}{\partial x}$$

Quantum:

1. State of particle is a vector $|\psi(t)\rangle$ in Hilbert space.
2. Independent variables x, p of classical mechanics become Hermitian operators X, P such that

$$\begin{aligned} \langle x|X|x'\rangle &= x\delta(x-x') \\ \langle x|P|x'\rangle &= -\hbar\delta'(x-x') \end{aligned}$$

(Note that this is just X and $P = \hbar K$ from before). Then $\omega(x, p)$ becomes

$$\Omega(X, P) = \omega(x \rightarrow X, p \rightarrow P)$$

3. If a particle is in the state $|\psi\rangle$, a perfect quantum measurement of Ω yields an eigenvalue ω with probability

$$P(\omega) \propto |\langle \omega|\psi\rangle|^2$$

The state of the system changes:

$$|\psi\rangle \rightarrow |\omega\rangle$$

Note that Ω is Hermitian, so the eigenvalues are real. In terms of the projection operator $\mathbb{P}_\omega = |\omega\rangle\langle\omega|$,

$$\begin{aligned} P(\omega) &\propto |\langle\omega|\psi\rangle|^2 = \langle\psi|\omega\rangle\langle\omega|\psi\rangle \\ &= \langle\psi|\mathbb{P}_\omega|\psi\rangle \\ &= \langle\psi|\mathbb{P}_\omega\mathbb{P}_\omega|\psi\rangle \\ &= \langle\mathbb{P}_\omega\psi|\mathbb{P}_\omega\psi\rangle \end{aligned}$$

4. Time evolution described by

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H |\psi(t)\rangle$$

$H(X, P) = H(x \rightarrow X, p \rightarrow P)$ from the classical Hamiltonian.

Potential problems:

1. $\Omega(x \rightarrow X, p \rightarrow P)$ is ambiguous. E.g. for $\omega = xp$, is $\Omega = XP$ or PX ? Solution is to symmetrize with $\Omega = (XP + PX)/2$.
2. Ω can be degenerate, e.g. two eigenvalues $\omega_1 = \omega_2 = \omega$. Then what is $P(\omega)$? Solution: find some basis $|\omega_1\rangle, |\omega_2\rangle$ in the eigenspace $\mathbb{V}_\omega$. Then

$$P(\omega) = |\langle\omega_1|\psi\rangle|^2 + |\langle\omega_2|\psi\rangle|^2$$

This is the same result we get by assuming ω_1, ω_2 are infinitesimally distinct, then asking for $P(\omega_1 \text{ or } \omega_2)$.

3. Eigenvalue spectrum of Ω can be continuous, e.g. X . Solution: expand

$$|\psi\rangle = \int |\omega\rangle \langle\omega|\psi\rangle d\omega$$

Introduce the **wave function in ω space** $\psi(\omega) = \langle\omega|\psi\rangle$ and the **probability density** $P(\omega) = |\langle\omega|\psi\rangle|^2$. $P(\omega) d\omega$ is the probability of measuring between ω and $\omega + d\omega$. Note $\int P(\omega) d\omega = 1$.

3.1.1 Ensembles

Used to test quantum theory. A quantum ensemble is a large number N of particles all in state $|\omega\rangle$.

A single particle, after measurement, could give any of the eigenvalues. Measure all N to get a statistical result.

Difference with classical:

- In classical ensemble of N particles, if we measure λ_1 in $N/3$ trials and λ_2 in $2N/3$ trials, we expect there were $N/3$ amount of λ_1 and $2N/3$ amount of λ_2 before measurement.
 - In quantum ensemble, all particles are assumed to be $|\omega\rangle$ before measurement.
-

To calculate average over ensemble, use the **expectation value**:

$$\begin{aligned}
\langle \Omega \rangle &= \sum_i P(\omega_i) \omega_i = \sum_i |\langle \omega_i | \psi \rangle|^2 \omega_i \\
&= \sum_i \langle \psi | \omega_i \rangle \langle \omega_i | \psi \rangle \omega_i \\
&= \sum_i \langle \psi | \Omega | \omega_i \rangle \langle \omega_i | \psi \rangle \\
&= \langle \psi | \Omega | \psi \rangle
\end{aligned}$$

To measure the uncertainty (fluctuation around the mean), use the **standard deviation**:

$$\Delta \Omega = \langle (\Omega - \langle \Omega \rangle)^2 \rangle^{1/2}$$

Classically, this is called “root-mean-square” deviation. In QM, it’s “uncertainty in Ω .”

$$(\Delta \Omega)^2 = \begin{cases} \sum_i P(\omega_i) (\omega_i - \langle \Omega \rangle)^2 & \text{(discrete spectrum)} \\ \int P(\omega) (\omega - \langle \Omega \rangle)^2 d\omega & \text{(continuous spectrum)} \end{cases}$$

E.g. if a particle has $\langle X \rangle = a$ and $\Delta X = \Delta$, then we know it’s likely to be spotted near a with deviation of order Δ .

3.1.2 Compatible Variables

Is there a filtering process by which we take $|\psi\rangle$ and produce a state with well-defined ω , λ for 2 variables Ω , Λ ?

- Not generally, because measurement changes the state. I.e. (if we do Ω then Λ)

$$\begin{aligned}
\text{first: } |\psi\rangle &\rightarrow \sum |\omega\rangle \langle \omega | \psi \rangle \\
\text{second: } |\omega\rangle &\rightarrow \sum |\lambda\rangle \langle \lambda | \omega \rangle
\end{aligned}$$

But if Ω and Λ have simultaneous eigenkets, then

$$\begin{aligned}
\Omega |\omega\lambda\rangle &= \omega |\omega\lambda\rangle \\
\Lambda |\omega\lambda\rangle &= \lambda |\omega\lambda\rangle \\
(\Omega\Lambda - \Lambda\Omega) |\omega\lambda\rangle &= 0
\end{aligned}$$

We can “know” both variables at the same time, if it has a simultaneous eigenket. Variables such that $[\Omega, \Lambda] = 0$ are called **compatible**.

E.g. The canonical commutation relation $[X, P] = i\hbar \neq 0$. Thus, a state can’t have well-defined eigenkets of X and P simultaneously. This is the Heisenberg uncertainty relation.

3.1.3 Density Matrices

Instead of an ensemble of N particles all in $|\psi\rangle$, we can consider an ensemble of N systems, n_i of which are $|i\rangle$.

Let $p_i = n_i/N$ be the probability of measuring a $|i\rangle$. Define the density matrix:

$$\rho = \sum_i p_i |i\rangle \langle i|$$

- “Pure”: all $n_i = 0$ except for one
- “Mixed”: otherwise.

The ensemble average of Ω is $\langle \bar{\Omega} \rangle = \sum_i p_i \langle i | \Omega | i \rangle$. Note that

$$\begin{aligned} \text{Tr}(\Omega \rho) &= \sum_j \langle j | \Omega \rho | j \rangle = \sum_i \sum_j \langle j | \Omega \rho | i \rangle \langle i | j \rangle p_i \\ &= \sum_i \sum_j \langle i | j \rangle \langle j | \Omega \rho | i \rangle p_i = \sum_i \langle i | \Omega \rho | i \rangle p_i \\ &= \langle \bar{\Omega} \rangle \end{aligned}$$

E.g. if we want the probability of obtaining ω , $\overline{P(\omega)}$, then

$$\begin{aligned} P(\omega) &= |\langle \omega | \psi \rangle|^2 = \langle \psi | \omega \rangle \langle \omega | \psi \rangle = \langle \psi | \mathbb{P}_\omega | \psi \rangle \\ \overline{P(\omega)} &= \text{Tr}(\mathbb{P}_\omega \rho) \end{aligned}$$

Features:

- $\rho^\dagger = \rho$
- $\rho^2 = \rho$ for pure state
- $\rho = \frac{1}{k} I$ for mixed state uniformly distributed over k sub-states
- $\text{Tr}(\rho) = 1$
- $\text{Tr}(\rho^2) \leq 1$, equality if ρ is pure.

3.2 Example: Gaussian Wavefunction

We'll find the Heisenberg uncertainty relation (sort of) for this model.

Put the wavefunction in the X -basis:

$$|\psi\rangle = \int_{-\infty}^{\infty} |x\rangle \langle x | \psi \rangle dx = \int_{-\infty}^{\infty} |x\rangle \psi(x) dx$$

Assume $\psi(x)$ is Gaussian and normalize to unity:

$$\psi(x) = \frac{1}{(\pi \Delta^2)^{1/4}} e^{-(x-a)^2/(2\Delta^2)}$$

In the X -basis, the effect of X is to multiply by x . I.e.

$$\begin{aligned}\langle x|X|\psi\rangle &= \int_{-\infty}^{\infty} \langle x|X|x'\rangle \langle x'|\psi\rangle dx' = \int_{-\infty}^{\infty} x\delta(x-x')\psi(x') dx' \\ &= x\psi(x)\end{aligned}$$

Expectation value is $\langle X \rangle = a$ (which makes sense; $\psi(x)$ is centered around a).

Also, $\langle X^2 \rangle = \Delta^2/2 + a^2$

Then the uncertainty is

$$\begin{aligned}\Delta X &= \left[\langle \psi | (X - \langle X \rangle)^2 | \psi \rangle \right]^{1/2} = \left[\langle X^2 \rangle - 2a\langle X \rangle + a^2 \right]^{1/2} \\ &= \left[\langle X^2 \rangle - a^2 \right]^{1/2} \\ &= \frac{\Delta}{\sqrt{2}}\end{aligned}$$

Now for P . The action in the X basis is $-i\hbar \frac{d}{dx}$. I.e.

$$\begin{aligned}\langle x|P|\psi\rangle &= \int_{-\infty}^{\infty} \langle x|P|x'\rangle \langle x'|\psi\rangle dx' = \int_{-\infty}^{\infty} -i\hbar\delta'(x-x')\psi(x') dx' \\ &= -i\hbar \frac{d\psi}{dx}\end{aligned}$$

Now find the eigenvalues. Note that $\langle x|p\rangle = \psi_p(x) \neq \psi(x)$.

$$\begin{aligned}P|p\rangle &= p|p\rangle \longrightarrow \langle x|P|p\rangle = p\langle x|p\rangle \\ -i\hbar \frac{d\psi_p(x)}{dx} &= p\psi_p(x) \\ \psi_p(x) &= \frac{1}{\sqrt{2\pi\hbar}} e^{ipx/\hbar}\end{aligned}$$

By finding $\psi(p) = \langle p|\psi\rangle$, we get a Gaussian of width $\hbar/(\Delta\sqrt{2})$ centered at 0. Thus $\langle P \rangle = 0$ and $\Delta P = \hbar/(\Delta\sqrt{2})$.

Put together: for this Gaussian wavefunction model, we get

$$\Delta X \cdot \Delta P = \frac{\hbar}{2}$$

This is a lower bound on the Heisenberg uncertainty relation.

3.3 Generalization to N Degrees of Freedom

Use mutually commuting operators $X_1, \dots, X_N$. In their simultaneous eigenbasis $|x_1, x_2, \dots, x_N\rangle$,

$$\langle x_1, x_2, \dots, x_N | x'_1, x'_2, \dots, x'_N \rangle = \delta(x_1 - x'_1) \cdots \delta(x_N - x'_N)$$

And

$$\begin{aligned} |\psi\rangle &\longrightarrow \langle x_1, \dots, x_N | \psi \rangle = \psi(x_1, \dots, x_N) \\ X_i |\psi\rangle &\longrightarrow \langle x_1, \dots, x_N | X_i | \psi \rangle = x_i \psi(x_1, \dots, x_N) \\ P_i |\psi\rangle &\longrightarrow \langle x_1, \dots, x_N | P_i | \psi \rangle = -i\hbar \frac{\partial}{\partial x_i} \psi(x_1, \dots, x_N) \end{aligned}$$

Then follow the procedure of replacing classical variables with quantum operators. E.g. for the classical operator

$$\omega = \frac{p_x^2 + p_y^2 + p_z^2}{2m} + x^2 + y^2 + z^2$$

We associate the quantum operator such that (with $X_1 = X$, $X_2 = Y$, and $X_3 = Z$)

$$\Omega |\omega\rangle = \omega |\omega\rangle \quad \Omega = \frac{P_x^2 + P_y^2 + P_z^2}{2m} + X^2 + Y^2 + Z^2$$

Then substitute $|\omega\rangle = \psi_\omega(x, y, z)$, $X \rightarrow x$, and $P_x \rightarrow -i\hbar \frac{d}{dx}$.

$$\left[-\frac{\hbar^2}{2m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) + x^2 + y^2 + z^2 \right] \psi_\omega(x, y, z) = \omega \psi_\omega(x, y, z) \quad (1)$$

Once we have this form, we can change to other coordinates, e.g. spherical. Use

$$\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \equiv \nabla^2 \equiv \frac{1}{r^2} \left(\frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right)$$

And substitute into (1) to get

$$-\frac{\hbar^2}{2m} \left[\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \psi_\omega}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \psi_\omega}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \psi_\omega}{\partial \phi^2} \right] + r^2 \psi_\omega = \omega \psi_\omega \quad (2)$$

3.4 The Schrödinger Equation

3.4.1 Hamiltonian Setup

Simply substitute $x \rightarrow X$, $p \rightarrow P$ in the classical Hamiltonian. E.g. for (1) harmonic oscillator, (2) constant force f , and (3) particle of charge q in an EM field,

$$\begin{aligned} 1) \quad H &= \frac{p^2}{2m} + \frac{1}{2} m \omega^2 x^2 &\longrightarrow H &= \frac{P^2}{2m} + \frac{1}{2} m \omega^2 X^2 \\ 2) \quad H &= \frac{p^2}{2m} - f x &\longrightarrow H &= \frac{P^2}{2m} - f X \\ 3) \quad H &= \frac{|\vec{p} - \frac{q}{c} \vec{A}(\vec{r}, t)|^2}{2m} + q\phi(\vec{r}, t) &\longrightarrow H &= \frac{1}{2m} \left(P \cdot P - \frac{q}{c} A \cdot P - \frac{q}{c} P \cdot A + \frac{q^2}{c^2} A \cdot A \right) + q\phi \end{aligned}$$

Note: P doesn't commute with A (which depends on X, Y, Z).

3.4.2 Time-Independent Schrödinger Equation

Start from Postulate IV, $i\hbar \frac{\partial}{\partial t} |\psi\rangle = H |\psi\rangle$. We seek to find a propagator of the form

$$|\psi(t)\rangle = U(t) |\psi(0)\rangle$$

Use the energy basis, since $H |E\rangle = E |E\rangle$.

$$|\psi(t)\rangle = \sum_E |E\rangle \langle E|\psi(t)\rangle \equiv \sum_E a_E(t) |E\rangle$$

Apply $(i\hbar \frac{\partial}{\partial t} - H)$ and use $(i\hbar \frac{\partial}{\partial t} - H) |\psi(t)\rangle = 0$ to get

$$\begin{aligned} 0 &= \sum_E (i\hbar \frac{\partial}{\partial t} - H) a_E(t) |E\rangle = \sum_E (i\hbar \dot{a}_E - E a_E) |E\rangle \\ a_E(t) &= a_E(0) e^{-iEt/\hbar} \\ \langle E|\psi(t)\rangle &= \langle E|\psi(0)\rangle e^{-iEt/\hbar} \end{aligned}$$

Now plug back in to $|\psi\rangle$ to get the propagator:

$$\begin{aligned} |\psi(t)\rangle &= \sum_E e^{-iEt/\hbar} |E\rangle \langle E|\psi(0)\rangle \\ U(t) &= \sum_E e^{-iEt/\hbar} |E\rangle \langle E| = e^{-iHt/\hbar} \end{aligned} \tag{1}$$

Because H is unitary, $U(t)$ is unitary, thus time evolution is unitary and the norm is preserved: $\langle \psi(t)|\psi(t)\rangle = \langle \psi(0)|\psi(0)\rangle$.

3.4.3 Time-Dependent Schrödinger Equation

To approximate a time-dependent Hamiltonian, split $(0, t)$ into N intervals of width $\Delta = t/N$. Then

$$\begin{aligned} |\psi(t)\rangle &= |\psi(0)\rangle + \Delta \left. \frac{d|\psi\rangle}{dt} \right|_0 = |\psi(0)\rangle - \frac{i\Delta}{\hbar} H(0) |\psi(0)\rangle \\ &= \left[1 - \frac{i\Delta}{\hbar} H(0) \right] |\psi(0)\rangle \\ &\approx \exp \left[-\frac{i\Delta}{\hbar} H(0) \right] |\psi(0)\rangle \end{aligned}$$

The overall propagator is found by summing the changes:

$$\begin{aligned} |\psi(t)\rangle &= \prod_{n=0}^{N-1} e^{-i\Delta H(n\Delta)/\hbar} |\psi(0)\rangle \\ U(t) &= \exp \left[-\frac{i}{\hbar} \int_0^t H(t') dt' \right], \quad N \rightarrow \infty \end{aligned} \tag{2}$$

4 Simple Problems in 1D

4.1 The Free Particle

4.1.1 Allowed Energy States

Start from the Schrodinger equation:

$$i\hbar \frac{d}{dt} |\psi\rangle = H |\psi\rangle = \frac{P^2}{2m} |\psi\rangle$$

The normal modes (in energy basis) are

$$|\psi\rangle = e^{-iEt/\hbar} |E\rangle$$

We seek to find allowed E states. $|p\rangle$ is an eigenvector of both P and P^2 , so

$$\begin{aligned} \frac{P^2}{2m} |p\rangle = E |p\rangle &\longrightarrow \frac{p^2}{2m} |p\rangle = E |p\rangle \\ p &= \pm\sqrt{2mE} \end{aligned}$$

Thus for each eigenvalue E , we have 2 eigenvectors:

$$\begin{aligned} |E, +\rangle &= |p = \sqrt{2mE}\rangle \\ |E, -\rangle &= |p = -\sqrt{2mE}\rangle \end{aligned}$$

I.e. a particle with energy E may be found moving left or right with momentum $\sqrt{2mE}$. So far, this agrees with a classical analysis. The difference is that in QM, we can have a state like

$$|E\rangle = \beta |p = \sqrt{2mE}\rangle + \gamma |p = -\sqrt{2mE}\rangle$$

4.1.2 Propagator

Use the propagator from the time-independent Schrodinger equation:

$$U(t) = \int_{-\infty}^{\infty} |p\rangle \langle p| e^{-iE(p)t/\hbar} dp = \int_{-\infty}^{\infty} |p\rangle \langle p| e^{-ip^2t/2m\hbar} dp$$

Note that in the X basis, the propagator for a time slice is

$$\begin{aligned} U(x, t; x') &\equiv \langle x|U(t)|x'\rangle = \int_{-\infty}^{\infty} \langle x|p\rangle \langle p|x'\rangle e^{-ip^2t/2m\hbar} dp \\ &= \frac{1}{2m\hbar} \int_{-\infty}^{\infty} e^{ip(x-x')/\hbar} \cdot e^{-ip^2t/2m\hbar} dp \\ &= \sqrt{\frac{m}{2\pi\hbar it}} e^{im(x-x')^2/2\hbar t} \end{aligned}$$

In terms of the propagator, given the initial state $\psi(x', 0)$, we can find an arbitrary state $\psi(x, t)$ as

$$\psi(x, t) = \int U(x, t; x') \psi(x', 0) dx'$$

4.2 The Particle in a Box

4.2.1 Wavefunction

Consider a box potential:

$$V(x) = \begin{cases} 0 & |x| < L/2 \\ \infty & |x| \geq L/2 \end{cases}$$

Let I and III be the regions outside the box, while II is the region within the box. Use the eigenvalue equation in the X basis:

$$\frac{d^2\psi}{dx^2} + \frac{2m}{\hbar^2}(E - V)\psi = 0$$

In III, we have

$$\begin{aligned} \frac{d^2\psi_{\text{III}}}{dx^2} - \frac{2m(V_0 - E)}{\hbar^2}\psi_{\text{III}} &= 0 \quad , \quad V_0 \rightarrow \infty \\ \psi_{\text{III}} &= Ae^{-\kappa x} + Be^{\kappa x} \quad , \quad \kappa = \sqrt{\frac{2m(V_0 - E)}{\hbar^2}} \end{aligned}$$

We must have $B = 0$ because otherwise the wavefunction blows up as $x \rightarrow \infty$. Then as $V_0 \rightarrow \infty$, $\psi_{\text{III}} = 0$. This is as expected. Similar results hold for I: $\psi_{\text{I}} = 0$.

In II, the solution is that of a free particle:

$$\psi_{\text{II}} = Ae^{ikx} + Be^{-ikx} \quad , \quad k = \sqrt{\frac{2mE}{\hbar^2}}$$

Further, it must obey the boundary conditions:

$$\begin{aligned} \psi_{\text{I}}(-L/2) = \psi_{\text{II}}(-L/2) = 0 &\longrightarrow Ae^{ikL/2} + Be^{ikL/2} = 0 \\ \psi_{\text{III}}(L/2) = \psi_{\text{II}}(L/2) = 0 &\longrightarrow Ae^{ikL/2} + Be^{-ikL/2} = 0 \\ \begin{bmatrix} e^{-ikL/2} & e^{ikL/2} \\ e^{ikL/2} & e^{-ikL/2} \end{bmatrix} \begin{bmatrix} A \\ B \end{bmatrix} &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} \end{aligned}$$

For this to happen, the determinant must be 0, i.e. $e^{ikL} - e^{-ikL} = 0$, thus $k = n\pi/L$, where $n = 0, \pm 1, \pm 2, \dots$. We can find that $A = -e^{in\pi}B$, leading to two families of solutions:

$$\psi_n(x) = \begin{cases} \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right) & n \text{ even} \\ \sqrt{\frac{2}{L}} \cos\left(\frac{n\pi x}{L}\right) & n \text{ odd} \end{cases}$$

4.2.2 Allowed Energies

Note that the energy is quantized:

$$E = \frac{\hbar^2 k_n^2}{2m} = \frac{\hbar^2 \pi^2 n^2}{2mL^2}$$

And further, the lowest bound on energy (for non-trivial solutions) is nonzero: the ground state energy E_g is

$$E_g = \frac{\hbar^2 \pi^2}{2mL^2}$$

An “intuitive” explanation for this is that Δx is bounded in the well, so by the uncertainty principle, we cannot have a well-defined momentum of 0.

The uncertainty principle also provides a quick estimate of the ground energy. Start with $H = P^2/2m$. From the Ehrenfest theorem, $\langle H \rangle = \langle P^2 \rangle/2m$. Since $\langle P \rangle = 0$ inside the box, we can further write

$$\langle H \rangle = \frac{\langle (P - \langle P \rangle)^2 \rangle}{2m} = \frac{(\Delta P)^2}{2m}$$

From the uncertainty relations, $\Delta X \cdot \Delta P \geq \hbar/2$, so we have

$$\langle H \rangle \geq \frac{\hbar^2}{8m(\Delta x)^2} \geq \frac{\hbar^2}{2mL^2}$$

In energy eigenstates, $\langle H \rangle = E$. Thus we have

$$E \geq \frac{\hbar^2}{2mL^2}$$

Which is the estimated order of the ground state, accurate up to a factor of π^2 .