

An introduction to chaining, and applications to sublinear algorithms

Jelani Nelson
Harvard

August 28, 2015

What's this talk about?

What's this talk about?

Given a collection of random variables X_1, X_2, \dots , we would like to say that $\max_i X_i$ is small with high probability. (Happens all over computer science, e.g. “Chernion” (Chernoff+Union) bound)

What's this talk about?

Given a collection of random variables X_1, X_2, \dots , we would like to say that $\max_i X_i$ is small with high probability. (Happens all over computer science, e.g. “Chernion” (Chernoff+Union) bound)

Today's topic: Beating the Union Bound

What's this talk about?

Given a collection of random variables X_1, X_2, \dots , we would like to say that $\max_i X_i$ is small with high probability. (Happens all over computer science, e.g. “Chernion” (Chernoff+Union) bound)

Today's topic: Beating the Union Bound

Disclaimer: This is an educational talk, about ideas which aren't mine.

A first example

- $T \subset B_{\ell_2^n}$

A first example

- $T \subset B_{\ell_2^n}$
- Random variables $(Z_x)_{x \in T}$
 $Z_x = \langle g, x \rangle$ for a vector g with i.i.d. $\mathcal{N}(0, 1)$ entries

A first example

- $T \subset B_{\ell_2^n}$
- Random variables $(Z_x)_{x \in T}$
 $Z_x = \langle g, x \rangle$ for a vector g with i.i.d. $\mathcal{N}(0, 1)$ entries
- Define *gaussian mean width* $\mathfrak{g}(T) = \mathbb{E}_g \sup_{x \in T} Z_x$

A first example

- $T \subset B_{\ell_2^n}$
- Random variables $(Z_x)_{x \in T}$
 $Z_x = \langle g, x \rangle$ for a vector g with i.i.d. $\mathcal{N}(0, 1)$ entries
- Define *gaussian mean width* $\mathfrak{g}(T) = \mathbb{E}_g \sup_{x \in T} Z_x$
- How can we bound $\mathfrak{g}(T)$?

A first example

- $T \subset B_{\ell_2^n}$
- Random variables $(Z_x)_{x \in T}$
 $Z_x = \langle g, x \rangle$ for a vector g with i.i.d. $\mathcal{N}(0, 1)$ entries
- Define *gaussian mean width* $g(T) = \mathbb{E}_g \sup_{x \in T} Z_x$
- How can we bound $g(T)$?
- **This talk:** four progressively tighter ways to bound $g(T)$, then applications of techniques to some TCS problems

Gaussian mean width bound 1: union bound

- $g(T) = \mathbb{E} \sup_{x \in T} Z_x = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$

Gaussian mean width bound 1: union bound

- $g(T) = \mathbb{E} \sup_{x \in T} Z_x = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Z_x is a gaussian with variance one

Gaussian mean width bound 1: union bound

- $g(T) = \mathbb{E} \sup_{x \in T} Z_x = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Z_x is a gaussian with variance one

$$\mathbb{E} \sup_{x \in T} Z_x = \int_0^\infty \mathbb{P}(\sup_{x \in T} Z_x > u) du$$

Gaussian mean width bound 1: union bound

- $g(T) = \mathbb{E} \sup_{x \in T} Z_x = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Z_x is a gaussian with variance one

$$\begin{aligned} \mathbb{E} \sup_{x \in T} Z_x &= \int_0^\infty \mathbb{P}(\sup_{x \in T} Z_x > u) du \\ &= \underbrace{\int_0^{u_*} \mathbb{P}(\sup_{x \in T} Z_x > u) du}_{\leq 1} + \int_{u_*}^\infty \underbrace{\mathbb{P}(\sup_{x \in T} Z_x > u)}_{\leq |T| \cdot e^{-u^2/2} \text{ (union bound)}} du \end{aligned}$$

Gaussian mean width bound 1: union bound

- $g(T) = \mathbb{E} \sup_{x \in T} Z_x = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Z_x is a gaussian with variance one

$$\begin{aligned} \mathbb{E} \sup_{x \in T} Z_x &= \int_0^\infty \mathbb{P}(\sup_{x \in T} Z_x > u) du \\ &= \underbrace{\int_0^{u_*} \mathbb{P}(\sup_{x \in T} Z_x > u) du}_{\leq 1} + \int_{u_*}^\infty \underbrace{\mathbb{P}(\sup_{x \in T} Z_x > u)}_{\leq |T| \cdot e^{-u^2/2} \text{ (union bound)}} du \\ &\leq u_* + |T| \cdot e^{-u_*^2/2} \\ &\lesssim \sqrt{\log |T|} \text{ (set } u_* = \sqrt{2 \log |T|}) \end{aligned}$$

Gaussian mean width bound 2: ε -net

- $g(T) = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Let S_ε be ε -net of (T, ℓ_2)

Gaussian mean width bound 2: ε -net

- $\mathfrak{g}(T) = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Let S_ε be ε -net of (T, ℓ_2)
- $\langle g, x \rangle = \langle g, x' \rangle + \langle g, x - x' \rangle$ ($x' = \operatorname{argmin}_{y \in T} \|x - y\|_2$)
$$\mathfrak{g}(T) \leq \mathfrak{g}(S_\varepsilon) + \mathbb{E}_g \sup_{x \in T} \underbrace{\langle g, x - x' \rangle}_{\leq \varepsilon \cdot \|g\|_2}$$

Gaussian mean width bound 2: ε -net

- $g(T) = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Let S_ε be ε -net of (T, ℓ_2)
- $\langle g, x \rangle = \langle g, x' \rangle + \langle g, x - x' \rangle$ ($x' = \operatorname{argmin}_{y \in T} \|x - y\|_2$)
$$g(T) \leq g(S_\varepsilon) + \mathbb{E}_g \sup_{x \in T} \underbrace{\langle g, x - x' \rangle}_{\leq \varepsilon \cdot \|g\|_2}$$
- $\lesssim \sqrt{\log |S_\varepsilon|} + \varepsilon (\mathbb{E}_g \|g\|_2^2)^{1/2}$
- $\lesssim \log^{1/2} \underbrace{\mathcal{N}(T, \ell_2, \varepsilon)}_{\text{smallest } \varepsilon\text{-net size}} + \varepsilon \sqrt{n}$

Gaussian mean width bound 2: ε -net

- $g(T) = \mathbb{E} \sup_{x \in T} \langle g, x \rangle$
- Let S_ε be ε -net of (T, ℓ_2)
- $\langle g, x \rangle = \langle g, x' \rangle + \langle g, x - x' \rangle$ ($x' = \operatorname{argmin}_{y \in T} \|x - y\|_2$)
$$g(T) \leq g(S_\varepsilon) + \mathbb{E}_g \sup_{x \in T} \underbrace{\langle g, x - x' \rangle}_{\leq \varepsilon \cdot \|g\|_2}$$
- $\lesssim \sqrt{\log |S_\varepsilon|} + \varepsilon (\mathbb{E}_g \|g\|_2^2)^{1/2}$
- $\lesssim \log^{1/2} \underbrace{\mathcal{N}(T, \ell_2, \varepsilon)}_{\text{smallest } \varepsilon\text{-net size}} + \varepsilon \sqrt{n}$
- Choose ε to optimize bound; can never be worse than last slide (which amounts to choosing $\varepsilon = 0$)

Gaussian mean width bound 3: ε -net sequence

- S_k is a $(1/2^k)$ -net of T , $k \geq 0$
 $\pi_k x$ is closest point in S_k to $x \in T$, $\Delta_k x = \pi_k x - \pi_{k-1} x$

Gaussian mean width bound 3: ε -net sequence

- S_k is a $(1/2^k)$ -net of T , $k \geq 0$
 $\pi_k x$ is closest point in S_k to $x \in T$, $\Delta_k x = \pi_k x - \pi_{k-1} x$
- wlog $|T| < \infty$ (else apply this slide to ε -net of T for ε small)
- $\langle g, x \rangle = \langle g, \pi_0 x \rangle + \sum_{k=1}^{\infty} \langle g, \Delta_k x \rangle$

Gaussian mean width bound 3: ε -net sequence

- S_k is a $(1/2^k)$ -net of T , $k \geq 0$
 $\pi_k x$ is closest point in S_k to $x \in T$, $\Delta_k x = \pi_k x - \pi_{k-1} x$
- wlog $|T| < \infty$ (else apply this slide to ε -net of T for ε small)
- $\langle g, x \rangle = \langle g, \pi_0 x \rangle + \sum_{k=1}^{\infty} \langle g, \Delta_k x \rangle$
- $$g(T) \leq \underbrace{\mathbb{E} \sup_{g \supset x \in T} \langle g, \pi_0 x \rangle}_0 + \sum_{k=1}^{\infty} \mathbb{E}_g \sup_{x \in T} \langle g, \Delta_k x \rangle$$

Gaussian mean width bound 3: ε -net sequence

- S_k is a $(1/2^k)$ -net of T , $k \geq 0$
 $\pi_k x$ is closest point in S_k to $x \in T$, $\Delta_k x = \pi_k x - \pi_{k-1} x$
- wlog $|T| < \infty$ (else apply this slide to ε -net of T for ε small)
- $\langle g, x \rangle = \langle g, \pi_0 x \rangle + \sum_{k=1}^{\infty} \langle g, \Delta_k x \rangle$
- $$g(T) \leq \underbrace{\mathbb{E} \sup_{g \supset x \in T} \langle g, \pi_0 x \rangle}_0 + \sum_{k=1}^{\infty} \mathbb{E}_g \sup_{x \in T} \langle g, \Delta_k x \rangle$$
- $$|\{\Delta_k x : x \in T\}| \leq \mathcal{N}(T, \ell_2, 1/2^k) \cdot \mathcal{N}(T, \ell_2, 1/2^{k-1})$$
$$\leq (\mathcal{N}(T, \ell_2, 1/2^k))^2$$

Gaussian mean width bound 3: ε -net sequence

- S_k is a $(1/2^k)$ -net of T , $k \geq 0$
 $\pi_k x$ is closest point in S_k to $x \in T$, $\Delta_k x = \pi_k x - \pi_{k-1} x$
- wlog $|T| < \infty$ (else apply this slide to ε -net of T for ε small)
- $\langle g, x \rangle = \langle g, \pi_0 x \rangle + \sum_{k=1}^{\infty} \langle g, \Delta_k x \rangle$
- $$g(T) \leq \underbrace{\mathbb{E} \sup_{x \in T} \langle g, \pi_0 x \rangle}_0 + \sum_{k=1}^{\infty} \mathbb{E}_g \sup_{x \in T} \langle g, \Delta_k x \rangle$$
- $$|\{\Delta_k x : x \in T\}| \leq \mathcal{N}(T, \ell_2, 1/2^k) \cdot \mathcal{N}(T, \ell_2, 1/2^{k-1})$$
$$\leq (\mathcal{N}(T, \ell_2, 1/2^k))^2$$
- $$g(T) \lesssim \sum_{k=1}^{\infty} (1/2^k) \cdot \log^{1/2} \mathcal{N}(T, \ell_2, 1/2^k)$$
$$\lesssim \int_0^{\infty} \log^{1/2} \mathcal{N}(T, \ell_2, u) du \text{ (Dudley's theorem)}$$

Gaussian mean width bound 4: generic chaining

- Again, wlog $|T| < \infty$. Define $T_0 \subseteq T_1 \subseteq \dots \subseteq T_{k_*} = T$
 $|T_0| = 1, |T_k| \leq 2^{2^k}$ (call such a sequence “admissible”)

Gaussian mean width bound 4: generic chaining

- Again, wlog $|T| < \infty$. Define $T_0 \subseteq T_1 \subseteq \dots \subseteq T_{k_*} = T$
 $|T_0| = 1, |T_k| \leq 2^{2^k}$ (call such a sequence “admissible”)
- Exercise: show Dudley’s theorem is equivalent to
$$g(T) \lesssim \inf_{\{T_k\} \text{ admissible}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$$
(should pick T_k to be the best $\varepsilon = \varepsilon(k)$ net of size 2^{2^k})

Gaussian mean width bound 4: generic chaining

- Again, wlog $|T| < \infty$. Define $T_0 \subseteq T_1 \subseteq \dots \subseteq T_{k_*} = T$
 $|T_0| = 1, |T_k| \leq 2^{2^k}$ (call such a sequence “admissible”)
- Exercise: show Dudley’s theorem is equivalent to
$$g(T) \lesssim \inf_{\{T_k\} \text{ admissible}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$$
(should pick T_k to be the best $\varepsilon = \varepsilon(k)$ net of size 2^{2^k})
- Fernique’76*: can pull the \sup_x *outside* the sum
- $g(T) \lesssim \inf_{\{T_k\}} \sup_{x \in T} \sum_{k=1}^{\infty} 2^{k/2} \cdot d_{\ell_2}(x, T_k) \stackrel{\text{def}}{=} \gamma_2(T, \ell_2)$

Gaussian mean width bound 4: generic chaining

- Again, wlog $|T| < \infty$. Define $T_0 \subseteq T_1 \subseteq \dots \subseteq T_{k_*} = T$
 $|T_0| = 1, |T_k| \leq 2^{2^k}$ (call such a sequence “admissible”)
- Exercise: show Dudley’s theorem is equivalent to
$$g(T) \lesssim \inf_{\{T_k\} \text{ admissible}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$$
(should pick T_k to be the best $\varepsilon = \varepsilon(k)$ net of size 2^{2^k})
- Fernique’76*: can pull the \sup_x *outside* the sum
- $g(T) \lesssim \inf_{\{T_k\}} \sup_{x \in T} \sum_{k=1}^{\infty} 2^{k/2} \cdot d_{\ell_2}(x, T_k) \stackrel{\text{def}}{=} \gamma_2(T, \ell_2)$
* equivalent upper bound proven by Fernique (who minimized some integral over all measures over T), but reformulated in terms of admissible sequences by Talgarand

Gaussian mean width bound 4: generic chaining

Proof of Fernique's bound

$$g(T) \leq \underbrace{\mathbb{E} \sup_{g, x \in T} \langle g, \pi_0 x \rangle}_0 + \mathbb{E} \sup_{g, x \in T} \sum_{k=1}^{\infty} \underbrace{\langle g, \Delta_k x \rangle}_{Y_k} \quad (\text{from before})$$

- $\forall t, \mathbb{P}(Y_k > t 2^{k/2} \|\Delta_k x\|_2) \leq e^{-t^2 2^k / 2}$ (gaussian decay)

Gaussian mean width bound 4: generic chaining

Proof of Fernique's bound

$$g(T) \leq \underbrace{\mathbb{E} \sup_{g, x \in T} \langle g, \pi_0 x \rangle}_0 + \mathbb{E} \sup_{g, x \in T} \sum_{k=1}^{\infty} \underbrace{\langle g, \Delta_k x \rangle}_{Y_k} \quad (\text{from before})$$

- $\forall t, \mathbb{P}(Y_k > t2^{k/2} \|\Delta_k x\|_2) \leq e^{-t^2 2^k / 2}$ (gaussian decay)
- $\mathbb{P}(\exists x, k \ Y_k > t2^{k/2} \|\Delta_k x\|_2) \leq \sum_k (2^{2^k})^2 e^{-t^2 2^k / 2}$

Gaussian mean width bound 4: generic chaining

Proof of Fernique's bound

$$g(T) \leq \underbrace{\mathbb{E} \sup_{g, x \in T} \langle g, \pi_0 x \rangle}_0 + \mathbb{E} \sup_{g, x \in T} \sum_{k=1}^{\infty} \underbrace{\langle g, \Delta_k x \rangle}_{Y_k} \quad (\text{from before})$$

- $\forall t, \mathbb{P}(Y_k > t2^{k/2} \|\Delta_k x\|_2) \leq e^{-t^2 2^k / 2}$ (gaussian decay)
- $\mathbb{P}(\exists x, k \ Y_k > t2^{k/2} \|\Delta_k x\|_2) \leq \sum_k (2^{2^k})^2 e^{-t^2 2^k / 2}$

$$\mathbb{E} \sup_{g, x \in T} \sum_k Y_k = \int_0^{\infty} \mathbb{P}(\sup_{x \in T} \sum_k Y_k > u) du$$

Gaussian mean width bound 4: generic chaining

$$\mathbb{E} \sup_{g} \sum_{x \in T} Y_k = \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > u) du$$

Gaussian mean width bound 4: generic chaining

$$\begin{aligned}\mathbb{E} \sup_{g} \sum_{k \in T} Y_k &= \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > u) du \\ &= \gamma_2(T, \ell_2) \cdot \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2) dt \\ &\quad (\text{change of variables: } u = t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2 \simeq t \gamma_2(T, \ell_2))\end{aligned}$$

Gaussian mean width bound 4: generic chaining

$$\begin{aligned}\mathbb{E} \sup_{g} \sum_{x \in T} Y_k &= \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > u) du \\&= \gamma_2(T, \ell_2) \cdot \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2) dt \\&\quad (\text{change of variables: } u = t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2 \simeq t \gamma_2(T, \ell_2)) \\&\leq \gamma_2(T, \ell_2) \cdot [2 + \int_2^\infty \left(\sum_{k=1}^\infty (2^{2^k})^2 e^{-t^2 2^k / 2} \right) dt] \\&\simeq \gamma_2(T, \ell_2)\end{aligned}$$

Gaussian mean width bound 4: generic chaining

$$\begin{aligned}\mathbb{E} \sup_{x \in T} \sum_k Y_k &= \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > u) du \\&= \gamma_2(T, \ell_2) \cdot \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2) dt \\&\quad (\text{change of variables: } u = t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2 \simeq t \gamma_2(T, \ell_2)) \\&\leq \gamma_2(T, \ell_2) \cdot [2 + \int_2^\infty \left(\sum_{k=1}^\infty (2^{2^k})^2 e^{-t^2 2^{k/2}} \right) dt] \\&\simeq \gamma_2(T, \ell_2)\end{aligned}$$

- Conclusion: $\mathfrak{g}(T) \lesssim \gamma_2(T, \ell_2)$

Gaussian mean width bound 4: generic chaining

$$\begin{aligned}\mathbb{E} \sup_{x \in T} \sum_k Y_k &= \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > u) du \\&= \gamma_2(T, \ell_2) \cdot \int_0^\infty \mathbb{P}(\sup_{x \in T} \sum_k Y_k > t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2) dt \\&\quad (\text{change of variables: } u = t \sup_{x \in T} \sum_k 2^{k/2} \|\Delta_k x\|_2 \simeq t \gamma_2(T, \ell_2)) \\&\leq \gamma_2(T, \ell_2) \cdot [2 + \int_2^\infty \left(\sum_{k=1}^\infty (2^{2^k})^2 e^{-t^2 2^k / 2} \right) dt] \\&\simeq \gamma_2(T, \ell_2)\end{aligned}$$

- Conclusion: $\mathfrak{g}(T) \lesssim \gamma_2(T, \ell_2)$
- Talagrand: $\mathfrak{g}(T) \simeq \gamma_2(T, \ell_2)$ (won't show today)
(“Majorizing measures theorem”)

Are these bounds really different?

- $\gamma_2(T, \ell_2)$: $\inf_{\{T_k\}} \sup_{x \in T} \sum_{k=1}^{\infty} 2^{k/2} \cdot d_{\ell_2}(x, T_k)$
- **Dudley:** $\inf_{\{T_k\}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$
 $\simeq \int_0^{\infty} \log^{1/2} \mathcal{N}(T, \ell_2, u) du$

Are these bounds really different?

- $\gamma_2(T, \ell_2)$: $\inf_{\{T_k\}} \sup_{x \in T} \sum_{k=1}^{\infty} 2^{k/2} \cdot d_{\ell_2}(x, T_k)$
- **Dudley:** $\inf_{\{T_k\}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$
 $\simeq \int_0^{\infty} \log^{1/2} \mathcal{N}(T, \ell_2, u) du$
- Dudley not optimal: $T = B_{\ell_1^n}$

Are these bounds really different?

- $\gamma_2(T, \ell_2)$: $\inf_{\{T_k\}} \sup_{x \in T} \sum_{k=1}^{\infty} 2^{k/2} \cdot d_{\ell_2}(x, T_k)$
- **Dudley:** $\inf_{\{T_k\}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$
 $\simeq \int_0^{\infty} \log^{1/2} \mathcal{N}(T, \ell_2, u) du$
- Dudley not optimal: $T = B_{\ell_1^n}$
- $\sup_{x \in B_{\ell_1^n}} \langle g, x \rangle = \|g\|_{\infty}$, so $\mathfrak{g}(T) \simeq \sqrt{\log n}$
- **Exercise:** Come up with admissible $\{T_k\}$ yielding $\gamma_2 \lesssim \sqrt{\log n}$ (must exist by majorizing measures)

Are these bounds really different?

- $\gamma_2(T, \ell_2)$: $\inf_{\{T_k\}} \sup_{x \in T} \sum_{k=1}^{\infty} 2^{k/2} \cdot d_{\ell_2}(x, T_k)$
- **Dudley:** $\inf_{\{T_k\}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$
 $\simeq \int_0^{\infty} \log^{1/2} \mathcal{N}(T, \ell_2, u) du$
- Dudley not optimal: $T = B_{\ell_1^n}$
- $\sup_{x \in B_{\ell_1^n}} \langle g, x \rangle = \|g\|_{\infty}$, so $\mathfrak{g}(T) \simeq \sqrt{\log n}$
- **Exercise:** Come up with admissible $\{T_k\}$ yielding $\gamma_2 \lesssim \sqrt{\log n}$ (must exist by majorizing measures)
- Dudley: $\log \mathcal{N}(B_{\ell_1^n}, \ell_2, u) \simeq (1/u^2) \log n$ for u not too small (consider just covering $(1/u^2)$ -sparse vectors with u^2 in each coordinate). Dudley can only give $\mathfrak{g}(B_{\ell_1^n}) \lesssim \log^{3/2} n$.

Are these bounds really different?

- $\gamma_2(T, \ell_2)$: $\inf_{\{T_k\}} \sup_{x \in T} \sum_{k=1}^{\infty} 2^{k/2} \cdot d_{\ell_2}(x, T_k)$
- **Dudley:** $\inf_{\{T_k\}} \sum_{k=1}^{\infty} 2^{k/2} \cdot \sup_{x \in T} d_{\ell_2}(x, T_k)$
 $\simeq \int_0^{\infty} \log^{1/2} \mathcal{N}(T, \ell_2, u) du$
- Dudley not optimal: $T = B_{\ell_1^n}$
- $\sup_{x \in B_{\ell_1^n}} \langle g, x \rangle = \|g\|_{\infty}$, so $\mathfrak{g}(T) \simeq \sqrt{\log n}$
- **Exercise:** Come up with admissible $\{T_k\}$ yielding $\gamma_2 \lesssim \sqrt{\log n}$ (must exist by majorizing measures)
- Dudley: $\log \mathcal{N}(B_{\ell_1^n}, \ell_2, u) \simeq (1/u^2) \log n$ for u not too small (consider just covering $(1/u^2)$ -sparse vectors with u^2 in each coordinate). Dudley can only give $\mathfrak{g}(B_{\ell_1^n}) \lesssim \log^{3/2} n$.
- Simple vanilla ε -net argument gives $\mathfrak{g}(B_{\ell_1^n}) \lesssim \text{poly}(n)$.

High probability

- So far just talked about $g(T) = \mathbb{E}_g \sup_{x \in T} Z_x$
But what if we want to know $\sup_{x \in T} Z_x$ is small whp, not just in expectation?

High probability

- So far just talked about $g(T) = \mathbb{E}_g \sup_{x \in T} Z_x$
But what if we want to know $\sup_{x \in T} Z_x$ is small whp, not just in expectation?
- Usual approach: bound $\mathbb{E}_g \sup_{x \in T} Z_x^p$ for large p and do Markov (“moment method”)
Can bound moments using chaining too; see (Dirksen’13)

Applications in computer science

- Fast RIP matrices (Candès, Tao'06), (Rudelson, Vershynin'06), (Cheragchi, Guruswami, Velingker'13), (N., Price, Wootters'14), (Bourgain'14), (Haviv, Regev'15)
- Fast JL (Ailon, Liberty'11), (Krahmer, Ward'11), (Bourgain, Dirksen, N.'15), (Oymak, Recht, Soltanolkotabi'15)
- Instance-wise JL bounds (Gordon'88), (Klartag, Mendelson'05), (Mendelson, Pajor, Tomczak-Jaegermann'07), (Dirksen'14)
- Approximate nearest neighbor (Indyk, Naor'07)
- Deterministic algorithm to estimate graph cover time (Ding, Lee, Peres'11)
- List-decodability of random codes (Wootters'13), (Rudra, Wootters'14)
- ...

A chaining result for quadratic forms

Theorem

[Krahmer, Mendelson, Rauhut'14] Let $\mathcal{A} \subset \mathbb{R}^{n \times n}$ be a family of matrices, and let $\sigma_1, \dots, \sigma_n$ be independent subgaussians. Then

$$\begin{aligned} & \mathbb{E} \sup_{A \in \mathcal{A}} \left| \|A\sigma\|_2^2 - \mathbb{E}_\sigma \|A\sigma\|_2^2 \right| \\ & \lesssim \gamma_2^2(\mathcal{A}, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) + \gamma_2(\mathcal{A}, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) \cdot \Delta_F(\mathcal{A}) + \Delta_{\ell_2 \rightarrow \ell_2}(\mathcal{A}) \cdot \Delta_F(\mathcal{A}) \end{aligned}$$

(Δ_X is diameter under X -norm)

A chaining result for quadratic forms

Theorem

[Krahmer, Mendelson, Rauhut'14] Let $\mathcal{A} \subset \mathbb{R}^{n \times n}$ be a family of matrices, and let $\sigma_1, \dots, \sigma_n$ be independent subgaussians. Then

$$\begin{aligned} \mathbb{E} \sup_{A \in \mathcal{A}} | \|A\sigma\|_2^2 - \mathbb{E} \|A\sigma\|_2^2 | \\ \lesssim \gamma_2^2(\mathcal{A}, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) + \gamma_2(\mathcal{A}, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) \cdot \Delta_F(\mathcal{A}) + \Delta_{\ell_2 \rightarrow \ell_2}(\mathcal{A}) \cdot \Delta_F(\mathcal{A}) \end{aligned}$$

(Δ_X is diameter under X -norm)

Won't show proof today, but it is similar to bounding $g(T)$ (with some extra tricks). See <http://people.seas.harvard.edu/~minilek/madalgo2015/>, Lecture 3.

Instance-wise bounds for JL

Corollary (Gordon'88, Klartag-Mendelson'05, Mendelson, Pajor, Tomczak-Jaegermann'07, Dirksen'14)

For $T \subseteq S^{n-1}$ and $0 < \varepsilon < 1/2$, let $\Pi \in \mathbb{R}^{m \times n}$ have independent subgaussian independent entries with mean zero and variance $1/m$ for $m \gtrsim (\mathfrak{g}^2(T)+1)/\varepsilon^2$. Then

$$\mathbb{E} \sup_{x \in T} |||\Pi x||_2^2 - 1| < \varepsilon$$

Instance-wise bounds for JL

Proof of Gordon's theorem

- For $x \in T$ let A_x denote the $m \times mn$ matrix:

$$A_x = \frac{1}{\sqrt{m}} \cdot \begin{bmatrix} x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & x_1 & \cdots & x_n \end{bmatrix}.$$

Instance-wise bounds for JL

Proof of Gordon's theorem

- For $x \in T$ let A_x denote the $m \times mn$ matrix:

$$A_x = \frac{1}{\sqrt{m}} \cdot \begin{bmatrix} x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & x_1 & \cdots & x_n \end{bmatrix}.$$

- Then $\|\Pi x\|_2^2 = \|A_x \sigma\|_2^2$, where σ is formed by concatenating rows of Π (multiplied by \sqrt{m}).

Instance-wise bounds for JL

Proof of Gordon's theorem

- For $x \in T$ let A_x denote the $m \times mn$ matrix:

$$A_x = \frac{1}{\sqrt{m}} \cdot \begin{bmatrix} x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & x_1 & \cdots & x_n \end{bmatrix}.$$

- Then $\|\Pi x\|_2^2 = \|A_x \sigma\|_2^2$, where σ is formed by concatenating rows of Π (multiplied by \sqrt{m}).
- $\|A_x - A_y\| = \|A_{x-y}\| = (1/\sqrt{m}) \cdot \|x - y\|_2$
 $\Rightarrow \gamma_2(\mathcal{A}_T, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) = \gamma_2(T, \ell_2) \simeq \mathfrak{g}(T)$

Instance-wise bounds for JL

Proof of Gordon's theorem

- For $x \in T$ let A_x denote the $m \times mn$ matrix:

$$A_x = \frac{1}{\sqrt{m}} \cdot \begin{bmatrix} x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & x_1 & \cdots & x_n \end{bmatrix}.$$

- Then $\|\Pi x\|_2^2 = \|A_x \sigma\|_2^2$, where σ is formed by concatenating rows of Π (multiplied by \sqrt{m}).
- $\|A_x - A_y\| = \|A_{x-y}\| = (1/\sqrt{m}) \cdot \|x - y\|_2$
 $\Rightarrow \gamma_2(\mathcal{A}_T, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) = \gamma_2(T, \ell_2) \simeq \mathfrak{g}(T)$
- $\Delta_F(\mathcal{A}_T) = 1$, $\Delta_{\ell_2 \rightarrow \ell_2}(\mathcal{A}_T) = 1/\sqrt{m}$

Instance-wise bounds for JL

Proof of Gordon's theorem

- For $x \in T$ let A_x denote the $m \times mn$ matrix:

$$A_x = \frac{1}{\sqrt{m}} \cdot \begin{bmatrix} x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & x_1 & \cdots & x_n \end{bmatrix}.$$

- Then $\|\Pi x\|_2^2 = \|A_x \sigma\|_2^2$, where σ is formed by concatenating rows of Π (multiplied by \sqrt{m}).
- $\|A_x - A_y\| = \|A_{x-y}\| = (1/\sqrt{m}) \cdot \|x - y\|_2$
 $\Rightarrow \gamma_2(\mathcal{A}_T, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) = \gamma_2(T, \ell_2) \simeq \mathfrak{g}(T)$
- $\Delta_F(\mathcal{A}_T) = 1$, $\Delta_{\ell_2 \rightarrow \ell_2}(\mathcal{A}_T) = 1/\sqrt{m}$
- Thus $\mathbb{E}_\Pi \sup_{x \in T} \left| \|\Pi x\|_2^2 - 1 \right| \lesssim \mathfrak{g}^2(T)/m + \mathfrak{g}(T)/\sqrt{m} + 1/\sqrt{m}$

Instance-wise bounds for JL

Proof of Gordon's theorem

- For $x \in T$ let A_x denote the $m \times mn$ matrix:

$$A_x = \frac{1}{\sqrt{m}} \cdot \begin{bmatrix} x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & x_1 & \cdots & x_n & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & 0 & x_1 & \cdots & x_n \end{bmatrix}.$$

- Then $\|\Pi x\|_2^2 = \|A_x \sigma\|_2^2$, where σ is formed by concatenating rows of Π (multiplied by \sqrt{m}).
- $\|A_x - A_y\| = \|A_{x-y}\| = (1/\sqrt{m}) \cdot \|x - y\|_2$
 $\Rightarrow \gamma_2(\mathcal{A}_T, \|\cdot\|_{\ell_2 \rightarrow \ell_2}) = \gamma_2(T, \ell_2) \simeq \mathfrak{g}(T)$
- $\Delta_F(\mathcal{A}_T) = 1$, $\Delta_{\ell_2 \rightarrow \ell_2}(\mathcal{A}_T) = 1/\sqrt{m}$
- Thus $\mathbb{E}_\Pi \sup_{x \in T} \left| \|\Pi x\|_2^2 - 1 \right| \lesssim \mathfrak{g}^2(T)/m + \mathfrak{g}(T)/\sqrt{m} + 1/\sqrt{m}$
- Set $m \gtrsim (\mathfrak{g}^2(T)+1)/\varepsilon^2$

Consequences of Gordon's theorem

$$m \gtrsim (\mathfrak{g}^2(T)+1)/\varepsilon^2$$

- $|T| < \infty$: $\mathfrak{g}^2(T) \lesssim \log |T|$ (JL)
- T a d -dim subspace: $\mathfrak{g}^2(T) \simeq d$ (subspace embeddings)
- T all k -sparse vectors: $\mathfrak{g}^2(T) \simeq k \log(n/k)$ (RIP)

Consequences of Gordon's theorem

$$m \gtrsim (\mathfrak{g}^2(T)+1)/\varepsilon^2$$

- $|T| < \infty$: $\mathfrak{g}^2(T) \lesssim \log |T|$ (JL)
- T a d -dim subspace: $\mathfrak{g}^2(T) \simeq d$ (subspace embeddings)
- T all k -sparse vectors: $\mathfrak{g}^2(T) \simeq k \log(n/k)$ (RIP)
- more applications to constrained least squares, manifold learning, model-based compressed sensing, ...
(see (Dirksen'14) and (Bourgain, Dirksen, N.'15))

Chaining isn't just for gaussians

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

“Restricted isometry property” useful in compressed sensing.

$$T = \{x : \|x\|_0 \leq k, \|x\|_2 = 1\}.$$

Theorem (Candès-Tao'06, Donoho'06, Candès'08)

If Π satisfies (ε_, k) -RIP for $\varepsilon_* < \sqrt{2} - 1$ then there is a linear program which, given Πx and Π as input, recovers \tilde{x} in polynomial time such that $\|x - \tilde{x}\|_2 \leq O(1/\sqrt{k}) \cdot \min_{\|y\|_0 \leq k} \|x - y\|_1$.*

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

“Restricted isometry property” useful in compressed sensing.

$$T = \{x : \|x\|_0 \leq k, \|x\|_2 = 1\}.$$

Theorem (Candès-Tao'06, Donoho'06, Candès'08)

If Π satisfies (ε_, k) -RIP for $\varepsilon_* < \sqrt{2} - 1$ then there is a linear program which, given Πx and Π as input, recovers \tilde{x} in polynomial time such that $\|x - \tilde{x}\|_2 \leq O(1/\sqrt{k}) \cdot \min_{\|y\|_0 \leq k} \|x - y\|_1$.*

Of interest to show sampling rows of discrete Fourier matrix is RIP

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

- (Unnormalized) Fourier matrix F , rows: z_1^*, \dots, z_n^*
- $\delta_1, \dots, \delta_n$ independent Bernoulli with expectation m/n

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

- (Unnormalized) Fourier matrix F , rows: z_1^*, \dots, z_n^*
- $\delta_1, \dots, \delta_n$ independent Bernoulli with expectation m/n
- Want

$$\mathbb{E}_{\delta} \sup_{\substack{T \subset [n] \\ |T| \leq k}} \left\| I_T - \frac{1}{m} \sum_{i=1}^n \delta_i z_i^{(T)} z_i^{(T)*} \right\| < \varepsilon$$

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

$$\text{LHS} = \mathbb{E}_{\delta} \sup_{\substack{T \subset [n] \\ |T| \leq k}} \left\| \overbrace{\mathbb{E}_{\delta'} \frac{1}{m} \sum_{i=1}^n \delta'_i z_i^{(T)} z_i^{(T)*}}^{I_T} - \frac{1}{m} \sum_{i=1}^n \delta_i z_i^{(T)} z_i^{(T)*} \right\|$$

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

$$\begin{aligned} \text{LHS} &= \mathbb{E}_{\delta} \sup_{\substack{T \subset [n] \\ |T| \leq k}} \left\| \overbrace{\mathbb{E}_{\delta'} \frac{1}{m} \sum_{i=1}^n \delta'_i z_i^{(T)} z_i^{(T)*}}^{I_T} - \frac{1}{m} \sum_{i=1}^n \delta_i z_i^{(T)} z_i^{(T)*} \right\| \\ &\leq \frac{1}{m} \mathbb{E}_{\delta, \delta'} \sup_T \left\| \sum_{i=1}^n (\delta'_i - \delta_i) z_i^{(T)} z_i^{(T)*} \right\| \quad (\text{Jensen}) \end{aligned}$$

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

$$\begin{aligned}
 \text{LHS} &= \mathbb{E}_{\delta} \sup_{\substack{T \subset [n] \\ |T| \leq k}} \left\| \overbrace{\mathbb{E}_{\delta'} \frac{1}{m} \sum_{i=1}^n \delta'_i z_i^{(T)} z_i^{(T)*}}^{I_T} - \frac{1}{m} \sum_{i=1}^n \delta_i z_i^{(T)} z_i^{(T)*} \right\| \\
 &\leq \frac{1}{m} \mathbb{E}_{\delta, \delta'} \sup_T \left\| \sum_{i=1}^n (\delta'_i - \delta_i) z_i^{(T)} z_i^{(T)*} \right\| \quad (\text{Jensen}) \\
 &= \sqrt{\frac{\pi}{2}} \cdot \frac{1}{m} \mathbb{E}_{\delta, \delta', \sigma} \sup_T \left\| \mathbb{E}_g \sum_{i=1}^n |g_i| \sigma_i (\delta'_i - \delta_i) z_i^{(T)} z_i^{(T)*} \right\|
 \end{aligned}$$

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

$$\begin{aligned}
 \text{LHS} &= \mathbb{E}_{\delta} \sup_{\substack{T \subset [n] \\ |T| \leq k}} \left\| \overbrace{\mathbb{E}_{\delta'} \frac{1}{m} \sum_{i=1}^n \delta'_i z_i^{(T)} z_i^{(T)*}}^{l_T} - \frac{1}{m} \sum_{i=1}^n \delta_i z_i^{(T)} z_i^{(T)*} \right\| \\
 &\leq \frac{1}{m} \mathbb{E}_{\delta, \delta'} \sup_T \left\| \sum_{i=1}^n (\delta'_i - \delta_i) z_i^{(T)} z_i^{(T)*} \right\| \text{ (Jensen)} \\
 &= \sqrt{\frac{\pi}{2}} \cdot \frac{1}{m} \mathbb{E}_{\delta, \delta', \sigma} \sup_T \left\| \mathbb{E}_{\mathbf{g}} \sum_{i=1}^n |g_i| \sigma_i (\delta'_i - \delta_i) z_i^{(T)} z_i^{(T)*} \right\| \\
 &\leq \sqrt{2\pi} \cdot \frac{1}{m} \mathbb{E}_{\delta, \mathbf{g}} \sup_T \left\| \sum_{i=1}^n g_i \delta_i z_i^{(T)} z_i^{(T)*} \right\| \text{ (Jensen+triangle ineq)}
 \end{aligned}$$

Chaining without gaussians: RIP (Rudelson, Vershynin'06)

$$\begin{aligned}
 \text{LHS} &= \mathbb{E}_{\delta} \sup_{\substack{T \subset [n] \\ |T| \leq k}} \left\| \overbrace{\mathbb{E}_{\delta'} \frac{1}{m} \sum_{i=1}^n \delta'_i z_i^{(T)} z_i^{(T)*}}^{l_T} - \frac{1}{m} \sum_{i=1}^n \delta_i z_i^{(T)} z_i^{(T)*} \right\| \\
 &\leq \frac{1}{m} \mathbb{E}_{\delta, \delta'} \sup_T \left\| \sum_{i=1}^n (\delta'_i - \delta_i) z_i^{(T)} z_i^{(T)*} \right\| \text{ (Jensen)} \\
 &= \sqrt{\frac{\pi}{2}} \cdot \frac{1}{m} \mathbb{E}_{\delta, \delta', \sigma} \sup_T \left\| \mathbb{E}_{\mathbf{g}} \sum_{i=1}^n |g_i| \sigma_i (\delta'_i - \delta_i) z_i^{(T)} z_i^{(T)*} \right\| \\
 &\leq \sqrt{2\pi} \cdot \frac{1}{m} \mathbb{E}_{\delta, \mathbf{g}} \sup_T \left\| \sum_{i=1}^n g_i \delta_i z_i^{(T)} z_i^{(T)*} \right\| \text{ (Jensen+triangle ineq)} \\
 &\simeq \frac{1}{m} \mathbb{E}_{\delta} \mathbb{E}_{\mathbf{g}} \sup_{x \in B_2^{n,k}} \left| \sum_{i=1}^n g_i \delta_i \langle z_i, x \rangle^2 \right| \text{ (gaussian mean width!)}
 \end{aligned}$$

The End

June 22nd+23rd: workshop on concentration of measure /
chaining at Harvard, after STOC'16. Details+website forthcoming.