
Lactylation regulator expression and DDR protein fractions




RNA: boxes show median and IQR; whiskers extend to 1.5 x IQR. Points are sample records; dashed lines connect material-level means.

DDR: calculate 100 x GO-DDR proteins / detected proteins per source unit, then plot mean +/- SEM (62 bars). Singletons and unreplicated conditions have no error bar.

t: SEM across technical preparations; m: mean across cell models, no SEM; q: pooled QC source. Other errors describe the included source observations; conditions are listed in the data table.

* Shared RNA reference: HCT116 (3 columns), HK-2 (2 columns). 31 display columns correspond to 28 RNA references; reference cohorts and conditions are listed in the mapping table.

Lactate metabolism / KAT-family lactyltransferases and DDR protein fractions




RNA: boxes show median and IQR; whiskers extend to 1.5 x IQR. Points are sample records; dashed lines connect material-level means.

DDR: calculate 100 x GO-DDR proteins / detected proteins per source unit, then plot mean +/- SEM (62 bars). Singletons and unreplicated conditions have no error bar.

t: SEM across technical preparations; m: mean across cell models, no SEM; q: pooled QC source. Other errors describe the included source observations; conditions are listed in the data table.


* Shared RNA reference: HCT116 (3 columns), HK-2 (2 columns). 31 display columns correspond to 28 RNA references; reference cohorts and conditions are listed in the mapping table.

Lactate metabolism / AARS lactyltransferases and DDR protein fractions



RNA: boxes show median and IQR; whiskers extend to 1.5 x IQR. Points are sample records; dashed lines connect material-level means.
DDR: calculate 100 x GO-DDR proteins / detected proteins per source unit, then plot mean +/- SEM (62 bars). Singletons and unreplicated conditions have no error bar.
t: SEM across technical preparations; m: mean across cell models, no SEM; q: pooled QC source. Other errors describe the included source observations; conditions are listed in the data table.
* Shared RNA reference: HCT116 (3 columns), HK-2 (2 columns). 31 display columns correspond to 28 RNA references; reference cohorts and conditions are listed in the mapping table.

Delactylation-associated regulators and DDR protein fractions



RNA: boxes show median and IQR; whiskers extend to 1.5 x IQR. Points are sample records; dashed lines connect material-level means.

DDR: calculate 100 x GO-DDR proteins / detected proteins per source unit, then plot mean +/- SEM (62 bars). Singletons and unreplicated conditions have no error bar.

t: SEM across technical preparations; m: mean across cell models, no SEM; q: pooled QC source. Other errors describe the included source observations; conditions are listed in the data table.

* Shared RNA reference: HCT116 (3 columns), HK-2 (2 columns). 31 display columns correspond to 28 RNA references; reference cohorts and conditions are listed in the mapping table.