Please check the examination details belo	w before ente	ring your candidate information
Candidate surname		Other names
Centre Number Candidate Nu Pearson Edexcel Level		
Tuesday 20 June 202	23	
Afternoon	Paper reference	9MA0/32
Mathematics Advanced PAPER 32: Mechanics		
You must have: Mathematical Formulae and Statistical	Tables (Gre	een), calculator

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of g is required, take $g = 9.8 \,\mathrm{m \, s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 50. There are 6 questions.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1.	A car is initially at rest on a straight horizontal road. The car then accelerates along the road with a constant acceleration of $3.2\mathrm{ms}^{-2}$ Find		
	(a) the speed of the car after 5 s,(b) the distance travelled by the car in the first 5 s.	(1)	
		(2)	_
			_
			_
			_
			_
			_
			_
			_
			_
			_
			_



$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$
DO NOT WRITE IN THIS AREA
X 62X
\times
XXXXX
XXXX
\times
\times
\times
$\times ac$
X5X
$\otimes \times \times$
$\otimes \otimes \otimes$
$\times=\times$
\times
\times
$\times $
$\times\!\!\times\!\!\times\!\!\times$
\times
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

\times
XX 8 EX
$\times \times \times$
Ė
i Z
E IN T
TE IN T
ITE IN THIS AREA
RITE IN TH
VRITE IN TH
WRITEINT
WRITE IN TH
T WRITE IN TH
OT WRITE IN TH
JOT WRITE IN TH
NOT WRITE IN TH
NOTWR
O NOT WR
DONOTWR
A DO NOT WR
A DO NOT WR
EA DO NOT WR
EA DO NOT WR
AREA DO NOT WR
AREA DO NOT WR
AREA DO NOT WR
AREA DO NOT WR
IIS AREA DO NOT WR
IIS AREA DO NOT WR
IIS AREA DO NOT WR
UTHIS AREA DO NOT WR
UTHIS AREA DO NOT WR
UTHIS AREA DO NOT WR
UTHIS AREA DO NOT WR
TE IN THIS AREA DO NOT WR
TE IN THIS AREA DO NOT WR
RITE IN THIS AREA DO NOT WR
RITE IN THIS AREA DO NOT WR
RITE IN THIS AREA DO NOT WR
RITE IN THIS AREA DO NOT WR
OT WRITE IN THIS AREA DO NOT WR
OT WRITE IN THIS AREA DO NOT WR
OT WRITE IN THIS AREA DO NOT WR
NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
NOT WRITE IN THIS AREA DO NOT WR

Question 1 continued	
	(Total for Question 1 is 3 marks)



2.



Figure 1

A particle *P* has mass 5 kg.

The particle is pulled along a rough horizontal plane by a horizontal force of magnitude 28 N.

The only resistance to motion is a frictional force of magnitude F newtons, as shown in Figure 1.

(a) Find the magnitude of the normal reaction of the plane on P

(1)

The particle is accelerating along the plane at $1.4 \,\mathrm{m\,s}^{-2}$

(b) Find the value of F

(2)

The coefficient of friction between P and the plane is μ

(c) Find the value of μ , giving your answer to 2 significant figures.

(1)





Κ				۲		
K	3	ø	۲	ø	=	
	Ġ	ď				
K	А		ĸ.		m)	
	V	۲	7	7		
1	2	К	1	1		
				į		
5	d	ķ	d	Ģ	ď	
2	4	į		S	3	
3	d		4		Ž	
2	3					
2	4		4 10			
3	4		1 W W W	10.00		
3	4		AMM.			
3	4					
3	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì	ŀ	S	2	Š	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì			2)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	
	Ì)	

Question 2 continued			
(Total for Question 2 is 4 marks)			



3. At time t seconds, where $t \ge 0$, a particle P has velocity $\mathbf{v} \, \mathbf{m} \, \mathbf{s}^{-1}$ where

$$\mathbf{v} = (t^2 - 3t + 7)\mathbf{i} + (2t^2 - 3)\mathbf{j}$$

Find

(a) the speed of P at time t = 0

(3)

(b) the value of t when P is moving parallel to $(\mathbf{i} + \mathbf{j})$

(2)

(c) the acceleration of P at time t seconds

(2)

(d) the value of t when the direction of the acceleration of P is perpendicular to \mathbf{i}

(2)

×××××	
	Question 3 continued
EA	
A B	
Ŧ I	
l	
ı	
ı	
I	

(Total for Question 3 is 9 marks)
_



4. [In this question, i and j are horizontal unit vectors and position vectors are given relative to a fixed origin O]

A particle P is moving on a smooth horizontal plane.

The particle has constant acceleration $(2.4i + j) \text{ m s}^{-2}$

At time t = 0, P passes through the point A.

At time t = 5 s, P passes through the point B.

The velocity of P as it passes through A is $(-16\mathbf{i} - 3\mathbf{j}) \,\mathrm{m \, s}^{-1}$

(a) Find the speed of P as it passes through B.

(4)

The position vector of A is $(44\mathbf{i} - 10\mathbf{j})$ m.

At time t = T seconds, where T > 5, P passes through the point C.

The position vector of C is $(4\mathbf{i} + c\mathbf{j})$ m.

(b) Find the value of *T*.

(3)

(c) Find the value of c.

(3)

\otimes		
	$\times\!\!\!\times$	8
×	\times	8
\otimes	\times	ξ
×	$\times\!\!\times$	ζ
\otimes	$\times\!\!\times$	ζ
\otimes	\approx	3
\otimes	AREA	Š
×	ш	K
\otimes	ÓĊ	Š
\times	₫	Š
\otimes	LO	Š
×	*	K
\otimes		8
\otimes		8
\otimes	Z	8
\otimes		8
X	ш	Š
×	X	8
\otimes	α	
\otimes	2	3
\otimes	NOTWRITEINTHIS	Š
\otimes	\boxtimes	Š
×	2	Ž
\otimes		8
\approx	0	8
X	00	Š
X		ξ
X	\otimes	ζ
×	\otimes	ζ
\otimes	\times	ζ
\otimes	\otimes	Š
\otimes	\times	Š
8	\otimes	Š
\otimes	$\times\!\!\times$	Š
\otimes	\bowtie	8
\otimes	\otimes	8
\otimes	\bowtie	8
X	\approx	S
	ST.	Š
\otimes	W	Š
\otimes	AREA	Š
\otimes	S.	Š
\otimes	S	Š
X		2
$\stackrel{\times}{\times}$	Į	
	E	
	I	
	工厂之山	
	WEITHER	
	OTWRITEINTE	
	TEN MEN MAKEN	
	ONOTWRITE IN THIS	
	Õ	
	DONOTWRITEINTH	
	Õ	
	Õ	
	Õ	
	Õ	
	Õ	
	Õ	
	Õ	
	Õ	
	Õ	
	Õ	
	00	
	00	
	REA	
	REA	
	SAREA	
	SAREA	
	HIS AREA DO	
	THIS AREA	
	THIS AREA	
	IN THIS AREA	
	TE IN THIS AREA	
	TE IN THIS AREA	
	TE IN THIS AREA	
	WRITE IN THIS AREA	
	WRITE IN THIS AREA	
	WRITE IN THIS AREA	
	NOT WRITE IN THIS AREA	
	NOT WRITE IN THIS AREA	
	WRITE IN THIS AREA	

Question 4 continued



Question 4 continued

Question 4 continued	
(Total for Question 4 is 10 marks)	



5.




Figure 2

A small ball is projected with speed $28\,\mathrm{m\,s}^{-1}$ from a point O on horizontal ground.

After moving for T seconds, the ball passes through the point A.

The point A is 40 m horizontally and 20 m vertically from the point O, as shown in Figure 2.

The motion of the ball from O to A is modelled as that of a particle moving freely under gravity.

Given that the ball is projected at an angle α to the ground, use the model to

(a) show that
$$T = \frac{10}{7\cos\alpha}$$

(2)

(b) show that $\tan^2 \alpha - 4 \tan \alpha + 3 = 0$

(5)

(c) find the greatest possible height, in metres, of the ball above the ground as the ball moves from O to A.

(3)

The model does not include air resistance.

(d) State one other limitation of the model.

(1)

Question 5 continued



Question 5 continued		

Question 5 continued	
	(Total for Question 5 is 11 marks)



6.




Figure 3

A rod AB has mass M and length 2a.

The rod has its end A on rough horizontal ground and its end B against a smooth vertical wall.

The rod makes an angle θ with the ground, as shown in Figure 3.

The rod is at rest in limiting equilibrium.

(a) State the direction (left or right on Figure 3 above) of the frictional force acting on the rod at A. Give a reason for your answer.

(1)

The magnitude of the normal reaction of the wall on the rod at *B* is *S*.

In an initial model, the rod is modelled as being uniform.

Use this initial model to answer parts (b), (c) and (d).

(b) By taking moments about A, show that

$$S = \frac{1}{2} Mg \cot \theta \tag{3}$$

The coefficient of friction between the rod and the ground is μ

Given that $\tan \theta = \frac{3}{4}$

(c) find the value of μ

(5)

(d) find, in terms of M and g, the magnitude of the resultant force acting on the rod at A.

(3)

In a new model, the rod is modelled as being **non-uniform**, with its centre of mass closer to B than it is to A.

A new value for S is calculated using this new model, with $\tan \theta = \frac{3}{4}$

(e) State whether this new value for S is larger, smaller or equal to the value that S would take using the initial model. Give a reason for your answer.

(1)

Question 6 continued



Question 6 continued		

Duestion 6 continued	
	-
	_
	_
	-
	-
	-
	_
	-
	-
	-
	-
	-
	-
	-
	_
	-
	-
	-
	-
	-
	-
	-
	_
	-
	-
	-



Question 6 continued	
	(Total for Question 6 is 13 marks)
	TOTAL FOR MECHANICS IS 50 MARKS