

The mirror map: a Fibonacci age law for an injective relative of the $3x + 1$ map, and numerical evidence for an open question of Reyes Jiménez

Darcy Thomas*

September 2026. Draft — not yet checked against all primary sources; see Section 6.

Abstract

Replacing the halving step $n \mapsto n/2$ of the $3x + 1$ map by $n \mapsto n - |n/2|$ leaves positive integers unchanged and turns the map on negative integers into the injective, magnitude-increasing map $m \mapsto 3m - 1$ (m odd), $m \mapsto 3m/2$ (m even), which we call the *mirror map*: it exchanges each factor 2 for a factor 3 instead of discarding it. Its inverse is a terminating process, giving every integer an *age*. We prove that the ages are governed by Fibonacci numbers: $\Pr(\text{age} \geq t) = F_{t+3}/(2 \cdot 3^t)$, and more generally $F_{t+3}/(2q^t)$ when 3 is replaced by any odd $q \geq 3$. The Fibonacci structure comes from the prime 2 only. We show that a recent forward Fibonacci count of Reyes Jiménez is the 2-adic counterpart of this law: avoiding 4 (mod 6) is the parity condition “no two consecutive halvings”, which makes that count a direct corollary of Winkler’s finite-state parity principle and identifies the counted set as the integers whose initial Syracuse exponents are all at most 2. That paper asks whether every $n > 1$ eventually has two consecutive halvings. We observe that the $3x + 1$ conjecture implies this, that it fails for negative integers, and we verify it exhaustively for all $n < 2^{56}$ with an algorithm whose cost is $O(\varphi^B)$ rather than $O(2^B)$. Beyond an integer’s own binary digits, the measured per-step survival probability is 0.809017, against $\varphi/2 = 0.809017$. The same method, run backwards in base 3, verifies a conjecture of Banerji for all odd $n < 2 \cdot 3^{32}$.

1 The map

Let $F(n) = 3n + 1$ for odd n and $F(n) = n - |n/2|$ for even n . For $n > 0$ this is the Collatz map. For $n < 0$ the even step is $n \mapsto 3n/2$, so on magnitudes $m = |n|$ the map is

$$G(m) = 3m - 1 \quad (m \text{ odd}), \quad G(m) = 3m/2 \quad (m \text{ even}). \quad (1)$$

We call G the *mirror map*. The name records two things: G is the Collatz map reflected through zero, and its structure mirrors that of the Collatz map, with the primes 2 and 3 exchanged and the direction of time reversed (Sections 3–5). For odd $q \geq 3$ and $c = \pm 1$ put $G_{q,c}(m) = qm + c$ for odd m and $qm/2$ for even m , so that $G = G_{3,-1}$; we use the same name for this family. The orbit of 1 under G is 1, 2, 3, 8, 12, 18, 27, 80, 120, 180, 270, 405, 1214, ...

The closest named relative is the map $x \mapsto (3x + 1)/2$ (x odd), $3x/2$ (x even) from Mahler’s Z -number problem [11]. The map (1) differs in the odd branch, and that difference is what makes it injective.

*musicalmathmind@gmail.com

2 Elementary properties

Proposition 1 $G_{q,c}$ is injective and $G_{q,c}(m) > m$ for all $m \geq 1$. Its image is $\{m : q \mid m\} \cup \{m \text{ even} : m \equiv c \pmod{q}\}$, of density $3/(2q)$. For $q = 3$, $c = -1$ exactly half of the positive integers have no predecessor, namely those with $m \equiv 1 \pmod{3}$ or $m \equiv 5 \pmod{6}$.

Proof. $qm + c > m$ and $qm/2 > m$. Odd m map to even values congruent to c modulo q ; even m map to multiples of q . These two sets are disjoint, each branch is injective, and every element of either set is attained. $\square$

Proposition 2 (Valuation transfer) Let b be odd and write $qb + c = 2^a u$ with u odd. The next odd value on the $G_{q,c}$ -orbit of b is $q^a u$, and $\gcd(u, q) = 1$. Hence its q -adic valuation equals $v_2(qb + c)$.

Proof. Each of the a even steps replaces one factor 2 by one factor q . Since $qb + c \equiv c \pmod{q}$, u is prime to q . $\square$

Thus G retains, as a power of 3, exactly the power of 2 that the Syracuse map discards. In terms of prime factorisations, the even step of G exchanges a factor 2 for a factor 3 and preserves the number Ω of prime factors, whereas halving decreases Ω by one.

3 The age law

Backward steps from m are: **A**, $m \mapsto 2m/q$, available iff $q \mid m$; and **B**, $m \mapsto (m - c)/q$, available iff m is even and $m \equiv c \pmod{q}$. At most one is available. Let $\text{age}(m)$ be the number of backward steps that can be taken from m , and let the *age word* of m record them, most recent first. The last value reached, which has no predecessor, is the *seed* of m . For G : $\text{age}(27) = 6$, with word AAABAB and seed 1.

Write $F_1 = F_2 = 1$ for the Fibonacci numbers.

Theorem 3 Let $q \geq 3$ be odd, $c = \pm 1$ and $t \geq 1$.

1. The age words are exactly the words over $\{A, B\}$ with no factor BB.
2. $\min(\text{age}(m), t)$ depends only on $m \bmod 2q^t$.
3. The number of residues modulo $2q^t$ with $\text{age} \geq t$ is F_{t+3} . Consequently the natural density of $\{m : \text{age}(m) \geq t\}$ is $F_{t+3}/(2q^t)$.

Proof. (i) After a step B the value $(m - c)/q$ is odd, since $m - c$ is odd and q is odd; but B requires an even value. So B never follows B. After a step A the value $2m/q$ is even, so the parity requirement of B is automatic there. Hence the only parity condition in an entire backward walk is on m itself, and only if the first step is B.

(ii) Each backward step has the form $m \mapsto (um + v)/q$ with $u \in \{1, 2\}$, a unit modulo q . After i steps the ancestor is $(Um + V)/q^i$ with U a unit modulo q . The condition for step $i + 1$, that the ancestor be congruent to 0 or to c modulo q , therefore selects exactly one of the q lifts of $m \bmod q^i$ to $m \bmod q^{i+1}$. By induction every BB-free word of length t occurs, determines $m \bmod q^t$ uniquely, and distinct words determine distinct residues.

(iii) By the Chinese remainder theorem a word beginning with A corresponds to two residues modulo $2q^t$ (the parity of m is free), and a word beginning with B to one (m even). There are F_{t+1} BB-free words of length t beginning with A and F_t beginning with B, and $2F_{t+1} + F_t = F_{t+3}$. $\square$

Corollary 4 1. The mean age is $\sum_{t \geq 1} F_{t+3}/(2q^t) = \frac{1}{2}(\frac{2+x}{1-x-x^2} - 2)$ at $x = 1/q$; for $q = 3$ it equals $11/10$.

2. The language of age words has factor complexity $p(n) = F_{n+2}$: it is the full golden-mean shift.
3. Every positive integer has unique coordinates (seed, age word), and the positive integers are a disjoint union of $G_{q,c}$ -orbits of seeds.
4. Forward, the run of even steps after an odd b has length $v_2(qb + c)$, which is geometric of ratio $1/2$ with respect to density. Backward, a run of steps A continues iff q divides the ancestor, which is geometric of ratio $1/q$.

The Fibonacci numbers here are produced by the prime 2 alone (an odd value cannot take the odd-type step); q enters only through the denominator.

Checks. Part (iii) was confirmed by exhaustive count for $q \in \{3, 5, 7, 9, 11\}$ and both signs of c , with $t \leq 9$ for $q = 3$; the mean age over $m \leq 2 \cdot 3^{11}$ is 1.09999; the run-length tails in Corollary 4(iv) agree to four decimals.

4 The forward counterpart

Let $T(n) = n/2$ for even n and $(3n + 1)/2$ for odd n . Reyes Jiménez [13, Theorem 4.7] proves that exactly F_{m+1} odd integers $n \in \{1, \dots, 2^m\}$ satisfy $T^i(n) \not\equiv 4 \pmod{6}$ for $1 \leq i \leq m - 1$, using the spectral radius φ of the modulo-6 transition graph with the vertex 4 removed.

Proposition 5 For odd n , some $T^i(n)$ with $1 \leq i \leq m - 1$ is congruent to 4 modulo 6 if and only if the parity word of n of length m contains 00.

Proof. After an odd step the value is $\equiv 2 \pmod{3}$. Halving such a value gives a value $\equiv 1 \pmod{3}$; values $\equiv 1 \pmod{3}$ arise in no other way, and no value after the first step is divisible by 3. A value is $\equiv 4 \pmod{6}$ iff it is $\equiv 1 \pmod{3}$ and even, that is, iff it was produced by a halving and the next step is a halving. $\square$

By the bijection of Terras and Everett [15, 4] between parity words of length m and residues modulo 2^m , words that begin with 1 and avoid 00 correspond to F_{m+1} odd residues. This reproves [13, Theorem 4.7] and identifies the counted set: the n whose initial Syracuse exponents are all at most 2. We verified the equality of the two sets for $m \leq 14$.

Remark 6 The counting step is an instance of Winkler’s finite-state parity principle [17, Proposition 5.1]: any condition on the first k parity symbols recognised by a finite automaton is a transfer-matrix count of residue classes modulo 2^k ; his own theorems use the forbidden factor 11. Winkler remarks that the theorem of [13] “is not a direct corollary” of that principle, because its states record residues modulo 6 rather than parity information. Proposition 5 removes the obstruction: avoiding $4 \pmod{6}$ is a parity condition, the forbidden factor 00, so [13, Theorem 4.7] is a direct corollary of [17, Proposition 5.1]. What is contributed here is only the translation.

	forward [13]	backward (Theorem 3)
forbidden factor	00	BB
digit bijection	parity word $\leftrightarrow n \bmod 2^m$	age word $\leftrightarrow m \bmod q^t$
count	F_{m+1} of 2^{m-1}	F_{t+3} of $2q^t$
status	selects a null set of orbits	holds for every integer

5 An open question of Reyes Jiménez

In [13, Section 5] it is asked whether every odd $n > 2$ eventually meets $4 \pmod{6}$, with the remark that an explicit bound $m_0(n) \leq f(n)$ on the first such level would settle it. By Proposition 5 this is:

Question Q *Does every integer $n > 1$ have two consecutive halvings in its T -orbit? Equivalently, is some Syracuse iterate of every odd $n > 1$ congruent to 5 modulo 8?*

Remark 7 1. The $3x+1$ conjecture implies a positive answer. *The odd Syracuse predecessors of 1 other than 1 are $(4^j - 1)/3$ with $j \geq 2$, reached with exponent $2j \geq 4$.*

2. The statement fails for negative integers: -1 is fixed with exponent 1, and $-5 \mapsto -7 \mapsto -5$ uses exponents 1, 2. *Any proof must therefore use positivity.*

3. *The 2-adic integers whose parity sequence avoids 00 form a set of Hausdorff dimension $\log_2 \varphi \approx 0.694$, because the parity-sequence map is a 2-adic isometry conjugating T to the shift [10].*

4. *An orbit giving a negative answer is either a non-trivial cycle all of whose exponents are at most 2, or is unbounded. Inside the family the exponent is 1 for $n \equiv 3, 7 \pmod{8}$ and 2 for $n \equiv 1 \pmod{8}$, so typical members grow; Question Q is of divergence type. The bounds on m -cycles of Simons-de Weger [14] and Hercher [7] do not apply, since such a cycle may have arbitrarily many local minima.*

Algorithm

The residues r modulo 2^j whose first j steps avoid 00 form a tree with F_{j+1} nodes at level j . With each node carry $x = T^j(r)$ and 3^s , where s is the number of odd steps so far. Replacing r by $r + 2^j$ replaces x by $x + 3^s$, which has the opposite parity, so both children are obtained in $O(1)$: the odd child always survives, and the even child survives iff the previous step was not a halving. Walk the tree to depth B and finish each of the F_{B+1} leaves, which are genuine integers $r < 2^B$ with $T^B(r)$ already known, by direct iteration. Integers that leave the family earlier are pruned. The search is exhaustive for $n < 2^B$ at cost $O(\varphi^B)$.

Results

B	leaves F_{B+1}	time	longest survival	attained at n	steps/ $\log_2 n$
32	3 524 578	< 0.1 s	100	1 548 635 775	3.28
40	165 580 141	0.2 s	129	971 145 319 023	3.24
48	7 778 742 049	10.5 s	152	19 914 105 817 593	3.44
56	365 435 296 162	525 s	173	66 107 737 450 688 865	3.10

(Rust, 32 threads, 128-bit arithmetic with an overflow guard that never triggered; counts and records agree with an independent Python implementation for $B = 24, 32, 40$.)

Question Q has a positive answer for every n below the largest bound in the table. A model in which each further step is survived with probability $\varphi/2$ predicts record survival times $\log_2 n / \log_2(2/\varphi) \approx 3.27 \log_2 n$, and the observed records lie on that line. We conjecture $m_0(n) \leq 4 \log_2 n$ for all sufficiently large n .

Self-generated digits

For $B = 56$, among the $3.65 \cdot 10^{11}$ integers that survive their own 56 binary digits, the conditional probability of surviving one more step is 0.809017, 0.809017, 0.809012, 0.809021 at levels 56, 64, 72, 80; and $\varphi/2 = 0.809017$. Beyond level B the parity is no longer read from the digits of n but from digits produced by the iteration, and these leave the family at exactly the rate at which free digits would.

The backward counterpart

Banerji [3] conjectured that backward Syracuse iteration restricted to exponents at most 2 always reaches a multiple of 3. On odd n this iteration is the deterministic map $b(n) = (2n - 1)/3$ if $n \equiv 2 \pmod{3}$, $b(n) = (4n - 1)/3$ if $n \equiv 1 \pmod{3}$, with $n = 1$ fixed. The same algorithm applies with base-3 digits: replacing r by $r + 2 \cdot 3^j d$ changes $b^j(r)$ by a power of 2 times d , so exactly one of the three lifts dies, and the survivors of t steps are 2^t of the 3^t odd residues modulo $2 \cdot 3^t$. The conjecture holds for every odd $1 < n < 2 \cdot 3^{32} \approx 2^{51.7}$; the longest survival is 83 steps, at $n = 116\,393\,404\,689\,949$; the per-step survival probability is 0.66667 against $2/3$. The exceptional set is 3-adic, of dimension $\log_3 2$.

6 Related work, and what has been checked

Forward Fibonacci counts for the $3x+1$ map are known: the set of itineraries of the unaccelerated map is the golden-mean shift [6]; Winkler [17] proves Fibonacci counts of residue classes modulo 2^{r+2} defined by a local cut time, via the Terras–Everett bijection and the forbidden factor 11, and formulates the general finite-state principle used in Remark 6; and [13] is discussed above. All of these are forward counts modulo powers of 2. Winkler leaves the extension to maps $n \mapsto (qn + 1)/2$ for separate investigation; Theorem 3 is a different statement (backward, modulo powers of q), but it does hold for all odd q . Albert, Gudmundsson and Ulfarsson [2] enumerate Collatz-induced permutations by Fibonacci numbers. That backward admissibility is decided 3-adically is the mechanism of Wirsching [16], for the map T , where runs of halvings are unrestricted and the counts are not Fibonacci. Step A of the inverse of G is the digit-0 edge of the base-3/2 number system of Akiyama, Frougny and Sakarovitch [1]; see also [5] for negative integers and 3-adic convergence. Merlini and Sala [12] obtain a Fibonacci-type recursion as a heuristic growth rate of the inverse tree.

We text-searched both annotated bibliographies of Lagarias [8, 9] and found no entry for the map (1), for an injective variant of this form, or for a Fibonacci count of backward residues. The sequence 1, 2, 3, 8, 12, 18, 27, 80, ... is not in the OEIS. We have read [17] (v3) and the introduction of [5]. **We have not read the primary texts of [16], [12], [3] or [6],** only annotations and abstracts, and [13] only in part.

Theorem 3 and Proposition 5 are elementary, and every ingredient is known. The prudent description is: an elementary backward, q -adic companion to the known forward Fibonacci counts, obtained by Wirsching’s mechanism applied to an injective relative of Mahler’s map.

7 Questions

1. Is there a single bijection explaining both Fibonacci counts, exchanging the roles of 2 and q ?
2. Does a single G -orbit have the Markov statistics ($\pi_A = 2/3$, $P(B | A) = 1/2$)? Three orbits of 200 000 steps pass a χ^2 test; a proof appears as hard as the distribution of $(3/2)^k$ modulo 1.
3. Can Question Q be settled for cycles alone?

Acknowledgements and reproducibility

Code, tests and raw results are in the author’s repository (`collatz/pinball/`, `scripts/`). The computations, the literature search and the drafting of this note were carried out with substantial assistance from Claude (Anthropic); all statements were checked by computation as described.

References

- [1] S. Akiyama, C. Frougny, J. Sakarovitch, Powers of rationals modulo 1 and rational base number systems, *Israel J. Math.* 168 (2008), 53–91.
- [2] M. H. Albert, B. Gudmundsson, H. Ulfarsson, Collatz meets Fibonacci, arXiv:1404.3054.
- [3] R. B. Banerji, Some properties of the $3n + 1$ function, *Cybernetics and Systems* 27 (1996), 473–486. [Known to us only through the annotation in [8].]
- [4] C. J. Everett, Iteration of the number-theoretic function $f(2n) = n$, $f(2n + 1) = 3n + 2$, *Adv. Math.* 25 (1977), 42–45.
- [5] C. Frougny, K. Klouda, Rational base number systems for p -adic numbers, *RAIRO Theor. Inform. Appl.* 46 (2012), 87–106.
- [6] D.-A. German, The golden mean shift is the set of $3x + 1$ itineraries, NKS 2004 conference.
- [7] C. Hercher, There are no Collatz m -cycles with $m \leq 91$, *J. Integer Seq.* 26 (2023), 23.3.5.
- [8] J. C. Lagarias, The $3x + 1$ problem: an annotated bibliography (1963–1999), arXiv:math/0309224.
- [9] J. C. Lagarias, The $3x + 1$ problem: an annotated bibliography, II (2000–2009), arXiv:math/0608208.
- [10] J. C. Lagarias, The $3x + 1$ problem and its generalizations, *Amer. Math. Monthly* 92 (1985), 3–23.
- [11] K. Mahler, An unsolved problem on the powers of $3/2$, *J. Austral. Math. Soc.* 8 (1968), 313–321.
- [12] D. Merlini, N. Sala, On the Fibonacci’s attractor and the long orbits in the $3n + 1$ problem, *Int. J. Chaos Theory Appl.* 4 (1999).
- [13] M.-A. Reyes Jiménez, A Fibonacci theorem for Collatz trajectories via modular graph structure, arXiv:2606.02621 (2026).

- [14] J. Simons, B. de Weger, Theoretical and computational bounds for m -cycles of the $3n + 1$ problem, *Acta Arith.* 117 (2005), 51–70.
- [15] R. Terras, A stopping time problem on the positive integers, *Acta Arith.* 30 (1976), 241–252.
- [16] G. J. Wirsching, *The Dynamical System Generated by the $3n + 1$ Function*, Lecture Notes in Math. 1681, Springer, 1998.
- [17] M. Winkler, Fibonacci enumeration of parity blocks in Collatz trajectories, arXiv:1412.0519v3 (10 August 2026).