

Exponential Families

version: 2018-01-23 · 21:35:08

Exponential Families

- **Definition:** If $\mathbf{Y} \sim f(\mathbf{y} | \boldsymbol{\theta})$, $\boldsymbol{\theta} \in \mathbb{R}^d$, then \mathbf{Y} is said to belong to an **exponential family** if

$$f(\mathbf{y} | \boldsymbol{\theta}) = \exp \{ \mathbf{T}'(\mathbf{y}) \boldsymbol{\eta} - \Psi(\boldsymbol{\eta}) \} \cdot h(\mathbf{y}),$$

where

- $\boldsymbol{\eta} = \boldsymbol{\eta}(\boldsymbol{\theta}) \in \mathbb{R}^d$ are the *natural parameters*.
($\boldsymbol{\eta}$ must have the same dimension as $\boldsymbol{\theta}$ for upcoming results to hold.)
- $\mathbf{T} = \mathbf{T}(\mathbf{y})$ are the *sufficient statistics*.
- $\Psi(\boldsymbol{\eta})$ is called the log-partition function, or sometimes the cumulant-generating function.
- **Natural Parametrization:** Since each value of $\boldsymbol{\theta}$ defines a different PDF, $\boldsymbol{\eta}(\boldsymbol{\theta})$ *must* be a bijection. Therefore, we might as well parametrize the exponential family by $\boldsymbol{\eta}$, in which case $f(\mathbf{y} | \boldsymbol{\eta})$ is said to be in its *canonical form*.

Examples

Binomial Distribution

$$Y \sim \text{Bin}(n, \rho) \implies$$

$$\begin{aligned} f(y | n) &= \binom{n}{y} \rho^y (1 - \rho)^{n-y} \\ &= \exp \left\{ y \cdot \underbrace{\log \left(\frac{\rho}{1 - \rho} \right)}_{\eta} - \underbrace{[-n \log(1 - \rho)]}_{\Psi(\eta)} \right\} \cdot \binom{n}{y} \end{aligned}$$

Examples

Multivariate Normal Distribution

$$\mathbf{Y} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \implies$$

$$\begin{aligned} f(\mathbf{y} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma}) &= \exp \left\{ -\frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu}) - \frac{1}{2} \log |\boldsymbol{\Sigma}| \right\} \cdot \underbrace{h(\mathbf{y})}_{(2\pi)^{d/2}} \\ &= \exp \left\{ -\frac{1}{2} \left[\underbrace{\text{tr}(\boldsymbol{\Sigma}^{-1} \mathbf{y} \mathbf{y}')}_{\text{vec}(\boldsymbol{\Sigma}^{-1})' \text{vec}(\mathbf{y} \mathbf{y}')} - 2\mathbf{y}'[\boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}] + \boldsymbol{\mu}' \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu} + \log |\boldsymbol{\Sigma}| \right] \right\} h(\mathbf{y}) \end{aligned}$$

\implies

$$\mathbf{T} = \left(-\frac{1}{2}\mathbf{y}\mathbf{y}', \mathbf{y}\right), \quad \boldsymbol{\eta} = (\boldsymbol{\Sigma}^{-1}, \boldsymbol{\Sigma}^{-1}\boldsymbol{\mu}), \quad \Psi(\boldsymbol{\eta}) = -\frac{1}{2}(\boldsymbol{\mu}' \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu} + \log |\boldsymbol{\Sigma}|).$$

(Some redundancy since $\mathbf{y}\mathbf{y}'$ and $\boldsymbol{\Sigma}^{-1}$ are symmetric matrices, but formulas get complicated)

Examples

► **Model:** $\mathbf{Y} \sim f(\mathbf{y} | \boldsymbol{\eta}) = \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y}), \quad \mathbf{T} = \mathbf{T}(\mathbf{y}).$

► **Exponential families:**

Poisson, Gamma (and Exponential), Multinomial (and Binomial),
Negative-Binomial (and Geometric), Dirichlet (and Beta), Wishart (and
Chi-Square).

► **Not Exponential families:**

Student- t (and Cauchy), Weibull, Unif(0, θ).

Moments of Sufficient Statistics

- ▶ **Exponential Family:** $\mathbf{Y} \sim f(\mathbf{y} | \boldsymbol{\eta}) = \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y})$, $\mathbf{T} = \mathbf{T}(\mathbf{y})$.
- ▶ **Expectation of \mathbf{T} :**

$$\begin{aligned} \text{(since RHS is a PDF)} \quad 1 &= \int \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y}) \, d\mathbf{y} \\ \text{(take } \frac{\partial}{\partial \boldsymbol{\eta}} \text{ on both sides)} \quad \mathbf{0} &= \frac{\partial}{\partial \boldsymbol{\eta}} \int \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y}) \, d\mathbf{y} \\ &= \int \frac{\partial}{\partial \boldsymbol{\eta}} \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y}) \, d\mathbf{y} \\ &= \int [\mathbf{T} - \nabla \Psi(\boldsymbol{\eta})]f(\mathbf{y} | \boldsymbol{\eta}) \, d\mathbf{y} \\ &\quad \underbrace{\int \mathbf{T} \cdot f(\mathbf{y} | \boldsymbol{\eta}) \, d\mathbf{y}}_{=E[\mathbf{T} | \boldsymbol{\eta}]} = \nabla \Psi(\boldsymbol{\eta}) \underbrace{\int f(\mathbf{y} | \boldsymbol{\eta}) \, d\mathbf{y}}_{=1} \end{aligned}$$

$$\implies E[\mathbf{T} | \boldsymbol{\eta}] = \nabla \Psi(\boldsymbol{\eta}).$$

Moments of Sufficient Statistics

► **Exponential Family:** $\mathbf{Y} \sim f(\mathbf{y} | \boldsymbol{\eta}) = \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y})$, $\mathbf{T} = \mathbf{T}(\mathbf{y})$.

► **Variance of \mathbf{T} :**

$$1 = \int \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y}) d\mathbf{y}$$

$$\mathbf{0} = \frac{\partial}{\partial \boldsymbol{\eta}} \int \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y}) d\mathbf{y}$$

$$= \int [\mathbf{T} - \nabla \Psi(\boldsymbol{\eta})]f(\mathbf{y} | \boldsymbol{\eta}) d\mathbf{y}$$

$$\text{(take } \frac{\partial}{\partial \boldsymbol{\eta}} \text{ on both sides again)} \quad = \int \frac{\partial}{\partial \boldsymbol{\eta}} [\mathbf{T} - \nabla \Psi(\boldsymbol{\eta})]f(\mathbf{y} | \boldsymbol{\eta}) d\mathbf{y}$$

$$\nabla^2 \Psi(\boldsymbol{\eta}) \int f(\mathbf{y} | \boldsymbol{\eta}) d\mathbf{y} = \int [\mathbf{T} - \underbrace{\nabla \Psi(\boldsymbol{\eta})}_{E[\mathbf{T} | \boldsymbol{\eta}]}][\mathbf{T} - \nabla \Psi(\boldsymbol{\eta})]' f(\mathbf{y} | \boldsymbol{\eta}) d\mathbf{y}$$

$$\implies \text{var}[\mathbf{T} | \boldsymbol{\eta}] = \nabla^2 \Psi(\boldsymbol{\eta}) \implies \nabla^2 \Psi(\boldsymbol{\eta}) \text{ is positive definite.}$$

Inference

- ▶ **Data:** $\mathbf{Y} = (\mathbf{Y}_1, \dots, \mathbf{Y}_n)$, $\mathbf{Y}_i \stackrel{\text{iid}}{\sim} \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y})$.
- ▶ **Loglikelihood:** $\ell(\boldsymbol{\eta} | \mathbf{Y}) = n[\bar{\mathbf{T}}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})]$, where $\bar{\mathbf{T}} = \frac{1}{n} \sum_{i=1}^n \mathbf{T}(\mathbf{Y}_i)$.
- ▶ **Score function:** $\nabla \ell(\boldsymbol{\eta} | \mathbf{Y}) = n[\bar{\mathbf{T}} - \nabla \Psi(\boldsymbol{\eta})]$
 \implies MLE satisfies $\nabla \Psi(\hat{\boldsymbol{\eta}}) = \bar{\mathbf{T}}$.
- ▶ **Expected Fisher Information:**

$$\mathcal{I}(\boldsymbol{\eta}) = E[-\nabla^2 \ell(\boldsymbol{\eta} | \mathbf{Y})] = n E[\nabla^2 \Psi(\boldsymbol{\eta})] = n \nabla^2 \Psi(\boldsymbol{\eta}).$$

\implies Asymptotic theory $\hat{\boldsymbol{\eta}} \approx \mathcal{N}(\boldsymbol{\eta}_0, \mathcal{I}(\boldsymbol{\eta}_0)^{-1})$ is more effectively applied in practice since Observed Fisher Information is $\hat{\mathcal{I}} = \mathcal{I}(\hat{\boldsymbol{\eta}}) = n \nabla^2 \Psi(\hat{\boldsymbol{\eta}})$.

(usually expectation can't be calculated analytically)

- ▶ **Question:** How to compute MLE $\hat{\boldsymbol{\eta}}$?

Newton-Raphson Method

- **Problem:** Find a minimum of $f : \mathbb{R}^d \rightarrow \mathbb{R}$.
- **Quadratic case:** $f(\mathbf{x}) = \mathbf{x}'\mathbf{A}\mathbf{x} - 2\mathbf{b}'\mathbf{x} + c$, with $\mathbf{A}_{d \times d}$ is positive definite.
(Using Cholesky $\mathbf{A} = \mathbf{L}\mathbf{L}'$, show that \mathbf{A}^{-1} exists and is +ve definite)
- *Multivariate complete-the-square:*

$$\begin{aligned} f(\mathbf{x}) &= \mathbf{x}'\mathbf{A}\mathbf{x} - 2\underbrace{\mathbf{b}'\mathbf{A}^{-1}}_{\boldsymbol{\mu}'}\mathbf{A}\mathbf{x} + c \\ &= \underbrace{(\mathbf{x} - \boldsymbol{\mu})'\mathbf{A}(\mathbf{x} - \boldsymbol{\mu})}_{\mathbf{x}'\mathbf{A}\mathbf{x} - 2\boldsymbol{\mu}'\mathbf{x} + \boldsymbol{\mu}'\mathbf{A}\boldsymbol{\mu}} - \boldsymbol{\mu}'\mathbf{A}\boldsymbol{\mu} + c, \end{aligned}$$

\implies Unique minimum of $f(\mathbf{x})$ is $\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$.

Newton-Raphson Method

- **Problem:** Find a minimum of $f : \mathbb{R}^d \rightarrow \mathbb{R}$.
- **Non-Quadratic case:** Iterative method.
 - *Initial guess:* \mathbf{x}_0
 - *Iterations:* At step $n + 1$, find 2nd order Taylor expansion of $f(\mathbf{x})$ around $\mathbf{x} = \mathbf{x}_n$:

$$\begin{aligned} f(\mathbf{x}) &\approx f(\mathbf{x}_n) + \underbrace{\mathbf{g}_n'}_{\nabla f(\mathbf{x}_n)'} (\mathbf{x} - \mathbf{x}_n) + \frac{1}{2} (\mathbf{x} - \mathbf{x}_n)' \underbrace{\mathbf{H}_n}_{\nabla^2 f(\mathbf{x}_n)} (\mathbf{x} - \mathbf{x}_n) \\ &= \frac{1}{2} [\mathbf{x}' \mathbf{H}_n \mathbf{x} - 2(\mathbf{H}_n \mathbf{x}_n - \mathbf{g}_n)' \mathbf{x}] + \text{const} \\ &= \frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})' \mathbf{H}_n (\mathbf{x} - \boldsymbol{\mu}) + \text{const}, \quad \boldsymbol{\mu} = \mathbf{H}_n^{-1} (-\mathbf{g}_n + \mathbf{H}_n \mathbf{x}_n) \\ &\quad = \mathbf{x}_n - \mathbf{H}_n^{-1} \mathbf{g}_n. \end{aligned}$$

$$\implies \text{Let } \mathbf{x}_{n+1} = \mathbf{x}_n - \mathbf{H}_n^{-1} \mathbf{g}_n = \mathbf{x}_n - [\nabla^2 f(\mathbf{x}_n)]^{-1} \nabla f(\mathbf{x}_n).$$

- **Stopping Condition:** Algorithm terminates when $\max_{1 \leq i \leq d} \frac{|x_{n,i} - x_{n-1,i}|}{|x_{n,i} + x_{n-1,i}|} < \varepsilon$.


Convex Functions

Newton-Raphson algorithm fails in all sorts of situations, but works relatively well when $f(\mathbf{x})$ is a **convex function**:

$$f(\rho \cdot \mathbf{x}_1 + (1 - \rho) \cdot \mathbf{x}_2) \leq \rho \cdot f(\mathbf{x}_1) + (1 - \rho) \cdot f(\mathbf{x}_2),$$

$\forall \mathbf{x}_1, \mathbf{x}_2$ and $\rho \in (0, 1)$.

$f(\mathbf{x})$ is **strictly convex** if “ \leq ” is replaced by “ $<$ ”. Examples of convex functions:



Convex Functions

► **Definition:** $f(\rho \cdot \mathbf{x}_1 + (1 - \rho) \cdot \mathbf{x}_2) \leq \rho \cdot f(\mathbf{x}_1) + (1 - \rho) \cdot f(\mathbf{x}_2),$

$\forall \mathbf{x}_1, \mathbf{x}_2$ and $\rho \in (0, 1)$. Strictly convex if “ \leq ” is replaced by “ $<$ ”.

► **Properties:**

1. If $\nabla^2 f(\mathbf{x})$ is positive definite then $f(\mathbf{x})$ is strictly convex.
2. Sum of convex functions is convex.
3. f, g convex and $\nabla g(\mathbf{x}) \leq 0 \implies h(\mathbf{x}) = g(f(\mathbf{x}))$ convex.
4. $f(\mathbf{x})$ (strictly) convex $\implies f(\mathbf{Ax} + \mathbf{b})$ (strictly) convex.
5. If $f(\mathbf{x})$ is convex and \mathbf{x}_0 is a local minimum of f , then \mathbf{x}_0 is a global minimum.
6. If $f(\mathbf{x})$ is strictly convex, then it has either a unique global minimum or no minimum at all.

Convex Functions

Application to Exponential Families

- ▶ **Data:** $\mathbf{Y} = (\mathbf{Y}_1, \dots, \mathbf{Y}_n)$, $\mathbf{Y}_i \stackrel{\text{iid}}{\sim} \exp\{\mathbf{T}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})\}h(\mathbf{y})$.
- ▶ **Loglikelihood:** $\ell(\boldsymbol{\eta} | \mathbf{Y}) = n[\bar{\mathbf{T}}'\boldsymbol{\eta} - \Psi(\boldsymbol{\eta})]$, $\bar{\mathbf{T}} = \frac{1}{n} \sum_{i=1}^n \mathbf{T}(\mathbf{Y}_i)$.
- ▶ **Expected Fisher-Information:** If $\boldsymbol{\eta}$ is the true parameter value, then

$$\mathcal{I}(\boldsymbol{\eta}) = -\nabla^2 \ell(\boldsymbol{\eta} | \mathbf{Y}) = n \nabla^2 \Psi(\boldsymbol{\eta}) = \text{var}(\mathbf{T} | \boldsymbol{\eta})^{-1}.$$

Therefore:

- ▶ $-\ell(\boldsymbol{\eta} | \mathbf{Y})$ is a strictly convex function.
- ▶ If the MLE $\hat{\boldsymbol{\eta}}$ exists, then it is unique.
- ▶ Newton-Raphson is well-suited to find $\hat{\boldsymbol{\eta}}$. The NR updates are given by

$$\boldsymbol{\eta}_{n+1} = \boldsymbol{\eta}_n + [\nabla^2 \Psi(\boldsymbol{\eta}_n)]^{-1}[\bar{\mathbf{T}} - \nabla \Psi(\boldsymbol{\eta}_n)].$$

Application

Generalized Linear Models

► **Model:**

$$y_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \exp\{T_i \eta_i - \Psi(\eta_i)\} h(y_i), \quad \eta_i = \mathbf{x}_i' \boldsymbol{\beta}.$$

► **Loglikelihood:**

$$\ell(\boldsymbol{\beta} | \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n T_i \mathbf{x}_i' \boldsymbol{\beta} - \Psi(\mathbf{x}_i' \boldsymbol{\beta})$$

► **Hessian:**

$$\frac{\partial^2}{\partial \boldsymbol{\beta}^2} \ell(\boldsymbol{\beta} | \mathbf{y}, \mathbf{X}) = -\mathbf{X}' [\Psi^{(2)}(\mathbf{X}\boldsymbol{\beta})] \mathbf{X}, \quad \text{where}$$

$$\Psi^{(2)}(\eta) = \frac{d^2}{d\eta^2} \Psi(\eta), \quad \Psi^{(2)}(\mathbf{X}\boldsymbol{\beta}) = \text{diag}(\Psi^{(2)}(\mathbf{x}_1' \boldsymbol{\beta}), \dots, \Psi^{(2)}(\mathbf{x}_n' \boldsymbol{\beta})).$$

$\implies -\ell(\boldsymbol{\beta} | \mathbf{y}, \mathbf{X})$ is strictly convex since $\mathbf{X}' [\Psi^{(2)}(\mathbf{X}\boldsymbol{\beta})] \mathbf{X} = \text{var}(\mathbf{X}' \mathbf{z})$, where $\mathbf{z}_i \stackrel{\text{ind}}{\sim} \mathcal{N}(0, \Psi^{(2)}(\mathbf{x}_i' \boldsymbol{\beta}))$.

GLM: Common Cases

1. Poisson Regression (for count data)

► **Model:** $y_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \text{Pois}(\lambda_i), \quad \lambda_i = \exp(\mathbf{x}_i' \boldsymbol{\beta}).$

$$\implies E[y | \mathbf{x}] = \exp(\mathbf{x}' \boldsymbol{\beta}).$$

► **Log-Likelihood:**

$$\ell(\boldsymbol{\beta} | \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n y_i \cdot \mathbf{x}_i' \boldsymbol{\beta} - \exp(\mathbf{x}_i' \boldsymbol{\beta})$$

► **R command:**

```
M <- glm(y ~ x1 + x2, family = "poisson")
```

GLM: Common Cases

2. Binomial Regression (for success/failure data)

- **Model:** $y_i | \mathbf{x}_i, N_i \stackrel{\text{ind}}{\sim} \text{Bin}(N_i, \rho_i),$

$$\rho_i = \frac{1}{1 + \exp(-\mathbf{x}'_i \boldsymbol{\beta})} \quad \Longleftrightarrow \quad \mathbf{x}'_i \boldsymbol{\beta} = \log \left(\frac{\rho_i}{1 - \rho_i} \right) = \text{logit}(\rho_i).$$

- **Log-Likelihood:**

$$\begin{aligned} \ell(\boldsymbol{\beta} | \mathbf{y}, \mathbf{X}) &= \sum_{i=1}^n y_i \log \left(\frac{\rho_i}{1 - \rho_i} \right) + N_i \log(1 - \rho_i) \\ &= \sum_{i=1}^n y_i \mathbf{x}'_i \boldsymbol{\beta} - N_i \log \{1 + \exp(\mathbf{x}'_i \boldsymbol{\beta})\} \end{aligned}$$

- **Logistic Regression:** Special name for the common case where $N_i \equiv 1$.

Logistic Regression

Example

- ▶ **Model:** $y_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \text{Bernoulli}(\rho_i), \quad \rho_i = [1 + \exp(-\mathbf{x}_i' \boldsymbol{\beta})]^{-1}.$
- ▶ **Titanic Data:** 4-way contingency table of the $n = 2201$ passengers on the Titanic in the following categories:
 - ▶ $\text{Class} \in \{\text{1st}, \text{2nd}, \text{3rd}, \text{Crew}\}.$
 - ▶ $\text{Sex} \in \{\text{Male}, \text{Female}\}.$
 - ▶ $\text{Age} \in \{\text{Child}, \text{Adult}\}.$
 - ▶ $\text{Survived} \in \{\text{No}, \text{Yes}\}.$

Application of GLM/NR

Heteroscedastic Linear Regression

- **Usual Linear Regression:** $y_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \mathcal{N}(\mathbf{x}_i' \boldsymbol{\beta}, \sigma^2).$

Model has *homoscedastic errors*: $\text{var}(y | \mathbf{x}) \equiv \sigma^2$ is constant.

- **Heteroscedastic Linear Regression:**

$$y_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \mathcal{N}(\mathbf{x}_i' \boldsymbol{\beta}, \sigma_i^2), \quad \sigma_i = \sigma(\mathbf{x}_i),$$

such that $\text{var}(y | \mathbf{x}) = \sigma^2(\mathbf{x})$ is not constant (depends on \mathbf{x}).

Application of GLM/NR

Heteroscedastic Linear Regression

- **Model:** (ignore mean term for now)

$$y_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \mathcal{N}(0, \sigma_i^2), \quad \sigma_i^2 = \exp(\mathbf{x}_i' \boldsymbol{\beta}).$$

- **Log-Likelihood:**

$$\ell(\boldsymbol{\beta} | \mathbf{y}, \mathbf{X}) = -\frac{1}{2} \sum_{i=1}^n \frac{y_i^2}{\exp(\mathbf{x}_i' \boldsymbol{\beta})} + \mathbf{x}_i' \boldsymbol{\beta}.$$

- **Convexity:**

Let $g(\eta) = a \cdot \exp(\eta) + \eta$, for $\eta \in \mathbb{R}$, $a > 0$.

$$\implies \frac{d^2}{d\eta^2} g(\eta) = a \cdot \exp(\eta) > 0 \implies g(\eta) \text{ is convex.}$$

$$\implies -\ell(\boldsymbol{\beta} | \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n g(\mathbf{x}_i' \boldsymbol{\beta}) \text{ is also convex.}$$

Heteroscedastic Linear Regression

- ▶ **Simplified Model:** $y_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \mathcal{N}(0, \exp(\mathbf{x}_i' \boldsymbol{\beta})) \implies$

$$y_i^2 | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \underbrace{\text{Gamma}\left(\frac{1}{2}, 2\mu_i\right)}_{\mu_i \cdot \chi_{(1)}^2}, \quad \mu_i = \exp(\mathbf{x}_i' \boldsymbol{\beta}).$$

- ▶ *Gamma parametrization:*

$$z \sim \text{Gamma}(\alpha, \lambda) \implies \begin{aligned} E[Y] &= \alpha\lambda \\ \text{var}(Y) &= \alpha\lambda^2. \end{aligned}$$

- ▶ **Gamma Regression:**

$$z_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \text{Gamma}(1/\tau, \tau\mu_i), \quad \mu_i = g^{-1}(\mathbf{x}_i' \boldsymbol{\beta})$$

$$\implies E[z | \mathbf{x}] = g^{-1}(\mathbf{x}' \boldsymbol{\beta}), \quad \text{var}(z | \mathbf{x}) = \tau \cdot E[z | \mathbf{x}]^2$$

- ▶ $g(\mu)$: Link function.
- ▶ τ : Dispersion parameter.

Gamma Regression

► **Model:** $z_i | \mathbf{x}_i \stackrel{\text{ind}}{\sim} \text{Gamma}(1/\tau, \tau\mu_i), \quad \mu_i = g^{-1}(\mathbf{x}_i'\boldsymbol{\beta}).$

► **Log-Likelihood:**

$$\ell(\boldsymbol{\beta}, \tau | \mathbf{z}, \mathbf{X}) = \sum_{i=1}^n \left[\frac{\log g^{-1}(\mathbf{x}_i'\boldsymbol{\beta}) - z_i/g^{-1}(\mathbf{x}_i'\boldsymbol{\beta})}{\tau} \right] - n \log \Gamma(1/\tau) + \sum_{i=1}^n \frac{\log(z_i)}{\tau}$$

► **Properties:**

► $\ell(\boldsymbol{\beta}, \tau | \mathbf{z}, \mathbf{X})$ **convex** if $\mu(\mathbf{x}) = \exp(\mathbf{x}'\boldsymbol{\beta})$.

► $\hat{\boldsymbol{\beta}} = \arg \max_{\boldsymbol{\beta}} \ell(\boldsymbol{\beta}, \tau | \mathbf{z}, \mathbf{X})$ doesn't depend on τ .

► Two **independent** convex problems:

(i) find $\hat{\boldsymbol{\beta}}$, then (ii) find $\hat{\tau} = \arg \max_{\tau} \ell(\hat{\boldsymbol{\beta}}, \tau | \mathbf{z}, \mathbf{X})$.

► **R Command:** `glm(z ~ X, family = Gamma("log"))`

Heteroscedastic Linear Regression

► Full Model:

$$y_i \mid \mathbf{x}_i, \mathbf{w}_i \stackrel{\text{ind}}{\sim} \mathcal{N}(\mathbf{x}_i' \boldsymbol{\beta}, \exp(\mathbf{w}_i' \boldsymbol{\gamma})).$$

Can think of \mathbf{x} and \mathbf{w} as subsets of a single set of covariates $\boldsymbol{\mathcal{X}}$, e.g.,

$$\mathbf{x} = (\text{Age}, \text{Height}, \text{Weight}), \quad \mathbf{w} = (\log(\text{Age}), \text{Height}/\text{Weight}).$$

► Maximum Likelihood Estimation:

► **Initial Value:** $\boldsymbol{\beta}_0 = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}, \quad \boldsymbol{\gamma}_0 = \mathbf{0}.$

► **Iterative fitting:** Given $(\boldsymbol{\beta}_n, \boldsymbol{\gamma}_n),$

$$\text{► } \boldsymbol{\beta}_{n+1} = (\mathbf{X}' \boldsymbol{\Lambda}_n \mathbf{X})^{-1} \mathbf{X}' \boldsymbol{\Lambda}_n \mathbf{y}, \quad \boldsymbol{\Lambda}_n = \begin{bmatrix} \exp(-\mathbf{w}_1' \boldsymbol{\gamma}_n) & & \\ & \ddots & \\ & & \exp(-\mathbf{w}_n' \boldsymbol{\gamma}_n) \end{bmatrix}.$$

This is just MLE of $\boldsymbol{\beta}$ for $y_i \stackrel{\text{ind}}{\sim} \mathcal{N}(\mathbf{x}_i' \boldsymbol{\beta}, \exp(\mathbf{w}_i' \boldsymbol{\gamma}_n)).$

► $\boldsymbol{\gamma}_{n+1} = \text{coef}(\text{glm}(\mathbf{u}_{n+1}^2 \sim \mathbf{W}, \text{family} = \text{Gamma}(\text{"log"}))), \quad \mathbf{u}_{n+1} = \mathbf{y} - \mathbf{X} \boldsymbol{\beta}_{n+1}.$

This is just MLE of $\boldsymbol{\gamma}$ for $u_{i,n+1}^2 \stackrel{\text{ind}}{\sim} \text{Gamma}(1, \exp(\mathbf{w}_i' \boldsymbol{\gamma})).$

Heteroscedastic Linear Regression

Example

- ▶ **Model:** $y_i | \mathbf{x}_i, \mathbf{w}_i \stackrel{\text{ind}}{\sim} \mathcal{N}(\mathbf{x}_i' \boldsymbol{\beta}, \exp(\mathbf{w}_i' \boldsymbol{\gamma}))$.
- ▶ **SENIC Dataset:** Study on the Efficiency of Nosocomial Infection Control (SENIC). $n = 113$ US hospitals with following measurements:
 - ▶ **length:** Average length of stay of patients in days.
 - ▶ **age:** Average age of patients.
 - ▶ **inf:** Probability of acquiring infection in hospital.
 - ▶ **cult:** Culturing ratio, i.e. $100 \times \frac{\text{cultures performed}}{\# \text{ of patients with no infection}}$.
 - ▶ **xray:** Chest X-ray ratio (defined as above).
 - ▶ **beds:** Number of beds.
 - ▶ **school:** Medical school affiliation (1 = no, 2 = yes).
 - ▶ **region:** US geographic region (1 = NC, 2 = NE, 3 = S, 4 = W).
 - ▶ **pat:** Number of patients.
 - ▶ **nurs:** Number of nurses.
 - ▶ **serv:** Available facilities (at given hospital).

More Resources

- ▶ Useful **R** functions for `lm`, `glm` and other regression models (e.g., in package `survival`): `coef`, `effects`, `residuals`, `fitted`, `vcov`, `summary`, `predict`, `formula`.
- ▶ [Article](#) by Carl Morris (1982) on Exponential Families with so-called “quadratic variance functions” (easy to read and considered a great breakthrough in statistical theory).
- ▶ [Simplified version](#) by Morris & Lock (2009) with a nice figure relating the different EF distributions.