

Chapter 5, Section 1

James Lee

August 9, 2026

1. Let C, D be any two divisors on a surface X , and let the corresponding invertible sheaves be $\mathcal{L}$ and $\mathcal{M}$. Show that

$$C.D = \chi(\mathcal{O}_X) - \chi(\mathcal{L}^{-1}) - \chi(\mathcal{M}^{-1}) + \chi(\mathcal{L}^{-1} \otimes \mathcal{M}^{-1}).$$

Proof. Applying the Riemann-Roch theorem to the divisor $-C - D$ gives

$$\chi(\mathcal{O}_X(-C - D)) = \frac{1}{2}(-C - D).(-C - D - K) + \chi(\mathcal{O}_X)$$

The first term on the right hand side simplifies to

$$(-C - D).(-C - D - K) = C^2 + 2C.D + D^2 + C.K + D.K.$$

Substituting back to above and isolating $C.D$ gives

$$\begin{aligned} C.D &= -\chi(\mathcal{O}_X) + \chi(\mathcal{O}_X(-C - D)) - \frac{1}{2}C.(C + K) - \frac{1}{2}D.(C + K) \\ &= \chi(\mathcal{O}_X) + \chi(\mathcal{O}_X(-C - D)) - \left(\chi(\mathcal{O}_X) - \frac{1}{2}C.(-C - K) \right) - \left(\chi(\mathcal{O}_X) - \frac{1}{2}D.(-D - K) \right) \\ &= \chi(\mathcal{O}_X) + \chi(\mathcal{O}_X(-C - D)) - \chi(\mathcal{O}_X(-C)) - \chi(\mathcal{O}_X(-D)). \end{aligned}$$

□

2. Let H be a very ample divisor on the surface X , corresponding to a projective embedding $X \subseteq \mathbb{P}^N$. If we write the Hilbert polynomial of X (III, Ex. 5.2) as

$$P(z) = \frac{1}{2}az^2 + bz + c,$$

show that $a = H^2$, $b = \frac{1}{2}H^2 + 1 - \pi$, where π is the genus of a nonsingular curve representing H , and $c = 1 + p_a$. Thus the *degree* of X in $\mathbb{P}^N$, as defined in (I, §7), is just H^2 . Show also that if C is any curve in X , then the degree of C in $\mathbb{P}^N$ is just $C.H$.

Proof. By the Riemann-Roch theorem, we have the equality

$$P(z) = \frac{1}{2}H^2z^2 - \frac{1}{2}(H.K)z + 1 + p_a.$$

Matching coefficients readily gives a, b , and c . We can rewrite the coefficient of z as

$$-\frac{1}{2}(H.K) = \frac{1}{2}H^2 - \frac{1}{2}H.(H + K),$$

and the adjunction formula (1.5) gives

$$2\pi - 2 = H.(H + K).$$

We conclude that

$$b = \frac{1}{2}H^2 + 1 - \pi.$$

If C is any curve in X , then the degree of C in $\mathbb{P}^N$ is just the degree of $\mathcal{O}_X(H) \otimes \mathcal{O}_C$ as a line bundle on C , which is equal to $C.H$ by (1.3). □

3. Recall that the *arithmetic genus* of a projective scheme D of dimension 1 is defined as $p_a(D) = 1 - \chi(\mathcal{O}_D)$ (III, Ex. 5.3).

- (a) If D is an effective divisor on a surface X , use (1.6) to show that $2p_a(D) - 2 = D \cdot (D + K)$.
- (b) $p_a(D)$ depends only on the linear equivalence class of D on X .
- (c) More generally, for *any* divisor D on X , we define the *virtual arithmetic genus* (which is equal to the ordinary arithmetic genus if D is effective) by the same formula: $2p_a - 2 = D \cdot (D + K)$. Show that for any two divisors C, D , we have

$$p_a(-D) = D^2 - p_a(D) + 2,$$

and

$$p_a(C + D) = p_a(C) + p_a(D) + C \cdot D - 1.$$

Proof.

- (a) We have the exact sequence

$$0 \longrightarrow \mathcal{O}_X(-D) \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_D \longrightarrow 0,$$

and taking Euler characteristics gives

$$\chi(\mathcal{O}_D) = \chi(\mathcal{O}_X) - \chi(\mathcal{O}_X(-D)).$$

On the other hand, the Riemann-Roch theorem gives

$$\chi(\mathcal{O}_X(-D)) = \frac{1}{2} D \cdot (D + K) + \chi(\mathcal{O}_X).$$

Substituting gives

$$1 - \chi(\mathcal{O}_D) = 1 + \frac{1}{2} D \cdot (D + K).$$

- (b) Follows from (a) and (1.1(4)), which states $D \cdot (D + K)$ depends only on the linear equivalence class of D on X .

- (c)

$$\begin{aligned} 2p_a(-D) - 2 &= -D \cdot (-D + K) \\ &= D \cdot (D - K) \\ &= 2D^2 - D \cdot (D + K) \\ &= 2D^2 - 2p_a(D) + 2 \\ 2p_a(C + D) - 2 &= (C + D) \cdot (C + D + K) \\ &= C \cdot (C + K) + D \cdot (D + K) + 2(C \cdot D) \\ &= 2p_a(C) - 2 + 2p_a(D) - 2 + 2(C \cdot D) \end{aligned}$$

□

4. (a) If a surface X of degree d in $\mathbb{P}^3$ contains a straight line $C = \mathbb{P}^1$, show that $C^2 = 2 - d$.
 (b) Assume $\text{char } k = 0$, and show for every $d \geq 1$, there exists a nonsingular surface X of degree d in $\mathbb{P}^3$ containing the line $x = y = 0$.

Proof.

- (a) By (Ex. 1.5), $C^2 = -C \cdot K - 2$. By (1.3), $C \cdot K = d - 4$, because C is a straight line and $\mathcal{O}_X(K) = \mathcal{O}_X(d - 4)$. Hence, $C^2 = 2 - d$.
- (b) We can prove the following more general statement:

Proposition 0.1. *For every $n \geq 3$ and $d > 0$, and for every linear subspace $H \subset \mathbb{P}^n$ of codimension at least 2, there exists a nonsingular hypersurface of degree d in $\mathbb{P}^n$ containing H .*

We can assume H has codimension two and is defined by the vanishing of $x = y = 0$. Bertini's theorem (III, 10.9.2) says a general hypersurface X of degree d in $\mathbb{P}^3$ containing H is nonsingular away from H . Also, any such X is defined by the vanishing of $xF + yG = 0$, where F, G define surfaces of degree $d-1$. If P is a point in H , then X is singular at P if and only if both F, G vanish at P . This is a closed condition, so we can find plenty of choices of F, G which give a nonsingular hypersurface containing H .

Example 0.2. For $n = 3$ and $d \geq 2$, the surface defined by

$$x(z^{d-1} + x^{d-1}) + y(w^{d-1} + y^{d-1}) = 0$$

is nonsingular and contains L . □

5. (a) If X is a surface of degree d in $\mathbb{P}^3$, then $K^2 = d(d-4)^2$.
 (b) If X is a product of two nonsingular curves C, C' , of genus g, g' respectively, then $K^2 = 8(g-1)(g'-1)$.

Proof.

- (a) The adjunction formula gives $\mathcal{O}_X(K) = \mathcal{O}_X(d-4)$. If H is a general hyperplane section of X in $\mathbb{P}^3$, then $K = (d-4)H$, and $H^2 = d$ by (Ex. 1.2). Hence,

$$K^2 = (d-4)^2 H^2 = d(d-4)^2.$$

- (b) By (II, Ex. 8.3), there is an isomorphism $\omega_X \cong p_1^* \omega_C \otimes p_2^* \omega_{C'}$. In particular, if $\mathfrak{e}'$ and $\mathfrak{e}'$ are canonical divisors of C and C' , respectively, then $p_1^* \mathfrak{e} + p_2^* \mathfrak{e}'$ is a canonical divisor of X . Denote $D = p_1^* \mathfrak{e}$ and $D' = p_2^* \mathfrak{e}'$. The support of D consists of $2g-2$ copies of C' which are the fibres of the points in $\mathfrak{e}$ with respect to $p_1 : X \rightarrow C$. It follows that

$$D \cdot D' = (\deg K_C) \cdot (\deg K_{C'}) = 4(g-1)(g'-1).$$

Also, we can always find two canonical divisors with disjoint support, and since the intersection pairing only depends on the linear equivalence class, we have $D^2 = D'^2 = 0$. We conclude that $K^2 = 2D \cdot D' = 8(g-1)(g'-1)$. □

6. (a) If C is a curve of genus g , show that the diagonal $\Delta \subseteq C \times C$ has self-intersection $\Delta^2 = 2 - 2g$. (Use the definition of ω_C in (II, §8).)
 (b) Let $l = C \times \text{pt}$ and $m = \text{pt} \times C$. If $g \geq 1$, show that l, m , and Δ are linearly independent in $\text{Num}(C \times C)$. Thus $\text{Num}(C \times C)$ has rank ≥ 3 , and in particular, $\text{Pic}(C \times C) \neq p_1^* \text{Pic } C \oplus p_2^* \text{Pic } C$.

Proof.

- (a) By definition, $\omega_C \cong \mathcal{O}_X(-\Delta) \otimes \mathcal{O}_\Delta$, and by (1.3), $\deg_\Delta(\mathcal{O}_X(-\Delta) \otimes \mathcal{O}_\Delta) = -\Delta^2$.
 (b) Suppose $D = al + bm + c\Delta = 0$ in $\text{Num}(C \times C)$ for some integers $a, b, c \in \mathbb{Z}$. It is not hard to see

$$l^2 = m^2 = 0, \quad l \cdot m = 1, \quad \Delta \cdot l = \Delta \cdot m = 1,$$

which implies

$$D \cdot m = b + c = 0, \quad D \cdot l = a + c = 0, \quad D \cdot \Delta = a + b + c(2 - 2g) = 0.$$

Combining the first two gives $a + b + 2c = 0$. The condition $g \geq 1$ forces $a, b, c = 0$, which is what we wanted to show. □

7. Algebraic Equivalence of Divisors. Let X be a surface. Recall that we have defined an algebraic family of effective divisors on X , parametrized by a nonsingular curve T , to be an effective Cartier divisor D on $X \times T$, flat over T (III, 9.8.5). In this case, for any two closed points $0, 1 \in T$, we say that the corresponding divisors D_0, D_1 on X are prealgebraically equivalent. Two arbitrary divisors are prealgebraically equivalent if they are differences of prealgebraically equivalent divisors. Two divisors D, D' are *algebraically equivalent* if there is a finite sequence $D = D_0, D_1, \dots, D_n = D'$ with D_i and D_{i+1} prealgebraically equivalent for each i .

- (a) Show that the divisors algebraically equivalent to 0 form a subgroup of $\text{Div } X$.
- (b) Show that linearly equivalent divisors are algebraically equivalent.
- (c) Show that algebraically equivalent divisors are numerically equivalent.

Proof.

- (a)
- (b)
- (c)

□

8. Cohomology Class of a Divisor. For any divisor D on the surface X , we define its cohomology class $c(D) \in H^1(X, \Omega_X)$ by using the isomorphism $\text{Pic } X \cong H^1(X, \mathcal{O}_X^*)$ of (III, Ex. 4.5) and the sheaf homomorphism $d \log : \mathcal{O}_X^* \rightarrow \Omega_X$ (III, Ex. 7.4c). Thus we obtain a group homomorphism $c : \text{Pic } X \rightarrow H^1(X, \Omega_X)$. On the other hand, $H^1(X, \Omega_X)$ is dual to itself by Serre duality (III, 7.13), so we have a nondegenerate bilinear map

$$\langle \ , \ \rangle : H^1(X, \Omega_X) \times H^1(X, \Omega_X) \rightarrow k.$$

- (a) Prove that this is compatible with the intersection pairing, in the following sense: for any two divisors D, E on X , we have

$$\langle c(D), c(E) \rangle = (D.E) \cdot 1$$

in k .

- (b) If $\text{char } k = 0$, use the fact that $H^1(X, \Omega_X)$ is a finite-dimensional vector space to show that $\text{Num } X$ is a finitely generated free abelian group.

Proof.

- (a) We need to first show numerically equivalent divisors give isomorphic cohomology classes. Suppose $D \equiv 0$ and $D \neq 0$ in $\text{Pic } X$. Let (U, f) be a locally defining equation of D . We want to show $df/f = 0$. Let P be any point in X ,

- (b)

□

9. (a) If H is an ample divisor on the surface X , and if D is any divisor, show that

$$(D^2)(H^2) \leq (D.H)^2.$$

- (b) Now let X be a product of two curves $X = C \times C'$. Let $l = C \times \text{pt}$, and $m = \text{pt} \times C'$. For any divisor D on X , let $a = D.l$, $b = D.m$. Then we say D has *type* (a, b) . If D has type (a, b) , with $a, b \in \mathbb{Z}$, show that

$$D^2 \leq 2ab,$$

and equality holds if and only if $D = bl + am$.

Proof.

- (a) By replacing H with mH for some $m \gg 0$ if necessary, we may assume that H is an irreducible nonsingular curve corresponding to a very ample line bundle on X . Consider the case when D is an irreducible curve.
- (b)

□

Example 0.3. Consider the quadric surface $Q = \mathbb{P}^1 \times \mathbb{P}^1$ and the divisor H of type $(1, 1)$. If D be a divisor of type (a, b) , then by (1.4.3), the left-hand side is equal to $2ab$, while the right-hand side is equal to $(a + b)^2$ (IV, 6.4.1) (Ex. 1).

- 10.** *Weil's Proof [2] of the Analogue of the Riemann Hypothesis for Curves.* Let C be a curve of genus g defined over the finite field $\mathbb{F}_q$, and let N be the number of points of C rational over $\mathbb{F}_q$. Then $N = 1 - a + q$, with $|a| \leq 2g\sqrt{q}$. To prove this, we consider C as a curve over the algebraic closure k of $\mathbb{F}_q$. Let $f: C \rightarrow C$ be the k -linear Frobenius morphism obtained by taking q th powers, which makes sense since C is defined over $\mathbb{F}_q$, so $X_q \cong X$ (IV, 2.4.1). Let $\Gamma \subseteq C \times C$ be the graph of f , and let $\Delta \subseteq C \times C$ be the diagonal. Show that $\Gamma^2 = q(2 - 2g)$, and $\Gamma \cdot \Delta = N$. Then apply (Ex. 1.9) to $D = r\Gamma + s\Delta$ for all r and s to obtain the result.

Proof. For any non-zero $r, s \in \mathbb{Z}$, consider the divisor $D = r\Gamma - s\Delta$. It has type $(r - s, r - s)$, so by (Ex. 1.9), we have the inequality

$$D^2 \leq 2(r - s)^2.$$

□

- 11.** In this problem, we assume that X is a surface for which $\text{Num } X$ is finitely generated.
- (a) If H is an ample divisor on X , and $d \in \mathbb{Z}$, show that the set of effective divisors D with $D \cdot H = d$, modulo numerical equivalence, is a finite set.
- (b) Now let C be a curve of genus $g \geq 2$, and use (a) to show that the group of automorphisms of C is finite, as follows. Given an automorphism σ of C , let $\Gamma \subseteq X = C \times C$ be its graph. First show that if $\Gamma \equiv \Delta$, then $\Gamma = \Delta$, using the fact that $\Delta^2 < 0$, since $g \geq 2$ (Ex. 1.6). Then use (a).
- 12.** If D is an ample divisor on the surface X , and $D' \equiv D$, then D' is also ample. Give an example to show, however, that if D is very ample, D' need not be very ample.

Proof.

□