

Chapter 2, Section 9

James Lee

September 7, 2026

1. Let X be a Noetherian scheme, Y a closed subscheme, and $\hat{X}$ the completion of X along Y . We call the ring $\Gamma(\hat{X}, \mathcal{O}_{\hat{X}})$ the ring of *formal-regular functions* on X along Y . In this exercise we show that if Y is a connected, nonsingular, positive dimensional subvariety of $X = \mathbb{P}_k^n$ over an algebraically closed field k , then $\Gamma(\hat{X}, \mathcal{O}_{\hat{X}}) = k$.

- (a) Let $\mathcal{J}$ be the ideal sheaf of Y . Use (8.13) and (8.17) to show that there is an inclusion of sheaves on Y , $\mathcal{J}/\mathcal{J}^2 \hookrightarrow \mathcal{O}_Y(-1)^{n+1}$.
- (b) Show that for any $r \geq 1$, $\Gamma(Y, \mathcal{J}^r/\mathcal{J}^{r+1}) = 0$.
- (c) Use the exact sequences

$$0 \rightarrow \mathcal{J}^r/\mathcal{J}^{r+1} \rightarrow \mathcal{O}_X/\mathcal{J}^{r+1} \rightarrow \mathcal{O}_X/\mathcal{J}^r \rightarrow 0$$

and induction on r to show that $\Gamma(Y, \mathcal{O}_X/\mathcal{J}^r) = k$ for all $r \geq 1$.

- (d) Conclude that $\Gamma(\hat{X}, \mathcal{O}_{\hat{X}}) = k$.

2. Use the result of (Ex. 9.1) to prove the following geometric result. Let $Y \subseteq X = \mathbb{P}_k^n$ be as above, and let $f : X \rightarrow Z$ be a morphism of k -varieties. Suppose that $f(Y)$ is a single closed point $P \in Z$. Then $f(X) = P$ also.

3. Prove the analogue of (5.6) for formal schemes, which says, if $\mathfrak{X}$ is an affine formal scheme, and if

$$0 \rightarrow \mathfrak{F}' \rightarrow \mathfrak{F} \rightarrow \mathfrak{F}'' \rightarrow 0$$

is an exact sequence of $\mathcal{O}_{\mathfrak{X}}$ -modules, and if $\mathfrak{F}'$ is coherent, then the sequence of global sections

$$0 \rightarrow \Gamma(\mathfrak{X}, \mathfrak{F}') \rightarrow \Gamma(\mathfrak{X}, \mathfrak{F}) \rightarrow \Gamma(\mathfrak{X}, \mathfrak{F}'') \rightarrow 0$$

is exact. For the proof, proceed in the following steps.

- (a) Let $\mathfrak{J}$ be an ideal of definition for $\mathfrak{X}$, and for each $n > 0$ consider the exact sequence

$$0 \rightarrow \mathfrak{F}'/\mathfrak{J}^n \mathfrak{F}' \rightarrow \mathfrak{F}/\mathfrak{J}^n \mathfrak{F} \rightarrow \mathfrak{F}''/\mathfrak{J}^n \mathfrak{F}'' \rightarrow 0.$$

Use (5.6), slightly modified, to show that for every open affine subset $\mathfrak{U} \subseteq \mathfrak{X}$, the sequence

$$0 \rightarrow \Gamma(\mathfrak{U}, \mathfrak{F}'/\mathfrak{J}^n \mathfrak{F}') \rightarrow \Gamma(\mathfrak{U}, \mathfrak{F}/\mathfrak{J}^n \mathfrak{F}) \rightarrow \Gamma(\mathfrak{U}, \mathfrak{F}''/\mathfrak{J}^n \mathfrak{F}'') \rightarrow 0$$

is exact.

- (b) Now pass to the limit, using (9.1), (9.2), and (9.6). Conclude that $\mathfrak{F} \cong \varprojlim \mathfrak{F}/\mathfrak{J}^n \mathfrak{F}$ and that the sequence of global sections above is exact.

4. Use (Ex. 9.3) to prove that if

$$0 \rightarrow \mathfrak{F}' \rightarrow \mathfrak{F} \rightarrow \mathfrak{F}'' \rightarrow 0$$

is an exact sequence of $\mathcal{O}_{\mathfrak{X}}$ -modules on a Noetherian formal scheme $\mathfrak{X}$, and if $\mathfrak{F}', \mathfrak{F}''$ are coherent, then $\mathfrak{F}$ is coherent.

5. If $\mathcal{F}$ is a coherent sheaf on a Noetherian formal scheme $\mathfrak{X}$, which can be generated by global sections, show in fact that it can be generated by a finite number of its global sections.

6. Let $\mathfrak{X}$ be a Noetherian formal scheme, let $\mathfrak{J}$ be an ideal of definition, and for each n , let Y_n be the scheme $(\mathfrak{X}, \mathcal{O}_{\mathfrak{X}}/\mathfrak{J}^n)$. Assume that the inverse system of groups $(\Gamma(Y_n, \mathcal{O}_{Y_n}))$ satisfies the Mittag-Leffler condition. Then prove that $\text{Pic } \mathfrak{X} = \varprojlim \text{Pic } Y_n$. As in the case of a scheme, we define $\text{Pic } \mathfrak{X}$ to be the group of locally free $\mathcal{O}_{\mathfrak{X}}$ -modules of rank 1 under the operation $\otimes$. Proceed in the following steps.
- (a) Use the fact that $\ker(\Gamma(Y_{n+1}, \mathcal{O}_{Y_{n+1}}) \rightarrow \Gamma(Y_n, \mathcal{O}_{Y_n}))$ is a nilpotent ideal to show that the inverse system $(\Gamma(Y_n, \mathcal{O}_{Y_n}^*))$ of units in the respective rings also satisfies (ML).
 - (b) Let $\mathfrak{F}$ be a coherent sheaf of $\mathcal{O}_{\mathfrak{X}}$ -modules, and assume that for each n , there is some isomorphism $\varphi_n : \mathfrak{F}/\mathfrak{J}^n \mathfrak{F} \cong \mathcal{O}_{Y_n}$. Be careful, because the φ_n may not be compatible with the maps in the two inverse systems $(\mathfrak{F}/\mathfrak{J}^n \mathfrak{F})$ and $(\mathcal{O}_{Y_n})$. Then show that there is an isomorphism $\mathfrak{F} \cong \mathcal{O}_{\mathfrak{X}}$. Conclude that the natural map $\text{Pic } \mathfrak{X} \rightarrow \varprojlim \text{Pic } Y_n$ is injective.
 - (c) Given an invertible sheaf $\mathcal{L}_n$ on Y_n for each n , and given isomorphism $\mathcal{L}_{n+1} \otimes \mathcal{O}_{Y_n} \cong \mathcal{L}_n$, construct maps $\mathcal{L}_{n'} \rightarrow \mathcal{L}_n$ for each $n' \geq n$ so as to make an inverse system, and show that $\mathcal{L} = \varprojlim \mathcal{L}_n$ is a coherent sheaf on $\mathfrak{X}$. Then show that $\mathcal{L}$ is locally free of rank 1, and thus conclude that the map $\text{Pic } \mathfrak{X} \rightarrow \varprojlim \text{Pic } Y_n$ is surjective. Again be careful, because even though each $\mathcal{L}_n$ is locally free of rank 1, the open sets needed to make them free might get smaller and smaller with n .
 - (d) Show that the hypothesis “ $(\Gamma(Y_n, \mathcal{O}_{Y_n}))$ satisfies (ML)” is satisfied if either $\mathfrak{X}$ is affine, or each Y_n is projective over a field k .