

Chapter 2, Section 7

James Lee

October 6, 2026

1. Let $(X, \mathcal{O}_X)$ be a locally ringed space, and let $f : \mathcal{L} \rightarrow \mathcal{M}$ be a surjective map of invertible sheaves on X . Show that f is an isomorphism.

Proof. We reduce to the following algebraic problem: any A -linear surjective map $\phi : A \rightarrow A$ is an isomorphism. Indeed, ϕ is determined by $r = \phi(1)$, and ϕ surjective implies there exists $a \in A$ such that $ar = \phi(a) = 1$. Thus, r is a unit, so ϕ is injective. $\square$

2. Let X be a scheme over a field k . Let $\mathcal{L}$ be an invertible sheaf on X , and let $\{s_0, \dots, s_n\}$ and $\{t_0, \dots, t_m\}$ be two sets of sections of $\mathcal{L}$, which generate the same subspace $V \subseteq \Gamma(X, \mathcal{L})$, and which generate the sheaf $\mathcal{L}$ at every point. Suppose $n \leq m$. Show that the corresponding morphisms $\varphi : X \rightarrow \mathbb{P}_k^n$ and $\psi : X \rightarrow \mathbb{P}_k^m$ differ by a suitable linear projection $\mathbb{P}^m - L \rightarrow \mathbb{P}^n$ and an automorphism of $\mathbb{P}^n$, where L is a linear subspace of $\mathbb{P}^m$ of dimension $m - n - 1$.
3. Let $\varphi : \mathbb{P}_k^n \rightarrow \mathbb{P}_k^m$ be a morphism. Then:
 - (a) either $\varphi(\mathbb{P}^n) = pt$ or $m \geq n$ and $\dim \varphi(\mathbb{P}^n) = n$;
 - (b) in the second case, φ can be obtained as the composition of (1) a d -uple embedding $\mathbb{P}^n \rightarrow \mathbb{P}^N$ for a uniquely determined $d \geq 1$, (2) a linear projection $\mathbb{P}^N - L \rightarrow \mathbb{P}^m$, and an automorphism of $\mathbb{P}^m$. Also, φ has finite fibers.

Proof.

- (a) Let $\mathcal{L}$ be the invertible sheaf on $\mathbb{P}^n$ associated to the morphism $\varphi : \mathbb{P}^n \rightarrow \mathbb{P}^m$, which is generated by global sections $s_i = \varphi^*(x_i)$, $i = 0, 1, \dots, m$. Pick $P \in \varphi^{-1}((1, 0, \dots, 0))$ (we can assume such P always exists by applying an automorphism of $\mathbb{P}^m$). Identifying $\mathcal{L}_P \cong \mathcal{O}_{P, \mathbb{P}^n}$, we see that $(s_i)_P$ for $i \neq 0$ is not a unit, and they generate the local ring of $\mathcal{O}_{P, \mathbb{P}^n}$. Since $\mathbb{P}^n$ is smooth, the minimal number of generators of the maximal ideal of $\mathcal{O}_{P, \mathbb{P}^n}$ is equal to the dimension n . The sections s_i generate $\mathcal{O}_{P, \mathbb{P}^n}$. Hence, $m \geq n$. Our proof also shows the induced homomorphism of local rings $\varphi_P : \mathcal{O}_{\varphi(P), \mathbb{P}^m} \rightarrow \mathcal{O}_{P, \mathbb{P}^n}$ is surjective, which implies $\mathcal{O}_{\varphi(P), \varphi(\mathbb{P}^n)} = \mathcal{O}_{\varphi(P), \mathbb{P}^m} / \ker \varphi_P \cong \mathcal{O}_{P, \mathbb{P}^n}$. Hence, $\varphi(P)$ has dimension n . $\square$

5. Establish the following properties of ample and very ample invertible sheaves on a noetherian scheme X . $\mathcal{L}, \mathcal{M}$ will denote invertible sheaves, and for (d), (e) we assume furthermore that X is of finite type over a noetherian ring A .
 - (a) If $\mathcal{L}$ is ample and $\mathcal{M}$ is generated by global sections, then $\mathcal{L} \otimes \mathcal{M}$ is ample.
 - (b) If $\mathcal{L}$ is ample and $\mathcal{M}$ is arbitrary, then $\mathcal{M} \otimes \mathcal{L}^n$ is ample for sufficiently large n .
 - (c) If $\mathcal{L}, \mathcal{M}$ are both ample, so is $\mathcal{L} \otimes \mathcal{M}$.
 - (d) If $\mathcal{L}$ is very ample and $\mathcal{M}$ is generated by global sections, then $\mathcal{L} \otimes \mathcal{M}$ is very ample.
 - (e) If $\mathcal{L}$ is ample, then there is an $n_0 > 0$ such that $\mathcal{L}^n$ is very ample for all $n \geq n_0$.

Proof. Fix a coherent sheaf $\mathcal{F}$ on X .

- (a) Let $n \geq 0$ such that $\mathcal{F} \otimes \mathcal{L}^n$ is generated by global sections. If $\mathcal{M}$ is generated by global sections, then so is $\mathcal{M}^n$ for any $n > 0$. More generally, the tensor product of two invertible sheaves generated by global sections is generated by global sections. Indeed, if t_i and r_j are generating global sections of $\mathcal{L}$ and $\mathcal{M}$, then $\mathcal{L} \otimes \mathcal{M}$ is generated by $t_i \otimes r_j$ for all pairs (i, j) (tensor products commutes with colimits). Thus, $\mathcal{F} \otimes \mathcal{L}^n \otimes \mathcal{M}^n \cong \mathcal{F} \otimes (\mathcal{L} \otimes \mathcal{M})^n$ is generated by global sections. Hence, $\mathcal{L} \otimes \mathcal{M}$ is ample.

- (b) Let $n > 0$ such that $\mathcal{M} \otimes \mathcal{L}^{n-1}$ is generated by global sections. By (a), $\mathcal{L} \otimes (\mathcal{M} \otimes \mathcal{L}^{n-1}) = \mathcal{M} \otimes \mathcal{L}^n$ is ample.
- (c) If $\mathcal{M}$ is ample, then $\mathcal{M}^n$ is generated by global sections for large enough n . Also, $\mathcal{L}^n$ is ample by (II, 7.5), so $\mathcal{L}^n \otimes \mathcal{M}^n$ is ample. Hence, $\mathcal{L} \otimes \mathcal{M}$ is ample by (II, 7.5) again.
- (d) Let $i : X \rightarrow \mathbb{P}_A^n$ be an immersion such that $\mathcal{L} \cong i^* \mathcal{O}_{\mathbb{P}_A^n}(1)$, and let $\varphi : X \rightarrow \mathbb{P}_A^m$ be the unique A -morphism corresponding to $\mathcal{M}$ (II, 7.1). By the universal property of the fiber product $\mathbb{P}_A^n \times_A \mathbb{P}_A^m$, there exists a unique A -morphism $\phi : X \rightarrow \mathbb{P}_A^n \times_A \mathbb{P}_A^m$ such that $i = p_n \circ \phi$ and $\varphi = p_m \circ \phi$, where $p_n : \mathbb{P}_A^n \times_A \mathbb{P}_A^m \rightarrow \mathbb{P}_A^n$ and $p_m : \mathbb{P}_A^n \times_A \mathbb{P}_A^m \rightarrow \mathbb{P}_A^m$ are the natural projection maps. It is not hard to see ϕ is an immersion, and composing it with a Segre embedding (Ex. 5.12), we obtain an immersion $\phi' : X \rightarrow \mathbb{P}_A^N$, where $N = nm + n + m$, such that $\mathcal{L} \otimes \mathcal{M} \cong \phi'^* \mathcal{O}(1)$.
- (e) If $\mathcal{L}$ is ample, then there exists $n_0 > 0$ such that $\mathcal{L}^{n-1}$ is generated by global sections for all $n \geq n_0$. By (d), $\mathcal{L}^n = \mathcal{L} \otimes \mathcal{L}^{n-1}$ is very ample for all $n \geq n_0$.

□