# Chapter 1, Section 5

James Lee

October 25, 2026

**1.** Locate the singular points and sketch the following curves in $\mathbb{A}^2$ (assume char $k \neq 2$):

(a) $x^2 = x^4 + y^4$;
(b) $xy = x^6 + y^6$;
(c) $x^3 = y^2 + x^4 + y^4$;
(d) $x^2y + xy^2 = x^4 + y^4$.

| $f$ | $\partial f/\partial x$ | $\partial f/\partial y$ | $\langle \partial f/\partial x, \partial f/\partial y \rangle = \langle 0,0 \rangle$ | Sing $f$ |
|---|---|---|---|---|
| $x^4 - x^2 + y^4$ | $4x^3 - 2x$ | $4y^3$ | $(0,0),\ (\pm 1/\sqrt{2}, 0)$ | $(0,0)$ |
| $x^6 + y^6 - xy$ | $6x^5 - y$ | $6y^5 - x$ | $(0,0),\ (6^{-1/4}, 6^{-1/4})$ | $(0,0)$ |
| $x^4 + y^4 - x^3 + y^2$ | $4x^3 - 3x^2$ | $4y^3 - 2y$ | $(0,0),(3/4,0),$ $(0,\pm 1/\sqrt{2}),(3/4,\pm 1/\sqrt{2})$ | $(0,0)$ |
| $x^4 + y^4 - x^2y - xy^2$ | $4x^3 - 2xy - y^2$ | $4y^3 - 2xy - x^2$ | $(0,0),\ (3/4, 3/4)$ | $(0,0)$ |

(a) is the tacnode, (b) is the node, (c) is the cusp, and (d) is the triple point.

**2.** Locate the singular points and describe the singularities of the following surfaces in $\mathbb{A}^3$ (assume char $k \neq 2$).

(a) $xy^2 = z^2$;
(b) $x^2 + y^2 = z^2$;
(c) $xy + x^3 + y^3 = 0$.

| $f$ | $\partial f/\partial x$ | $\partial f/\partial y$ | $\partial f/\partial z$ | Sing $f$ |
|---|---|---|---|---|
| $xy^2 - z^2$ | $y^2$ | $2xy$ | $-2z$ | $(t,0,0),\ t \in k$ |
| $x^2 + y^2 - z^2$ | $2x$ | $2y$ | $-2z$ | $(0,0,0)$ |
| $x^3 + y^3 + xy$ | $3x^2 + y$ | $3y^2 + x$ | $0$ | $(0,0,t),\ t \in k$ |

(a) is the pinch point, (b) is the conical double point, and (c) is the double line.

**3.** *Multiplicities.* Let $Y \subseteq \mathbb{A}^2$ be a curve defined by the equation $f(x,y) = 0$. Let $P = (a,b)$ be a point of $\mathbb{A}^2$. Make a linear change of coordinates so that $P$ becomes the point $(0,0)$. Then write $f$ as a sum $f = f_0 + f_1 + \cdots + f_d$, where $f_i$ is a homogenous polynomial of degree $i$ in $x$ and $y$. Then we define the *multiplicity* of $P$ on $Y$, denoted $\mu_P(Y)$, to be the least $r$ such that $f_r \neq 0$. (Note that $P \in Y \Leftrightarrow \mu_P(Y) > 0$.) The linear factors of $f_r$ are called the *tangent directions at* $P$.

(a) Show that $\mu_P(Y) = 1 \Leftrightarrow P$ is a nonsingular point of $Y$.
(b) Find the multiplicity of each of the singular points in (Ex. 5.1) above.

*Proof.*

(a) Let $f'$ be the polynomial $f(x,y)$ after change of coordinates so that $P$ becomes the origin and let $O = (0,0)$ be the origin. Then, $P$ is a nonsingular point of $Y \Leftrightarrow (\partial f/\partial x)(P),\ (\partial f/\partial y)(P) \neq 0 \Leftrightarrow (\partial f'/\partial x)(O),\ (\partial f'/\partial y)(O) \neq 0 \Leftrightarrow \partial f'/\partial x,\ \partial f'/\partial y =$ higher degree terms + constant $\Leftrightarrow f'$ has linear terms $\Leftrightarrow \mu_P(Y) = 1$.

(b) (a), (b), and (c) has multiplicity 2, and (d) has multiplicity 3 at $O$.

□

**6.** *Blowing Up Curve Singularities.*

(a) Let $Y$ be the cusp or node of (Ex. 5.1). Show that the curve $\widetilde{Y}$ obtained by blowing up $Y$ at $O = (0,0)$ is nonsingular.

(b) We define a *node* (also called *ordinary double point*) to be a double point (i.e., a point of multiplicity 2) of a plane curve with distinct tangent directions. If $P$ is a node on a plane curve $Y$, show that $\varphi^{-1}(P)$ consists of two distinct nonsingular points on the blown-up curve $\widetilde{Y}$. We say that "blowing up $P$ resolves the singularity at $P$".

(c) Let $P \in Y$ be the tacnode of (Ex. 5.1). If $\varphi : \widetilde{Y} \to Y$ is the blowing-up at $P$, show that $\varphi^{-1}(P)$ is a node. Using (b) we see that the tacnode can be resolved by two successive blowing-ups.

(d) Let $Y$ be the plane curve $y^3 = x^5$, which has a "high order cusp" at $O$. Show that $O$ is a triple point; that blowing up $O$ gives rise to a double point (what kind?) and that one further blowing up resolves the singularity.

*Proof.*

(a) Let $Y$ be the curve defined by the equation $xy = x^6 + y^6$. It is the node of (Ex. 5.1). We consider the equations $xy = x^6 + y^6$ and $xu = ty$ in $\mathbb{A}^2 \times \mathbb{P}^1$, where $t$, $u$ are homogenous coordinates for $\mathbb{P}^1$. Now $\mathbb{P}^1$ is covered by open sets $t \neq 0$ and $u \neq 0$, which we can consider separately. If $t \neq 0$, we can set $t = 1$, and use $u$ as an affine parameter. Then we have the equations

$$xy = x^6 + y^6$$
$$y = xu$$

in $\mathbb{A}^3$ with coordinates $x$, $y$, $u$. Substituting, we get $x^2u = x^6 + x^6u^6$, which factors. Thus, we obtain two irreducible components, one defined by $x = 0$, $y = 0$, $u$ arbitrary, which is the exceptional curve $E$, and the other defined by $u = x^4 + x^4u^6$. This is $\widetilde{Y}$, which meets $E$ at the point $u = 0$. This curve is nonsingular since if we set $f(x, y, u) = x^4(1 + u^6) - u$, then

$$\frac{\partial f}{\partial x} = 0, \quad \frac{\partial f}{\partial y} = 0, \quad \frac{\partial f}{\partial u} = -1,$$

so $\widetilde{Y}$ is nonsingular for all points $t \neq 0$. The curve $Y$ is symmetric with respect to $x = y$, so $\widetilde{Y}$ is also nonsingular for all points $u \neq 0$, hence $\widetilde{Y}$ is nonsingular.

(b) We can assume $P$ is the origin so that $Y$ is defined by $f = f_2 + f_3 + \cdots + f_r$. Thus, we can write $f_2 = (ax + by)(cx + dy)$, where $(a, b) \neq (c, d)$ as points in $\mathbb{P}^1$. Take $t$, $u$ to be homogenous coordinates for $\mathbb{P}^1$. If $t \neq 0$, substituting $y = xu$ we have the equation

$$F_u = (a + bu)(c + du) + f_3' + \cdots + f_r' = 0,$$

where $f_i'$ is obtained by substituting $y = xu$ into $f_i$ and dividing by $x^2$, which is possible since $f_i$ is homogenous of degree $i > 2$. This equation defines $\widetilde{Y}$, which meets the exceptional curve at $(b, -a)$ and $(d, -c)$. Similarly, if $u \neq 0$, then we obtain the equation

$$F_t = (at + b)(ct + d) + \hat{f_3} + \cdots + \hat{f_r} = 0,$$

thus $(b, -a)$ and $(d, -c)$ are the only points $\widetilde{Y}$ meets the exceptional curve, that is $\varphi^{-1}(P) = \{(a, b), (c, d)\}$. Also, $\widetilde{Y}$ is nonsingular at both points since

$$\frac{\partial F_u}{\partial u} = b\left(c - \frac{ad}{b}\right) \neq 0, \quad \frac{\partial F_t}{\partial u} = b\left(a - \frac{bc}{d}\right) \neq 0.$$

(c) Let $Y$ be the plane curve defined by $x^2 = x^4 + y^4$ and let $t$, $u$, be homogenous coordinates for $\mathbb{P}^1$. If $t \neq 0$, then we have the equations

$$x^2 = x^4 + y^4$$
$$y = xu$$

in $\mathbb{A}^3$ with coordinates $x$, $y$, $u$. Substituting, we get $x^2 = x^4 + x^4u^4$, thus $\widetilde{Y}$ is defined by $1 = x^2 + x^2u^4$, which intersects the exceptional curve at $u = 0$. Similarly, if $u \neq 0$, then $\widetilde{Y}$ is defined by $t^2 = t^2y^2 + y^2$ and intersects the exceptional curve at $t = 0$. These two points correspond to the slopes of the two branches of $Y$ at $P$, which are the two distinct tangent directions on the node $P$.

(d) If $t \neq 0$, then we have the equation $x^2 = u^3$, which is a cusp. If $u \neq 0$, then we have the equation $1 = t^5 y^2$, which does not meet the exceptional curve.

□

**8.** Let $Y \subseteq \mathbb{P}^n$ be a projective variety of dimension $r$. Let $f_1, \ldots, f_t \in S = k[x_0, \ldots, x_n]$ be homogenous polynomials which generate the ideal of $Y$. Let $P \in Y$ be a point, with homogenous coordinates $P = (a_0, \ldots, a_n)$. Show that $P$ is nonsingular on $Y$ if and only if the rank of the matrix $\left\|(\partial f_i/\partial x_j)(a_0, \ldots, a_n)\right\|$ is $n - r$.

*Proof.* We follow the hint. The matrix $\|(\partial f_i/\partial x_i)(a_0, \ldots, a_n)\|$ has rank $n - r$ if and only if there exists a $(n - r) \times (n - r)$ submatrix with nonzero determinant. The determinant of a matrix of homogenous polynomials where entries of the same row have same degree is also a homogenous polynomial, hence the rank of $\left\|(\partial f_i/\partial x_j)(a_0, \ldots, a_n)\right\|$ is independent of the choice of homogenous coordinates of $P$. Assuming $a_0 \neq 0$, we can pass to an open affine $U_0 \subseteq \mathbb{P}^n$ containing $P$ and use the affine Jacobian matrix, where $U_0$ is the open affine subset consisting of all points with nonzero 0th coordinate. In particular, f we set $Y_0 = Y \cap U_0$, then $g_i(y_1, \ldots, y_n) = f_i(1, y_1, \ldots, y_n)$ $(1 \leq i \leq t)$ generate the ideal of $Y_0$. Thus, $P$ is nonsingular if and only if the rank of the matrix $\left\|(\partial g_i/\partial y_j)(a_1/a_0, \ldots, a_n/a_0)\right\|$ is $n - r$. Then, we have

$$\frac{\partial g_i}{\partial y_j} = \frac{\partial f_i(1, y_1, \ldots, y_n)}{\partial y_j} = \frac{\partial f_i(1, x_1, \ldots, x_n)}{\partial x_j}.$$

Since the rank of the Jacobian matrix is independent of the homogenous coordinates chosen for $P$, we conclude that $||(\partial g_i/\partial y_j)(P)||$ has an invertible $(n - r) \times (n - r)$ minor if and only if $\left\|(\partial f_i/\partial x_j)(P)\right\|$ has an invertible $(n - r) \times (n - r)$ minor. □

**10.** For a point $P$ on a variety $X$, let $\mathfrak{m}$ be the maximal ideal of the local ring $\mathcal{O}_P$. We define the *Zariski tangent space* $T_P(X)$ of $X$ at $P$ to be the dual $k$-vector space of $\mathfrak{m}/\mathfrak{m}^2$.

(a) For any point $P \in X$, $\dim T_P(X) \geq \dim X$, with equality if and only if $P$ is nonsingular.

(b) For any morphism $\varphi : X \to Y$, there is a natural induced $k$-linear map $T_P(\varphi) : T_P(X) \to T_{\varphi(P)}(Y)$.

(c) If $\varphi$ is the vertical projection of the parabola $x = y^2$ onto the $x$-axis, show that the induced map $T_0(\varphi)$ of tangent spaces at the origin is the zero map.

*Proof.*

(a) Since $\mathfrak{m}/\mathfrak{m}^2$ is a finite dimension $k$-vector space, $\dim_k T_p(X) = \dim_k \mathfrak{m}/\mathfrak{m}^2 \geq \dim X$ by (5.2A), and by (Ex. 3.12),

$$\dim_k \mathfrak{m}/\mathfrak{m}^2 = \dim \mathcal{O}_{P,X} = \dim X.$$

(b) A morphism $\varphi : X \to Y$ induces a map $\varphi_P^* : \mathcal{O}_{\varphi(P),Y} \to \mathcal{O}_{P,X}$ between local rings, where the maximal ideal $\mathfrak{n}$ of $\mathcal{O}_{\varphi(P),Y}$ is mapped into the maximal ideal $\mathfrak{m}$ of $\mathcal{O}_{P,X}$. Thus, we can define $T_P(\varphi)F$ as

$$(T_P(\varphi)F)f = F(d_\varphi f) = F(f \circ \varphi),$$

where $d_\varphi : \mathfrak{n}/\mathfrak{n}^2 \to \mathfrak{m}/\mathfrak{m}^2$ is the map induced by $\varphi_P^*$.

(c) By the formula above, it suffices to show $d_\varphi$ is the zero map at 0. Let $X$ be the parabola $x = y^2$ and let $Y$ by the $x$-axis. The maximal ideal $\mathfrak{m}$ of the local ring $\mathcal{O}_{0,X}$ is generated by $y$, thus $\mathfrak{m}/\mathfrak{m}^2$ is a one dimensional $k$-vector space spanned by $y$, and similarly $\mathfrak{n}/\mathfrak{n}^2$ of $\mathcal{O}_{0,Y}$ is a one dimensional $k$-vector space spanned by $x$. Thus, we have $\varphi^* x = y^2 \in \mathfrak{m}^2$, hence $d_\varphi = 0$, hence $T_0(\varphi) = 0$.

□