Chapter 3, Section 7

James Lee

October 25, 2026

1. Let X be an integral projective scheme of dimension ≥ 1 over a field k, and let $\mathcal{L}$ be an ample invertible sheaf on X. Then $\mathrm{H}^0(X, \mathcal{L}^{-1}) = 0$.

Proof. A global section of $\mathcal{L}^{-1}$ defines an $\mathcal{O}_X$-module morphism

$$\varphi : \mathcal{L} \to \mathcal{O}_X,$$

and hence induces morphisms

$$\varphi^{\otimes n} : \mathcal{L}^{\otimes n} \to \mathcal{O}_X$$

for all $n > 0$. Since $\mathcal{L}$ is locally free of rank one, φ is locally given by multiplication by a section of $\mathcal{O}_X$. If for some $x \in X$ the stalk map $\varphi_x^{\otimes n}$ is not injectve, it must be the zero map. As X is integral, this forces φ to vanish at the generic point, hence $\varphi = 0$. Assume therefore $\varphi \neq 0$. Since the global section functor is left exact, $\varphi^{\otimes n}$ induces an injective map

$$\Gamma(\varphi^{\otimes n}) : \mathrm{H}^0(X, \mathcal{L}^{\otimes n}) \to \mathrm{H}^0(X, \mathcal{O}_X).$$

However, $\mathrm{H}^0(X, \mathcal{O}_X)$ is finite dimensional over k (5.2), and for $n \gg 0$,

$$\dim_k \mathrm{H}^0(X, \mathcal{L}^{\otimes n}) = \chi(\mathcal{L}^{\otimes n}),$$

which is a polynomial in n (5.3), and hence it eventually exceeds $\dim_k \mathrm{H}^0(X, \mathcal{O}_X)$. This is a contradiction, and therefore $\varphi = 0$. □

2. Let $f : X \to Y$ be a finite morphism of projective schemes of the same dimension over a field k, and let ω_Y° be a dualizing sheaf for Y.

(a) Show that $f^!\omega_Y^\circ$ is a dualizing sheaf for X, where $f^!$ is defined as in (Ex. 6.10).

(b) If X and Y are both nonsingular, and k algebraically closed, conclude that there is a natural trace map $t : f_*\omega_X \to \omega_Y$.

Proof.

(a) We must first define a trace map $t_X : \mathrm{H}^n(X, f^!\omega_Y^\circ) \to k$. By (Ex. 4.1), we have a natural isomorphism

$$\mathrm{H}^n(X, f^!\omega_Y^\circ) \xrightarrow{\sim} \mathrm{H}^n(Y, f_* f^!\omega_Y^\circ).$$

There is a natural morphism of $\mathcal{O}_Y$-modules $f_* f^!\omega_Y \to \omega_Y$, which induces a map between cohomology

$$\mathrm{H}^n(Y, f_* f^!\omega_Y^\circ) \to \mathrm{H}^n(Y, \omega_Y^\circ).$$

Composing with the trace map of ω_Y° gives us a trace map for $f^!\omega_Y^\circ$. It remains to show t_X satisfies the universal property, so let $\mathcal{F}$ be a coherent sheaf on X. Then $f_*\mathcal{F}$ is also a coherent sheaf on Y, so by (Ex. 6.10), we have

$$\mathrm{Hom}_X(\mathcal{F}, f^!\omega_Y) \cong \mathrm{Hom}_Y(f_*\mathcal{F}, \omega_Y) \cong \mathrm{H}^n(Y, f_*\mathcal{F})^\vee \cong \mathrm{H}^n(X, \mathcal{F})^\vee.$$

(b) By (Ex. 6.10), defining a morphim $t : f_*\omega_X \to \omega_Y$ is the same as defining a morphism $\omega_X \to f^!\omega_Y$, so we can let t be the morphism corresponding to an isomorphism $\omega_X \to f^!\omega_Y$.

□

3. Let $X = \mathbb{P}^n_k$. Show that $\mathrm{H}^q(X, \Omega^p_X) = 0$ for $p \neq q$, k for $p = q$, $0 \leq p, q \leq n$.

Proof. Consider the Euler sequence of (II, 8.13)

$$0 \longrightarrow \Omega_X \longrightarrow \mathcal{O}_X(-1)^{\oplus n+1} \longrightarrow \mathcal{O}_X \longrightarrow 0.$$

For each $0 \leq p \leq n$, let $\mathcal{G}^{p+1} = \bigwedge^{p+1} \mathcal{O}_X(-1)^{\oplus n+1}$, which is isomorphic to the direct sum of $\binom{n+1}{p+1}$ copies of $\mathcal{O}_X(-1)$. By (II, Ex. 5.16c), for each $0 \leq p \leq n$, there exists a filtration $\{\mathcal{F}^q\}$ of $\mathcal{G}^{p+1}$ such that

$$\mathcal{F}^q/\mathcal{F}^{q+1} \cong \Omega^q_X \otimes \bigwedge^{p+1-q} \mathcal{O}_X = \begin{cases} \Omega^q_X & q = p, p+1 \\ 0 & \text{otherwise.} \end{cases}.$$

Since $\mathcal{O}_X$ has rank one,

$$\mathcal{G}^{p+1} = \mathcal{F}^0 = \mathcal{F}^1 = \cdots = \mathcal{F}^p, \quad \mathcal{F}^p/\mathcal{F}^{p+1} \cong \Omega^p_X, \quad \mathcal{F}^{p+1} \cong \Omega^{p+1}_X,$$

which can be put into the short exact sequence

$$0 \longrightarrow \Omega^{p+1}_X \longrightarrow \mathcal{G}^{p+1} \longrightarrow \Omega^p_X \longrightarrow 0.$$

Taking the long exact sequence of cohomology gives

$$\cdots \longrightarrow \mathrm{H}^i(X, \mathcal{G}^{p+1}) \longrightarrow \mathrm{H}^i(X, \Omega^p_X) \longrightarrow \mathrm{H}^{i+1}(X, \Omega^{p+1}_X) \longrightarrow \mathrm{H}^{i+1}(X, \mathcal{G}^{p+1}) \longrightarrow \cdots$$

Direct sums commute with cohomology (2.9), and $\mathrm{H}^i(X, \mathcal{O}_X(-1)) = 0$ for all i, which means $\mathrm{H}^i(X, \mathcal{G}^{p+1}) = 0$, so from the long exact sequence, we obtain the isomorphisms

$$\mathrm{H}^i(X, \Omega^p_X) \cong \mathrm{H}^{i+1}(X, \Omega^{p+1}_X).$$

Hence, if $q \geq p$,

$$\mathrm{H}^q(X, \Omega^p_X) \cong \mathrm{H}^{q-p}(X, \mathcal{O}_X) = \begin{cases} k & p = q, \\ 0 & \text{otherwise.} \end{cases}$$

The other Hodge numbers are obtained via Serre duality. □