

Notes on Linear Algebra Done Right¹

Hao Su

July 17, 2025

¹4th ed. Some exercises are omitted as trivial.

Contents

1	Vector Spaces	1
1A	\mathbb{R}^n and \mathbb{C}^n	1
1B	Definition of Vector Space	2
1C	Subspaces	5
2	Finite-Dimensional Vector Spaces	9
2A	Span and Linear Independence	9

Chapter 1

Vector Spaces

1A \mathbb{R}^n and \mathbb{C}^n

Defn 1.1 *Addition and multiplication on \mathbb{C} are defined by*

$$\begin{aligned}(a + bi) + (c + di) &= (a + c) + (b + d)i \\ (a + bi) \cdot (c + di) &= (ac - bd) + (ad + bc)i\end{aligned}$$

where $a, b, c, d \in \mathbb{R}$. ◇

By properties of \mathbb{R} and 1.1 we obtain properties of \mathbb{C} . By the existence of inverses we define $-\alpha$ and $\frac{1}{\alpha}$, and subtraction and division accordingly.

Note 1.2 Use \mathbb{F} (i.e., fields) to denote either \mathbb{R} or \mathbb{C} . Elements of \mathbb{F} are *scalars*. Say that x_k is the k^{th} *coordinate* of the *list* (x_1, \dots, x_n) (n , the *length* of the list, has to be finite). Lists, when thought of as arrows, are *vectors*. Addition and scalar multiplication on lists are defined componentwise in the standard way. ◇

Exercises

5 Additive inverse of complex arithmetic. ◁

This is due to the additive inverse of real arithmetic.

6 Multiplicative inverse of complex arithmetic. ◁

For $\alpha = a + bi$, where $a \neq 0$, we have $\frac{a - bi}{a^2 + b^2}$ as the multiplicative inverse of α . Thus one exists. Also all the multiplicative inverses of α equals it by real arithmetic.

8 Find two distinct square roots of i . ◁

We solve the equation $(a + bi)^2 = i$ to obtain $\pm \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right)$.

1B Definition of Vector Space

We need a space where things act like vectors, which is defined as follows.

Defn 1.3 A *vector space* over a field \mathbb{F} is a set V equipped with two operations: vector addition $+: V \times V \rightarrow V$, and scalar multiplication $\cdot: \mathbb{F} \times V \rightarrow V$, satisfying the following axioms:

1. $(V, +)$ is an abelian group, i.e.:

- (a) (Associativity) $u + (v + w) = (u + v) + w$ for all $u, v, w \in V$,
- (b) (Commutativity) $u + v = v + u$ for all $u, v \in V$,
- (c) (Identity) There exists an element $0 \in V$ such that $v + 0 = v$ for all $v \in V$,
- (d) (Inverses) For each $v \in V$, there exists an element $-v \in V$ such that $v + (-v) = 0$.

2. Scalar multiplication satisfies:

- (a) (Multiplicative identity) $1 \cdot v = v$ for all $v \in V$,
- (b) (Associativity) $a \cdot (b \cdot v) = (ab) \cdot v$ for all $a, b \in \mathbb{F}$, $v \in V$,
- (c) (Distributivity over vector addition) $a \cdot (u + v) = a \cdot u + a \cdot v$ for all $a \in \mathbb{F}$, $u, v \in V$,
- (d) (Distributivity over field addition) $(a + b) \cdot v = a \cdot v + b \cdot v$ for all $a, b \in \mathbb{F}$, $v \in V$. ◇

The simplest vector space is $\{0\}$.

Defn 1.4 Elements of a vector space are called *vectors* or *points*. ◇

Eg 1.5 For $f, g \in \mathbb{F}^S$ and $\lambda \in \mathbb{F}$, $f + g \in \mathbb{F}^S$ is defined by $\forall x \in S, (f + g)(x) = f(x) + g(x)$ and $\lambda f \in \mathbb{F}^S$ by $\forall x (\lambda f)(x) = \lambda f(x)$. If $S \neq \emptyset$, then \mathbb{F}^S is a vector space over \mathbb{F} . The vector space \mathbb{F}^n is a special case of \mathbb{F}^S , where $S = \{1, 2, \dots, n\}$. ◇

Thm 1.6 A vector space has a unique additive identity. \diamond

Proof. Suppose 0 and $0'$ are both additive identities for some vector space V . Then

$$0' = 0' + 0 = 0 + 0' = 0. \quad \dashv$$

Thm 1.7 Every element in a vector space has a unique additive inverse. \diamond

Proof. Suppose $a \in V$ has two additive inverses b and c . Then

$$b = b + (a + c) = (b + a) + c = c. \quad \dashv$$

Notations like $-v$ and $w - v$ make sense due to the uniqueness of additive inverses. From now on, V denotes a vector space over \mathbb{F} .

Thm 1.8 $\forall v \in V, 0v = 0$. \diamond

Proof. We have for any $v \in V$

$$0v = (0 + 0)v = 0v + 0v.$$

Adding the additive inverse of $0v$ to both sides of the equation above gives $0 = 0v$. \dashv

Comment. We have to use distributivity for that's where vector addition and scalar multiplication are connected in 1.3. The first equation holds because $0 \in \mathbb{F}$.

Thm 1.9 $\forall a \in \mathbb{F}, a0 = 0$. \diamond

Proof. We have for any $a \in \mathbb{F}$

$$a0 = a(0 + 0) = a0 + a0.$$

Adding the additive inverse of $a0$ to both sides of the equation above gives $0 = a0$. \dashv

Similarly, $0 = (1 + (-1))v$ gives us

Thm 1.10 $\forall v \in V, (-1)v = -v$. \diamond

Exercises

1 Prove that $\forall v \in V, -(-v) = v$. \triangleleft

$$0 = (-v) + (-(-v)) = v + (-v).$$

2 Suppose $a \in \mathbb{F}, v \in V$, and $av = 0$. Prove that $a = 0$ or $v = 0$. \triangleleft

If $a \neq 0$ we have that $v = (a \cdot \frac{1}{a})v = \frac{1}{a} \cdot (av) = 0$, which is equivalent to what we are asked to prove.

5 Show that (d) of item 1 in 1.3 can be replaced with 1.8. \triangleleft

We are to show the existence of additive inverse from 1.8 and the rest of 1.3.

$$0 = 0v = (1 - 1)v = v + (-1)v.$$

Thus the additive inverse of v exists, namely $(-1)v$.

6 Let ∞ and $-\infty$ denote two distinct objects, neither of which is in \mathbb{R} . Define an addition and scalar multiplication on $\mathbb{R} \cup \{\infty, -\infty\}$: the sum and product of two real numbers is as usual, and for $t \in \mathbb{R}$ define

$$t\infty = \begin{cases} -\infty & \text{if } t < 0, \\ 0 & \text{if } t = 0, \\ \infty & \text{if } t > 0, \end{cases} \quad t(-\infty) = \begin{cases} \infty & \text{if } t < 0, \\ 0 & \text{if } t = 0, \\ -\infty & \text{if } t > 0, \end{cases}$$

and

$$\begin{aligned} t + \infty &= \infty + t = \infty + \infty = \infty, \\ t + (-\infty) &= (-\infty) + t = (-\infty) + (-\infty) = -\infty, \\ \infty + (-\infty) &= (-\infty) + \infty = 0. \end{aligned}$$

With these operations of addition and scalar multiplication, is $\mathbb{R} \cup \{\infty, -\infty\}$ a vector space over \mathbb{R} ? Explain. \triangleleft

Consider $(\mathbb{R} \cup \{\infty, -\infty\}, +)$. Commutativity and existence of identity and inverses holds by definition. However $(u, v, w) = (3, \infty, -\infty)$ violates commutativity, hence $(\mathbb{R} \cup \{\infty, -\infty\}, +)$ is not an abelian group, thus $\mathbb{R} \cup \{\infty, -\infty\}$ not a vector space.

8 Suppose V is a real vector space.

- The *complexification* of V , denoted by $V_{\mathbb{C}}$, equals $V \times V$. An element of $V_{\mathbb{C}}$ is an ordered pair (u, v) , where $u, v \in V$, but we write this as $u + iv$.

- Addition on $V_{\mathbb{C}}$ is defined by

$$(u_1 + iv_1) + (u_2 + iv_2) = (u_1 + u_2) + i(v_1 + v_2)$$

for all $u_1, v_1, u_2, v_2 \in V$.

- Complex scalar multiplication on $V_{\mathbb{C}}$ is defined by

$$(a + bi)(u + iv) = (au - bv) + i(av + bu)$$

for all $a, b \in \mathbb{R}$ and all $u, v \in V$.

Prove that with the definitions of addition and scalar multiplication as above, $V_{\mathbb{C}}$ is a complex vector space. \triangleleft

We verify each of the requirements specified by 1.3 by properties of V as an vector space, very much like verifying the properties of \mathbb{C} by those of \mathbb{R} .

Comment. Think of V as a subset of $V_{\mathbb{C}}$ by identifying $u \in V$ with $u + i0$. The construction of $V_{\mathbb{C}}$ can then be thought of as generalizing the construction of \mathbb{C}^n from \mathbb{R}^n (thought of as a subset of \mathbb{C}^n .)

1C Subspaces

Defn 1.11 A subset U of V is called a *subspace* of V if U is also a vector space with the same additive identity, addition, and scalar multiplication as on V . \diamond

Thm 1.12 $U \subseteq V$ is a subspace of V iff it (1) has the additive identity of V (or is nonempty, because we can take $u \in U$ then $0u \in U$), (2) is closed under addition, and (3) is closed under scalar multiplication. \diamond

Proof. Both directions hold by definition. In particular, closure ensures that addition and multiplication are reasonably defined ($U \times U \rightarrow U$ and $\mathbb{F} \times U \rightarrow U$) and properties such as associativity hold because they hold on V and $U \subseteq V$. \dashv

Eg 1.13 The set of differentiable real-valued functions f on the interval $(0,3)$ such that $f'(2) = b$ is a subspace of $\mathbb{R}^{(0,3)}$ iff $b = 0$ for closure under scalar multiplication, which shows the linear structure underlying parts of calculus. The subspaces of \mathbb{R}^2 are precisely $\{0\}$ all lines in \mathbb{R}^2 containing the origin and \mathbb{R} , which intuitively justifies the word “linear”. \diamond

Defn 1.14 Suppose V_1, \dots, V_m are subspaces of V . The *sum* of them is

$$V_1 + \dots + V_m = \{v_1 + \dots + v_m \mid \bigwedge v_i \in V_i\} \quad \diamond$$

Thm 1.15 Suppose V_1, \dots, V_m are subspaces of V . Then $V_1 + \dots + V_m$ is the smallest subspace of V containing V_1, \dots, V_m . \diamond

Proof. That $V_1 + \dots + V_m$ is a subspace and contains V_1, \dots, V_m is trivial. Suppose that V' contains V_1, \dots, V_m and is a subspace. By 1.14 and closure under addition we have that $V_1 + \dots + V_m \subseteq V'$, thus the minimality. \dashv

Defn 1.16 Suppose V_1, \dots, V_m are subspaces of V . The sum $V_1 + \dots + V_m$ is called a *direct sum* if each element of $V_1 + \dots + V_m$ can be written in only one way as a sum $v_1 + \dots + v_m$ where each $v_k \in V_k$, denoted $V_1 \oplus \dots \oplus V_m$. \diamond

The definition of direct sum requires every vector in the sum to have a unique representation as an appropriate sum.

Thm 1.17 Suppose V_1, \dots, V_m are subspaces of V . Then $V_1 + \dots + V_m$ is a direct sum iff the only way to write 0 as a sum $v_1 + \dots + v_m$, where each $v_k \in V_k$, is by taking each v_k equal to 0. \diamond

Proof. (\Rightarrow is trivial. Consider \Leftarrow .) Suppose for sake of contradiction that $V_1 + \dots + V_m$ is *not* a direct sum. Then there exists $v \in V_1 + \dots + V_m$ such that $v = v'_1 + \dots + v'_m = v''_1 + \dots + v''_m$, where $v'_k \in V_k$ and $v''_k \in V_k$ for each k and $(v'_1, \dots, v'_m) \neq (v''_1, \dots, v''_m)$. Then we have $0 = (v'_1 - v''_1) + \dots + (v'_m - v''_m)$ and at least a j such that $v'_j - v''_j \neq 0$. Hence contradiction. \dashv

Thm 1.18 Suppose U and W are subspaces of V . Then

$$U + W \text{ is a direct sum} \Leftrightarrow U \cap W = \{0\}. \quad \diamond$$

Proof. \Rightarrow : Say that $v \in U \cap W$ and $v \neq 0$, then $0 = v + (-v)$, where $v \in U$ and $-v \in W$, hence $U + W$ is not a direct sum, and contradiction.

\Leftarrow : Say that $0 = a + b$, where $a \in U$, $b \in W$, and $a \neq 0$. Then $b = -a \in U$. Hence $b \in U \cap W$ and $b \neq 0$, and contradiction. \dashv

Sums of subspaces are analogous to unions of subsets. Similarly, direct sums of subspaces are analogous to disjoint unions of subsets.

Exercises

6 Is $\{(a, b, c) \in \mathbb{R}^3 \mid a^3 = b^3\}$ a subspace of \mathbb{R}^3 ? How about \mathbb{C}^3 ? \triangleleft

By $a^3 = b^3$ we have that $(a - b)(a^2 + ab + b^2) = 0$. That is, if $b \neq 0$ then $\frac{a}{b} \in \left\{1, \frac{-1 \pm \sqrt{3}i}{2}\right\}$. We need a unique $\frac{a}{b}$ so that it is closed under addition, which holds in \mathbb{R} but not in \mathbb{C} .

7 Prove or give a counterexample: If U is a nonempty subset of \mathbb{R}^2 such that U is closed under addition and contains additive inverses, then U is a subspace of \mathbb{R}^2 . \triangleleft

A counterexample is \mathbb{Z}^2 .

8 Give an example of a nonempty subset U of \mathbb{R}^2 such that U is closed under scalar multiplication, but U is not a subspace of \mathbb{R}^2 . \triangleleft

C.f. [Ex.1C.6](#). Easily we have that $\{(a, b) \mid a = 2b \vee a = b\}$ is not closed under addition. Note that $\{(a, b) \mid a = 2b\} + \{(a, b) \mid a = b\}$ is a direct sum, yet their union is not a subspace.

12 Prove that the union of two subspaces of V is a subspace of V iff one of the subspaces is a subset of another. \triangleleft

Suppose otherwise that U and W are subspaces of V , $u \in U \setminus W$, and $w \in W \setminus U$. Then $u + w \notin U \cup W$, for that otherwise contradiction will arise. Hence $U \cup W$ is not a subspace of V .

13 Prove that the union of three subspaces of V is a subspace of V iff one of the subspaces contains the other two. \triangleleft

to do

18 Does the operation of addition on the subspaces of V have an additive identity? Which subspaces have additive inverses? \triangleleft

$\{0\}$ is *the* additive identity of addition on subspaces of V . Also only $\{0\}$ has an additive inverse, namely itself. Suppose otherwise for sake of contradiction that $W \subseteq V$ has an additive inverse $-W$, $w \in W$, and $w \neq 0$. Then $w = w + 0 \in W + (-W) = \{0\}$, hence contradiction.

19 Prove or give a counterexample: If V_1, V_2, U are subspaces of V such that $V_1 + U = V_2 + U$ then $V_1 = V_2$. \triangleleft

$V_1 + U = V_2 + U$ should always hold, as long as $V_1 \subseteq U$ and $V_2 \subseteq U$, considering 1.15.

20 Suppose $U = \{(x, x, y, y) \in \mathbb{F}^4 \mid x, y \in \mathbb{F}\}$. Find a subspace W of \mathbb{F}^4 such that $\mathbb{F}^4 = U \oplus W$. \triangleleft

It is easy to verify that $\{(x, -x, y, -y) \in \mathbb{F}^4 \mid x, y \in \mathbb{F}\}$ suffices.

21 Suppose

$$U = \{(x, y, x + y, x - y, 2x) \in \mathbb{F}^5 : x, y \in \mathbb{F}\}.$$

Find a subspace W of \mathbb{F}^5 such that $\mathbb{F}^5 = U \oplus W$. \triangleleft

to do

22 Suppose

$$U = \{(x, y, x + y, x - y, 2x) \in \mathbb{F}^5 : x, y \in \mathbb{F}\}.$$

Find three subspaces W_1, W_2, W_3 of \mathbb{F}^5 , none of which equals $\{0\}$, such that

$$\mathbb{F}^5 = U \oplus W_1 \oplus W_2 \oplus W_3. \quad \triangleleft$$

to do

23 Prove or give a counterexample: If V_1, V_2, U are subspaces of V such that $V = V_1 \oplus U = V_2 \oplus U$ then $V_1 = V_2$. \triangleleft

A counterexample is that $V_1 = \{(x, 0) \mid x \in \mathbb{R}\}$, $V_2 = \{(x, x) \mid x \in \mathbb{R}\}$, $U = \{(0, y) \mid y \in \mathbb{R}\}$, and $V = \mathbb{R}^2$. The intuition is well illustrated by the counterexample: there are different sets of vectors that span a plane, which may share same components.

24 Let V_e denote the set of real-valued even functions on \mathbb{R} and V_o the set of real-valued odd functions. Show that $\mathbb{R}^{\mathbb{R}} = V_e \oplus V_o$. \triangleleft

Trivial considering that $f(x) = \frac{1}{2}(f(x) + f(-x)) + \frac{1}{2}(f(x) - f(-x))$.

Chapter 2

Finite-Dimensional Vector Spaces

2A Span and Linear Independence

We will usually write lists of vectors without surrounding parentheses.

Defn 2.1 A *linear combination* of a list v_1, \dots, v_m of vectors in V is a vector of the form

$$a_1v_1 + \dots + a_mv_m,$$

where $a_1, \dots, a_m \in \mathbb{F}$. ◇

Defn 2.2 The set of all linear combinations of a list of vectors v_1, \dots, v_m in V is called the *span* of v_1, \dots, v_m , denoted $\text{span}(v_1, \dots, v_m)$. The span of the empty list $()$ is defined to be $\{0\}$. ◇

Thm 2.3 The span of a list of vectors in V is the smallest subspace of V containing all vectors in the list. ◇

The proof is omitted as trivial.

Defn 2.4 If $\text{span}(v_1, \dots, v_m)$ equals V , say that the list v_1, \dots, v_m *spans* V . ◇

Defn 2.5 A vector space is called *finite-dimensional* if some list of vectors in it spans the space. A vector space is called *infinite-dimensional* if it is not finite-dimensional. ◇

Defn 2.6 A function $p : \mathbb{F} \rightarrow \mathbb{F}$ is called a *polynomial* with coefficients in \mathbb{F} if

there exist $a_0, \dots, a_m \in \mathbb{F}$ such that

$$p(z) = a_0 + a_1z + a_2z^2 + \dots + a_mz^m$$

for all $z \in \mathbb{F}$. $\mathcal{P}(\mathbb{F})$ is the set of all polynomials with coefficients in \mathbb{F} . \diamond

If a polynomial is represented by two set of coefficients, then subtracting one representation of the polynomial from the other produces a polynomial that is identically zero as a function on \mathbb{F} and hence has all zero coefficients (which is obvious). Thus the coefficients of a polynomial are uniquely determined by the polynomial.

Defn 2.7 A polynomial $p \in \mathcal{P}(\mathbb{F})$ is said to have *degree* m if there exist scalars $a_0, a_1, \dots, a_m \in \mathbb{F}$ with $a_m \neq 0$ such that for every $z \in \mathbb{F}$, we have

$$p(z) = a_0 + a_1z + \dots + a_mz^m.$$

The polynomial that is identically 0 is said to have degree $-\infty$. The degree of a polynomial p is denoted by $\deg p$. For m a nonnegative integer, $\mathcal{P}_m(\mathbb{F})$ denotes the set of all polynomials with coefficients in \mathbb{F} and degree at most m . \diamond

$\mathcal{P}_m(\mathbb{F}) = \text{span}(1, z, \dots, z^m)$ and thus a finite-dimensional vector space for each $m \in \mathbb{N}$. Consider any list of elements of $\mathcal{P}(\mathbb{F})$. Let m denote the highest degree of the polynomials in this list. Then every polynomial in the span of this list has degree at most m . Thus z^{m+1} is not in the span of our list. Hence no list spans $\mathcal{P}(\mathbb{F})$. Thus $\mathcal{P}(\mathbb{F})$ is infinite-dimensional.

Defn 2.8 A list v_1, \dots, v_m of vectors in V is called *linearly independent* if the only choice of $a_1, \dots, a_m \in \mathbb{F}$ that makes

$$a_1v_1 + \dots + a_mv_m = 0$$

is $a_1 = \dots = a_m = 0$. The empty list $()$ is declared to be linearly independent. \diamond

$\text{span}(v_1, \dots, v_m)$ is by definition $\text{span}(v_1) + \dots + \text{span}(v_m)$ and is a direct sum iff v_1, \dots, v_m is linearly independent. C.f. 1.17.

If some vectors are removed from a linearly independent list, the remaining list is also linearly independent.

Defn 2.9 A list of vectors in V is called *linearly dependent* if it is not linearly independent. \diamond

If some vector in a list of vectors in V is a linear combination of the other vectors, then the list is linearly dependent.

Thm 2.10 Suppose v_1, \dots, v_m is a linearly dependent list in V . Then there exists $k \in \{1, 2, \dots, m\}$ such that

$$v_k \in \text{span}(v_1, \dots, v_{k-1}).$$

Furthermore, if k satisfies the condition above and the k^{th} term is removed from v_1, \dots, v_m , then the span of the remaining list equals $\text{span}(v_1, \dots, v_m)$. \diamond

Proof.

—