





## Example of a Carnot engine





$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$






$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$

$$\text{adiabats: } V_A T_h^c = V_D T_c^c \text{ (exponent } c = \frac{C_V}{nR})$$

$$\text{and } V_C T_c^c = V_B T_h^c$$



$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$


$$\text{adiabats: } V_A T_h^c = V_D T_c^c \text{ (exponent } c = \frac{C_V}{nR})$$

$$\text{and } V_C T_c^c = V_B T_h^c$$

$$\text{obtain : } V_A V_C T_h^c T_c^c = V_D V_B T_h^c T_c^c$$

$$\text{and } \therefore \frac{V_A}{V_B} = \frac{V_D}{V_C} \implies q_c = -nRT_c \ln \frac{V_B}{V_A}$$

$$\therefore \frac{q_h}{q_c} = -\frac{T_h}{T_c}$$



$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$

adiabats:  $V_A T_h^c = V_D T_c^c$  (exponent  $c = \frac{C_V}{nR}$ )

and  $V_C T_c^c = V_B T_h^c$


obtain :  $V_A V_C T_h^c T_c^c = V_D V_B T_h^c T_c^c$

and  $\therefore \frac{V_A}{V_B} = \frac{V_D}{V_C} \implies q_c = -nRT_c \ln \frac{V_B}{V_A}$

$$\therefore \frac{q_h}{q_c} = -\frac{T_h}{T_c}$$

efficiency,  $\eta = \frac{\text{work performed}}{\text{heat absorbed}} = \frac{|w|}{q_h}$

$$= \frac{q_h + q_c}{q_h} = 1 + \frac{q_c}{q_h} = 1 - \frac{T_c}{T_h}$$



$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$

$$\text{adiabats: } V_A T_h^c = V_D T_c^c \quad (\text{exponent } c = \frac{C_V}{nR})$$

$$\text{and} \quad V_C T_c^c = V_B T_h^c$$

$$\text{obtain : } V_A V_C T_h^c T_c^c = V_D V_B T_h^c T_c^c$$

$$\text{and } \therefore \frac{V_A}{V_B} = \frac{V_D}{V_C} \implies q_c = -nRT_c \ln \frac{V_B}{V_A}$$

$$\therefore \frac{q_h}{q_c} = -\frac{T_h}{T_c}$$

$$\begin{aligned} \text{efficiency, } \eta &= \frac{\text{work performed}}{\text{heat absorbed}} = \frac{|w|}{q_h} \\ &= \frac{q_h + q_c}{q_h} = 1 + \frac{q_c}{q_h} = 1 - \frac{T_c}{T_h} \end{aligned}$$

$$\Delta S = \frac{q_h}{T_h} + 0 + \frac{q_c}{T_c} + 0 = 0$$

2nd Law : all reversible engines have same efficiency regardless of their construction




2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction

Let  $\eta_A > \eta_B$

engine A takes heat  $q_h$  from hot source and dumps  $q_c$  in cold reservoir




2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction

Let  $\eta_A > \eta_B$

engine A takes heat  $q_h$  from hot source and dumps  $q_c$  in cold reservoir

Reverse engine B



2nd Law : all reversible engines have same efficiency regardless of their construction


► Equivalent to Kelvin-Planck statement : Proof by contradiction

Let  $\eta_A > \eta_B$

engine A takes heat  $q_h$  from hot source and dumps  $q_c$  in cold reservoir

Reverse engine B

engine B acquires energy heat  $q_c$  from cold reservoir and releases  $q'_h < q_h$  to the hot end



2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction


Let  $\eta_A > \eta_B$

engine A takes heat  $q_h$  from hot source and dumps  $q_c$  in cold reservoir

Reverse engine B

engine B acquires energy heat  $q_c$  from cold reservoir and releases  $q'_h < q_h$  to the hot end

Form composite engine C : A+B



2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction

Let  $\eta_A > \eta_B$


engine A takes heat  $q_h$  from hot source and dumps  $q_c$  in cold reservoir

Reverse engine B

engine B acquires energy heat  $q_c$  from cold reservoir and releases  $q'_h < q_h$  to the hot end

Form composite engine C : A+B

C takes heat  $q_h - q'_h$  from hot source and converts it completely into work



2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction

Let  $\eta_A > \eta_B$

engine A takes heat  $q_h$  from hot source and dumps  $q_c$  in cold reservoir



Reverse engine B

engine B acquires energy heat  $q_c$  from cold reservoir and releases  $q'_h < q_h$  to the hot end

Form composite engine C : A+B


C takes heat  $q_h - q'_h$  from hot source and converts it completely into work

⇒ violation of Kelvin-Planck statement of second law




universality of 2nd law :

any reversible cycle  $\approx$   
collection of Carnot cycles  
 $\oint$ (arbitrary path)  
 $= \sum \oint$  (each cycle)



universality of 2nd law :

any reversible cycle  $\approx$   
collection of Carnot cycles  
 $\oint$ (arbitrary path)  
 $= \sum \oint$  (each cycle)




$\Delta S$  along an individual path inside is  
cancelled by  $\Delta S$  along the path it shares with  
neighbouring cycle



universality of 2nd law :

any reversible cycle  $\approx$   
collection of Carnot cycles  
 $\oint$ (arbitrary path)  
 $= \sum \oint$  (each cycle)




$\Delta S$  along an individual path inside is  
cancelled by  $\Delta S$  along the path it shares with  
neighbouring cycle

The approxmn. becomes exact as individual  
cycles become infinitesimal

$$\oint$$
(arbitrary path) $= \sum \oint$  (each cycle) $= 0$

universality of 2nd law :

any reversible cycle  $\approx$   
collection of Carnot cycles  
 $\oint$ (arbitrary path)  
 $= \sum \oint$  (each cycle)



$\Delta S$  along an individual path inside is  
cancelled by  $\Delta S$  along the path it shares with  
neighbouring cycle

The approxmn. becomes exact as individual  
cycles become infinitesimal

$$\oint$$
(arbitrary path) $= \sum \oint$  (each cycle) $= 0$


$\therefore$  all entropy changes cancel except for those  
along perimeter of the overall cycle

$$\sum_{\text{all}} \frac{q_{\text{rev}}}{T} = \sum_{\text{perimeter}} \frac{q_{\text{rev}}}{T} = 0$$

universality of 2nd law :

any reversible cycle  $\approx$   
collection of Carnot cycles  
 $\oint$ (arbitrary path)

$$= \sum \oint \text{(each cycle)}$$



$\Delta S$  along an individual path inside is cancelled by  $\Delta S$  along the path it shares with neighbouring cycle

The approxmn. becomes exact as individual cycles become infinitesimal

$$\oint \text{(arbitrary path)} = \sum \oint \text{(each cycle)} = 0$$

$\therefore$  all entropy changes cancel except for those along perimeter of the overall cycle

$$\sum_{\text{all}} \frac{q_{\text{rev}}}{T} = \sum_{\text{perimeter}} \frac{q_{\text{rev}}}{T} = 0$$

In the limit of infinitesimal cycles, the non-cancelling edges of Carnot cycles match the overall cycle exactly, and the sum becomes an integral

$\Rightarrow dS$  is exact differential and  $S$  is state function

Clausius inequality

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

## Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$  and  $dw$  are negative when energy leaves the system as work

## Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$  and  $dw$  are negative when energy leaves the system as work

$$\therefore -dw_{\text{rev}} \geq -dw \text{ or, } dw \geq dw_{\text{rev}}$$



## Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$  and  $dw$  are negative when energy leaves the system as work

$$\therefore -dw_{\text{rev}} \geq -dw \text{ or, } dw \geq dw_{\text{rev}}$$

$$dU = dq + dw = dq_{\text{rev}} + dw_{\text{rev}}$$

## Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$  and  $dw$  are negative when energy leaves the system as work

$$\therefore -dw_{\text{rev}} \geq -dw \text{ or, } dw \geq dw_{\text{rev}}$$


$$dU = dq + dw = dq_{\text{rev}} + dw_{\text{rev}}$$

$$dq_{\text{rev}} \geq dq, \therefore \Delta S = \int \frac{dq_{\text{rev}}}{T} \geq \int \frac{dq}{T}$$

Consider transfer of energy as heat from  
one system - hot source-at  $T_h$   
to another system - the cold sink- at  $T_c$

Consider transfer of energy as heat from  
one system - hot source-at  $T_h$   
to another system - the cold sink- at  $T_c$


thermodynamic refrigerator



energy removed from cool source at temp.  $T_c$   
 $= \frac{|q_c|}{T_c}$

Consider transfer of energy as heat from  
one system - hot source-at  $T_h$   
to another system - the cold sink- at  $T_c$

thermodynamic refrigerator




energy removed from cool source at temp.  $T_c$   
 $= \frac{|q_c|}{T_c}$

and deposited in warmer sink at temp.  $T_h$

Consider transfer of energy as heat from  
one system - hot source-at  $T_h$   
to another system - the cold sink- at  $T_c$

thermodynamic refrigerator




energy removed from cool source at temp.  $T_c$   
 $= \frac{|q_c|}{T_c}$

and deposited in warmer sink at temp.  $T_h$

$$\Delta S = -\frac{|q_c|}{T_c} + \frac{|q_c|}{T_h} < 0 \implies \text{not spontaneous}$$

Consider transfer of energy as heat from  
one system - hot source-at  $T_h$   
to another system - the cold sink- at  $T_c$

thermodynamic refrigerator



energy removed from cool source at temp.  $T_c$   
 $= \frac{|q_c|}{T_c}$


and deposited in warmer sink at temp.  $T_h$

$$\Delta S = -\frac{|q_c|}{T_c} + \frac{|q_c|}{T_h} < 0 \implies \text{not spontaneous}$$

to generate more entropy, energy must be  
added to the stream that enters the warm sink.


task: find minimum energy to be supplied

## Thermodynamic refrigerator






## Thermodynamic refrigerator



coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

## Thermodynamic refrigerator




coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is  $c$

## Thermodynamic refrigerator




coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is  $c$

and more efficient the refrigerator.

## Thermodynamic refrigerator



coefficient of performance,


$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is  $c$

and more efficient the refrigerator.

$$\frac{1}{c} = \frac{|q_h| - |q_c|}{|q_c|} = \frac{|q_h|}{|q_c|} - 1$$

## Thermodynamic refrigerator



coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is  $c$

and more efficient the refrigerator.

$$\frac{1}{c} = \frac{|q_h| - |q_c|}{|q_c|} = \frac{|q_h|}{|q_c|} - 1$$

using  $\frac{q_h}{q_c} = -\frac{T_h}{T_c}$ , we get,  $c = \frac{T_c}{T_h - T_c}$

work to maintain a low temperature is relevant to design of refrigerators

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate  $\propto \Delta T$

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate  $\propto \Delta T$

If rate at which energy leaks =  $A(T_h - T_c)$

where  $A$  depends on sample size and insulation

then minimum power,  $P$ , required to maintain original  $\Delta T$  by pumping out energy by heating surroundings

$$P = \frac{1}{c} \cdot A (T_h - T_c) = \frac{A(T_h - T_c)^2}{T_c} \propto (\Delta T)^2$$



work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate  $\propto \Delta T$

If rate at which energy leaks =  $A(T_h - T_c)$

where  $A$  depends on sample size and insulation

then minimum power,  $P$ , required to maintain original  $\Delta T$  by pumping out energy by heating surroundings

$$P = \frac{1}{c} \cdot A(T_h - T_c) = \frac{A(T_h - T_c)^2}{T_c} \propto (\Delta T)^2$$

power increases as  $(\Delta T)^2$

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate  $\propto \Delta T$

If rate at which energy leaks =  $A(T_h - T_c)$

where  $A$  depends on sample size and insulation

then minimum power,  $P$ , required to maintain original  $\Delta T$  by pumping out energy by heating surroundings


$$P = \frac{1}{c} \cdot A(T_h - T_c) = \frac{A(T_h - T_c)^2}{T_c} \propto (\Delta T)^2$$

power increases as  $(\Delta T)^2$


$\therefore$  air-conditioners more expensive to run on hot days than on mild days

## Equivalence of Kelvin-Planck and Clausius statements

## Equivalence of Kelvin-Planck and Clausius statements




## Equivalence of Kelvin-Planck and Clausius statements




—C—K


## Equivalence of Kelvin-Planck and Clausius statements




—CC —K



## Equivalence of Kelvin-Planck and Clausius statements




—C $\subset$ —K




—K $\subset$ —C

entropy change in isothermal expansion :






entropy change in isothermal expansion :



$$\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$$


entropy change in isothermal expansion :



$$\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

entropy change in isothermal expansion :




$$\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

$$\Delta U = 0$$

$\therefore$  If isothermal expansion occurs freely,  
 $w_{\text{surr}} = 0$  and irreversibly, then  $q = 0$

entropy change in isothermal expansion :



$$\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$$


$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

$$\Delta U = 0$$

$\therefore$  If isothermal expansion occurs freely,  
 $w_{\text{surr}} = 0$  and irreversibly, then  $q = 0$

$$\therefore \Delta S_{\text{surr (irrev)}} = 0$$

entropy change in isothermal expansion :



$$\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

$$\Delta U = 0$$

$\therefore$  If isothermal expansion occurs freely,  
 $w_{\text{surr}} = 0$  and irreversibly, then  $q = 0$

$$\therefore \Delta S_{\text{surr (irrev)}} = 0$$

$$\Delta S_{\text{total (rev)}} = 0 \text{ rev}$$
$$> 0 \text{ irrev}$$

Phase transition :  $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Phase transition :  $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same  $\Delta_{\text{vap}}S$  ( $\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$ )

Phase transition :  $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same  $\Delta_{\text{vap}}S$  ( $\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$ )

Ex. liq.  $\text{Br}_2$ :

- no complexity like hydrogen bonding in liquid  $\text{Br}_2$
- there is no unusual behaviour in the gas phase
- safe to use Trouton's rule



Phase transition :  $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same  $\Delta_{\text{vap}}S$  ( $\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$ )

Ex. liq.  $\text{Br}_2$ :

- no complexity like hydrogen bonding in liquid  $\text{Br}_2$   
there is no unusual behaviour in the gas phase
- safe to use Trouton's rule

$$\begin{aligned} \text{- boils at } 59.2^\circ\text{C}, \Delta_{\text{vap}}H^\ominus &= T_b \times 85 \text{ J mol}^{-1} \\ &= 332.4 \times 85 = 28 \text{ kJ mol}^{-1} \end{aligned}$$

Phase transition :  $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same  $\Delta_{\text{vap}}S$  ( $\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$ )

Ex. liq.  $\text{Br}_2$ :

- no complexity like hydrogen bonding in liquid  $\text{Br}_2$   
there is no unusual behaviour in the gas phase
- safe to use Trouton's rule

$$\begin{aligned} \text{- boils at } 59.2^\circ\text{C}, \Delta_{\text{vap}}H^\ominus &= T_b \times 85 \text{ J mol}^{-1} \\ &= 332.4 \times 85 = 28 \text{ kJ mol}^{-1} \end{aligned}$$

experimental value :  $29.45 \text{ kJ mol}^{-1}$


entropy change as a function of temperature (varying heat capacity) :

$$S(T_f) = S(T_i) + \int_{T_i}^{T_f} \frac{dq_{\text{rev}}}{T}$$

entropy change as a function of temperature (varying heat capacity) :

$$S(T_f) = S(T_i) + \int_{T_i}^{T_f} \frac{dq_{\text{rev}}}{T}$$

$$\begin{aligned} \text{const. } p(V) : S(T_f) &= S(T_i) + \int_{T_i}^{T_f} \frac{C_{p(V)} dT}{T} \\ &= S(T_i) + C_{p(V)} \ln \frac{T_f}{T_i} \end{aligned}$$



label : heat capacity

solved problem : Calculate  $\Delta S$  when  $0.500 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  expands to  $1.000 \text{ dm}^3$  and is simultaneously heated to  $100^\circ\text{C}$

solved problem : Calculate  $\Delta S$  when  $0.500 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  expands to  $1.000 \text{ dm}^3$  and is simultaneously heated to  $100^\circ\text{C}$


$S$  is a state function

$\therefore$  choose a convenient path from initial state

solved problem : Calculate  $\Delta S$  when  $0.500 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  expands to  $1.000 \text{ dm}^3$  and is simultaneously heated to  $100^\circ\text{C}$

$S$  is a state function

$\therefore$  choose a convenient path from initial state



solved problem : Calculate  $\Delta S$  when  $0.500 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  expands to  $1.000 \text{ dm}^3$  and is simultaneously heated to  $100^\circ\text{C}$

$S$  is a state function

$\therefore$  choose a convenient path from initial state

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 1 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{heat}} 1 \text{ dm}^3 \text{ Ar}; 100^\circ\text{C}$

$$\Delta S_1 = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i};$$



solved problem : Calculate  $\Delta S$  when  $0.500 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  expands to  $1.000 \text{ dm}^3$  and is simultaneously heated to  $100^\circ\text{C}$

$S$  is a state function

$\therefore$  choose a convenient path from initial state

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 1 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{heat}} 1 \text{ dm}^3 \text{ Ar}; 100^\circ\text{C}$

$$\Delta S_{\text{I}} = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{\text{II}} = C_V \ln \frac{T_f}{T_i}$$

solved problem : Calculate  $\Delta S$  when  $0.500 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  expands to  $1.000 \text{ dm}^3$  and is simultaneously heated to  $100^\circ\text{C}$

$S$  is a state function

$\therefore$  choose a convenient path from initial state


$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 1 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{heat}} 1 \text{ dm}^3 \text{ Ar}; 100^\circ\text{C}$

$$\Delta S_I = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{II} = C_V \ln \frac{T_f}{T_i}$$

$$\begin{aligned} \Delta S &= \Delta S_I + \Delta S_{II} = nR \left( \ln \frac{V_f}{V_i} + \frac{3}{2} \ln \frac{T_f}{T_i} \right) = \frac{p_i V_i}{T_i} \ln \left( \frac{V_f}{V_i} \left[ \frac{T_f}{T_i} \right]^{\frac{3}{2}} \right) \\ &= \frac{10^5 \text{ Pa} \times 0.5 \times 10^{-3} \text{ m}^3}{298} \ln \left( \frac{1}{0.5} \left[ \frac{373}{298} \right]^{\frac{3}{2}} \right) \\ &= \frac{10^5 \text{ Pa} \times 0.5 \times 10^{-3} \text{ m}^3}{298} \ln \left( \frac{1}{0.5} \left[ \frac{373}{298} \right]^{\frac{3}{2}} \right) = 0.173 \text{ JK}^{-1} \end{aligned}$$

solved problem : Calculate  $\Delta S$  when  $0.5 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  is compressed to  $0.05 \text{ dm}^3$  and is simultaneously cooled to  $-25^\circ\text{C}$

solved problem : Calculate  $\Delta S$  when  $0.5 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  is compressed to  $0.05 \text{ dm}^3$  and is simultaneously cooled to  $-25^\circ\text{C}$



solved problem : Calculate  $\Delta S$  when  $0.5 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  is compressed to  $0.05 \text{ dm}^3$  and is simultaneously cooled to  $-25^\circ\text{C}$

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 0.05 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{cool}} 0.05 \text{ dm}^3 \text{ Ar}; -25^\circ\text{C}$

$$\Delta S_1 = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i};$$

solved problem : Calculate  $\Delta S$  when  $0.5 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  is compressed to  $0.05 \text{ dm}^3$  and is simultaneously cooled to  $-25^\circ\text{C}$

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 0.05 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{cool}} 0.05 \text{ dm}^3 \text{ Ar}; -25^\circ\text{C}$

$$\Delta S_{\text{I}} = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{\text{II}} = C_V \ln \frac{T_f}{T_i}$$


solved problem : Calculate  $\Delta S$  when  $0.5 \text{ dm}^3$  of Ar at  $25^\circ\text{C}$  and  $1.00 \text{ bar}$  is compressed to  $0.05 \text{ dm}^3$  and is simultaneously cooled to  $-25^\circ\text{C}$

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 0.05 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{cool}} 0.05 \text{ dm}^3 \text{ Ar}; -25^\circ\text{C}$

$$\Delta S_I = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{II} = C_V \ln \frac{T_f}{T_i}$$

$$\begin{aligned} \Delta S &= \Delta S_I + \Delta S_{II} = nR \left( \ln \frac{V_f}{V_i} + \frac{3}{2} \ln \frac{T_f}{T_i} \right) = \frac{p_i V_i}{T_i} \ln \left( \frac{V_f}{V_i} \left[ \frac{T_f}{T_i} \right]^{\frac{3}{2}} \right) \\ &= \frac{10^5 \text{ Pa} \times 0.5 \times 10^{-3} \text{ m}^3}{298} \ln \left( \frac{0.05}{0.5} \left[ \frac{248}{298} \right]^{\frac{3}{2}} \right) = \\ &= -0.44 = -0.44 \text{ JK}^{-1} \end{aligned}$$


Heat capacity vs. temp.




entropy vs.  $T$




Heat capacity vs. temp.




entropy vs.  $T$



Heat capacity vs. temp.



entropy vs.  $T$



$$S(T) = S(0) + \int_0^{T_f} \frac{C_p(s)dT}{T} + \frac{\Delta_{\text{fus}} H}{T_{\text{fus}}} + \int_{T_1}^{T_b} \frac{C_p(l)dT}{T} + \frac{\Delta_{\text{vap}} H}{T_{\text{vap}}} + \int_{T_b}^T \frac{C_p(g)dT}{T}$$

All terms except  $S(0)$  can be measured calorimetrically

All terms except  $S(0)$  can be measured calorimetrically

integrals can be evaluated either graphically or by fitting a polynomial to the data and integrating analytically

All terms except  $S(0)$  can be measured calorimetrically

integrals can be evaluated either graphically or by fitting a polynomial to the data and integrating analytically

$\therefore \frac{dT}{T} = d \ln T$ , one can evaluate area under a plot of  $C_p$  against  $\ln T$

All terms except  $S(0)$  can be measured calorimetrically

integrals can be evaluated either graphically or by fitting a polynomial to the data and integrating analytically

$\therefore \frac{dT}{T} = d \ln T$ , one can evaluate area under a plot of  $C_p$  against  $\ln T$

Debye extrapolation :  $\lim_{T \rightarrow 0} C_p \longrightarrow aT^3$

Ex.  $S_m^\ominus$  for  $\text{N}_2(\text{g})$  at  $25^\circ\text{C}$  calculated from following data :

Ex.  $S_m^\ominus$  for  $\text{N}_2(\text{g})$  at  $25^\circ\text{C}$  calculated from following data :

$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
---



Ex.  $S_m^\ominus$  for  $\text{N}_2(\text{g})$  at  $25^\circ\text{C}$  calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ C$  calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ C$  calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ\text{C}$  calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ\text{C}$  calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ\text{C}$  calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ\text{C}$  calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ C$  calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20



Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ C$  calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20
Correction for gas imperfection	0.92

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ C$  calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20
Correction for gas imperfection	0.92
Total	192.06

Ex.  $S_m^\ominus$  for  $N_2(g)$  at  $25^\circ C$  calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20
Correction for gas imperfection	0.92
Total	192.06

$$S_m(298.15K) = S_m(0) + 192.06 J K^{-1} mol^{-1}$$

Nernst heat theorem :  $\lim_{T \rightarrow 0} \Delta S \longrightarrow 0$  for any transformation provided all substances involved are perfectly crystalline

Nernst heat theorem :  $\lim_{T \rightarrow 0} \Delta S \longrightarrow 0$  for any transformation provided all substances involved are perfectly crystalline

3rd law (?) :

The entropy of all perfect crystalline substances is zero at  $T = 0$

Boltzmann formula :  $S = k_B \ln \Omega$

$\Omega = \#$ ways the molecules can be arranged in a crystalline form

Boltzmann formula :  $S = k_B \ln \Omega$

$\Omega = \# \text{ways the molecules can be arranged in a crystalline form}$

In most cases,  $\lim_{T \rightarrow 0} \Omega = 1$ ,  $\therefore \lim_{T \rightarrow 0} S = 0$

Boltzmann formula :  $S = k_B \ln \Omega$

$\Omega = \# \text{ways the molecules can be arranged in a crystalline form}$

In most cases,  $\lim_{T \rightarrow 0} \Omega = 1$ ,  $\therefore \lim_{T \rightarrow 0} S = 0$

when  $\Omega \neq 1$ , then  $S = S(0)$ , residual entropy



Ice-I<sub>h</sub> :

each H atom can lie either close to or far  
from its 'parent' O atom

total # of arrangements in sample of  $N$

H<sub>2</sub>O molecules with  $2N$  H atoms =  $2^{2N}$

Ice-I<sub>h</sub> :

each H atom can lie either close to or far from its 'parent' O atom

total # of arrangements in sample of  $N$

H<sub>2</sub>O molecules with  $2N$  H atoms =  $2^{2N}$

consider a single central O atom.

total number of arrangements of

locations of H atoms around central O

atom of one H<sub>2</sub>O molecule is  $2^4 = 16$

Ice-I<sub>h</sub> :

each H atom can lie either close to or far from its 'parent' O atom

total # of arrangements in sample of  $N$

H<sub>2</sub>O molecules with  $2N$  H atoms =  $2^{2N}$

consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H<sub>2</sub>O molecule is  $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds

Ice-I<sub>h</sub> :

each H atom can lie either close to or far from its 'parent' O atom


total # of arrangements in sample of  $N$  H<sub>2</sub>O molecules with  $2N$  H atoms =  $2^{2N}$

consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H<sub>2</sub>O molecule is  $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds

only  $\frac{6}{16} = \frac{3}{8}$  of all arrangements are possible, and for  $N$  molecules only  $(\frac{3}{8})^N$  of all arrangements are possible



Ice-I<sub>h</sub> :


each H atom can lie either close to or far from its 'parent' O atom

total # of arrangements in sample of  $N$  H<sub>2</sub>O molecules with  $2N$  H atoms =  $2^{2N}$   
consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H<sub>2</sub>O molecule is  $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds

only  $\frac{6}{16} = \frac{3}{8}$  of all arrangements are possible, and for  $N$  molecules only  $(3/8)^N$  of all arrangements are possible



∴ for  $N$  water molecules  
# of possible configurations  
 $= 2^{2N} (3/8)^N = \left(\frac{3}{2}\right)^N$

Ice-I<sub>h</sub> :


each H atom can lie either close to or far from its 'parent' O atom

total # of arrangements in sample of  $N$  H<sub>2</sub>O molecules with  $2N$  H atoms  $= 2^{2N}$   
consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H<sub>2</sub>O molecule is  $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds

only  $\frac{6}{16} = \frac{3}{8}$  of all arrangements are possible, and for  $N$  molecules only  $(3/8)^N$  of all arrangements are possible



$\therefore$  for  $N$  water molecules  
# of possible configurations  
 $= 2^{2N} (3/8)^N = \left(\frac{3}{2}\right)^N$

$$S(0) = Nk_B \ln \left(\frac{3}{2}\right) = 3.37 \text{ J mol}^{-1}\text{K}^{-1}$$

molecular interpretation : Consider the increase in entropy with volume

molecular interpretation : Consider the increase in entropy with volume


As the system size increases  
energy levels move closer together and more  
become accessible to the molecules



molecular interpretation : Consider the increase in entropy with volume

As the system size increases  
energy levels move closer together and more  
become accessible to the molecules

# ways of achieving same energy ( $\Omega$ ) increases



**Maxwell's demon** : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers

**Maxwell's demon** : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers

Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

**Maxwell's demon** : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers

Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

expansion is irreversible as there is no way to get the gas back into the initial chamber without doing work.

**Maxwell's demon** : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum


Open the tap and gas in first chamber expands to fill both chambers

Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

expansion is irreversible as there is no way to get the gas back into the initial chamber without doing work.

Maxwell imagined that the tap was operated by a microscopic intelligent creature. If it sees a molecule heading from the second chamber back into the first, it opens the tap and then shuts it, just letting the molecule through. If it sees a molecule heading from the first chamber back into the second chamber, it keeps the tap closed



**Maxwell's demon** : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers


Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

expansion is irreversible as there is no way to get the gas back into the initial chamber without doing work.

Maxwell imagined that the tap was operated by a microscopic intelligent creature. If it sees a molecule heading from the second chamber back into the first, it opens the tap and then shuts it, just letting the molecule through. If it sees a molecule heading from the first chamber back into the second chamber, it keeps the tap closed

Demon does no work and yet it makes molecules in second chamber all go back into the first chamber



Thus it creates a pressure difference between the two chambers where none existed before the demon started its mischief

Thus it creates a pressure difference between the two chambers where none existed before the demon started its mischief

as if the demon could therefore cause entropy to decrease in a system with no consequent increase in entropy anywhere else



- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it
- ▶ However, erasing information is irreversible

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it
- ▶ However, erasing information is irreversible
- ▶ Erasing information has an associated increase in entropy

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it
- ▶ However, erasing information is irreversible
- ▶ Erasing information has an associated increase in entropy
- ▶ Maxwell's demon can operate reversibly therefore, but only if it has a large enough hard disc that it doesn't ever need to clear space to continue operating.

**The Landauer principle** : “information is physical” and it has an energy equivalent  
quantifies the thermodynamic cost of the recording/erasure of one bit of information,



**The Landauer principle** : “information is physical” and it has an energy equivalent

quantifies the thermodynamic cost of the recording/erasure of one bit of information,

- ▶ Erasure of one bit of information requires a minimum energy cost equal to  $k_B T \ln 2 \approx 0.018$  eV, where  $T$  is the temperature of a thermal reservoir used in the process.

- ▶ Information theory is usually supplied in a form that is independent of any physical embodiment.

- ▶ Information theory is usually supplied in a form that is independent of any physical embodiment.
- ▶ Landauer argued that information is physical and it has an energy equivalent. It may be stored in physical systems such as books and memory chips and it is transmitted by physical devices exploiting electrical or optical signals.

- ▶ Information theory is usually supplied in a form that is independent of any physical embodiment.
- ▶ Landauer argued that information is physical and it has an energy equivalent. It may be stored in physical systems such as books and memory chips and it is transmitted by physical devices exploiting electrical or optical signals.
- ▶  $\therefore$  it must obey the laws of physics and, first and foremost, the laws of thermodynamics.

(1) any logically irreversible process must result in an entropy increase in the non-information-bearing degrees of freedom of the information-processing system or its environment

- (1) any logically irreversible process must result in an entropy increase in the non-information-bearing degrees of freedom of the information-processing system or its environment
- (2) any logically reversible process can be implemented thermodynamically reversibly.

(1) any logically irreversible process must result in an entropy increase in the non-information-bearing degrees of freedom of the information-processing system or its environment

(2) any logically reversible process can be implemented thermodynamically reversibly.

- an amount of energy equal to  $k_B T \ln 2$  ( $k_B T$  = thermal noise per unit bandwidth) is needed to transmit a bit of information, and more if quantized channels are used with photon energies  $h\nu > kT$