

A general property of exact differentials :

consider $f(x, y)$ and $df = gdx + hdy$, where $g \equiv g(x, y)$ and $h \equiv h(x, y)$

A general property of exact differentials :

consider $f(x, y)$ and $df = gdx + hdy$, where $g \equiv g(x, y)$ and $h \equiv h(x, y)$ e.g., for

$$U(S, V), \quad dU = TdS - pdV$$

A general property of exact differentials :

consider $f(x, y)$ and $df = gdx + hdy$, where $g \equiv g(x, y)$ and $h \equiv h(x, y)$ e.g., for

$U(S, V)$, $dU = TdS - pdV$ when df is an exact differential,

$$\left(\frac{\partial g}{\partial y}\right)_x = \underbrace{\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 f}{\partial x \partial y}}_{\text{both x and y, vary}} = \left(\frac{\partial h}{\partial x}\right)_y$$

Maxwell relations :
for reversible changes in systems with no non-mechanical work

Maxwell relations :

for reversible changes in systems with no non-mechanical work

$$dU = TdS - pdV$$

$$\left(\frac{\partial T}{\partial V}\right)_S = -\left(\frac{\partial p}{\partial S}\right)_V$$

Maxwell relations :

for reversible changes in systems with no non-mechanical work

$$dU = TdS - pdV$$

$$\left(\frac{\partial T}{\partial V}\right)_S = -\left(\frac{\partial p}{\partial S}\right)_V$$

$$dH = TdS + Vdp$$

$$\left(\frac{\partial T}{\partial p}\right)_S = \left(\frac{\partial V}{\partial S}\right)_p$$

Maxwell relations :

for reversible changes in systems with no non-mechanical work

$$dU = TdS - pdV$$

$$\left(\frac{\partial T}{\partial V}\right)_S = -\left(\frac{\partial p}{\partial S}\right)_V$$

$$dH = TdS + Vdp$$

$$\left(\frac{\partial T}{\partial p}\right)_S = \left(\frac{\partial V}{\partial S}\right)_p$$

$$dA = -SdT - pdV$$

$$\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$$

Maxwell relations :

for reversible changes in systems with no non-mechanical work

$$dU = TdS - pdV$$

$$\left(\frac{\partial T}{\partial V}\right)_S = -\left(\frac{\partial p}{\partial S}\right)_V$$

$$dH = TdS + Vdp$$

$$\left(\frac{\partial T}{\partial p}\right)_S = \left(\frac{\partial V}{\partial S}\right)_p$$

$$dA = -SdT - pdV$$

$$\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$$

$$dG = Vdp - SdT$$

$$\left(\frac{\partial V}{\partial T}\right)_p = -\left(\frac{\partial S}{\partial p}\right)_T$$

Another definition of temperature

Another definition of temperature

$$dU = TdS - pdV \implies \boxed{\frac{1}{T} = \left(\frac{\partial S}{\partial U}\right)_V}$$

Another definition of temperature

$$dU = TdS - pdV \implies \boxed{\frac{1}{T} = \left(\frac{\partial S}{\partial U}\right)_V}$$

$$dH = TdS + Vdp \implies \frac{1}{T} = \left(\frac{\partial S}{\partial H}\right)_p$$

Thermodynamic equation of state :

$$dU = TdS - pdV$$

$$\left(\frac{\partial U}{\partial V}\right)_T = T \left(\frac{\partial S}{\partial V}\right)_T - p \quad \text{or, } \pi_T = T \left(\frac{\partial S}{\partial V}\right)_T - p$$

Thermodynamic equation of state :

$$dU = TdS - pdV$$

$$\left(\frac{\partial U}{\partial V}\right)_T = T \left(\frac{\partial S}{\partial V}\right)_T - p \quad \text{or, } \pi_T = T \left(\frac{\partial S}{\partial V}\right)_T - p$$

Using Maxwell relation $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$

$$\pi_T = T \left(\frac{\partial p}{\partial T}\right)_V - p$$

Thermodynamic equation of state :

$$dU = TdS - pdV$$

$$\left(\frac{\partial U}{\partial V}\right)_T = T \left(\frac{\partial S}{\partial V}\right)_T - p \quad \text{or, } \pi_T = T \left(\frac{\partial S}{\partial V}\right)_T - p$$

Using Maxwell relation $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$

$$\pi_T = T \left(\frac{\partial p}{\partial T}\right)_V - p$$

For perfect gas, show that this leads to $\pi_T = 0$

Thermodynamic equation of state :

$$dU = TdS - pdV$$

$$\left(\frac{\partial U}{\partial V}\right)_T = T \left(\frac{\partial S}{\partial V}\right)_T - p \quad \text{or, } \pi_T = T \left(\frac{\partial S}{\partial V}\right)_T - p$$

Using Maxwell relation $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$

$$\pi_T = T \left(\frac{\partial p}{\partial T}\right)_V - p$$

For perfect gas, show that this leads to $\pi_T = 0$

Find for van der Waals gas., $\pi_T = a \frac{n^2}{V^2}$

Thermodynamic equation of state :

$$dU = TdS - pdV$$

$$\left(\frac{\partial U}{\partial V}\right)_T = T \left(\frac{\partial S}{\partial V}\right)_T - p \quad \text{or, } \pi_T = T \left(\frac{\partial S}{\partial V}\right)_T - p$$

Using Maxwell relation $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T$

$$\pi_T = T \left(\frac{\partial p}{\partial T}\right)_V - p$$

For perfect gas, show that this leads to $\pi_T = 0$

Find for van der Waals gas., $\pi_T = a \frac{n^2}{V^2} > 0 \implies \left(\frac{\partial U}{\partial V}\right)_T > 0$

Gibbs' Free energy and entropy


Gibbs' Free energy and entropy

$$\left(\frac{\partial G}{\partial T}\right)_p = -S < 0$$


Gibbs' Free energy and entropy

$$\left(\frac{\partial G}{\partial T}\right)_p = -S < 0$$


$$\left(\frac{\partial G}{\partial p}\right)_T = V > 0$$




chemical potential and phase equilibrium : $\left(\frac{\partial \mu}{\partial T}\right)_p = -S_m$



sublimation before melting



$$\left(\frac{\partial G}{\partial p}\right)_T = V > 0$$



Gibbs-Helmholtz equation :

$$\left(\frac{\partial G}{\partial T}\right)_p = \frac{G-H}{T}$$

Use this to show : $\left(\frac{\partial\left[\frac{G}{T}\right]}{\partial T}\right)_p = -\frac{H}{T^2}$


$$\boxed{\left(\frac{\partial\left[\frac{\Delta G}{T}\right]}{\partial T}\right)_p = -\frac{\Delta H}{T^2}}$$

Change in G with p : $G(p_f) = G(p_i) + \int_{p_i}^{p_f} V dp$

Change in G with p : $G(p_f) = G(p_i) + \int_{p_i}^{p_f} V dp$


for condensed phases

gases




Change in G with p : $G(p_f) = G(p_i) + \int_{p_i}^{p_f} V dp$

for condensed phases




gases




Change in G with p : $G(p_f) = G(p_i) + \int_{p_i}^{p_f} V dp$

for condensed phases




$$G_m(p_f) = G_m(p_i) + V_m (p_f - p_i)$$

gases



$$G_m(p_f) = G_m(p_i) + RT \int_{p_i}^{p_f} \frac{dp}{p} = G_m(p_i) + RT \ln \frac{p_f}{p_i}$$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{p}{p^\ominus}$



molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

$$\int_{p'}^p V_m dp = G_m(p) - G_m(p') = \left\{ G_m^\ominus + RT \ln \frac{f}{p^\ominus} \right\} - \left\{ G_m^\ominus + RT \ln \frac{f'}{p^\ominus} \right\}$$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

$$\int_{p'}^p V_m dp = G_m(p) - G_m(p') = \left\{ G_m^\ominus + RT \ln \frac{f}{p^\ominus} \right\} - \left\{ G_m^\ominus + RT \ln \frac{f'}{p^\ominus} \right\}$$

$$\int_{p'}^p V_{\text{perfect}, m} dp = RT \int_{p'}^p \frac{dp}{p} = RT \ln \frac{p}{p'}$$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

$$\int_{p'}^p V_m dp = G_m(p) - G_m(p') = \left\{ G_m^\ominus + RT \ln \frac{f}{p^\ominus} \right\} - \left\{ G_m^\ominus + RT \ln \frac{f'}{p^\ominus} \right\}$$

$$\int_{p'}^p V_{\text{perfect}, m} dp = RT \int_{p'}^p \frac{dp}{p} = RT \ln \frac{p}{p'}$$

$$\int_{p'}^p (V_m - V_{\text{perfect}, m}) dp = RT \left(\ln \frac{f}{f'} - \ln \frac{p}{p'} \right)$$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

$$\int_{p'}^p V_m dp = G_m(p) - G_m(p') = \left\{ G_m^\ominus + RT \ln \frac{f}{p^\ominus} \right\} - \left\{ G_m^\ominus + RT \ln \frac{f'}{p^\ominus} \right\}$$

$$\int_{p'}^p V_{\text{perfect}, m} dp = RT \int_{p'}^p \frac{dp}{p} = RT \ln \frac{p}{p'}$$

$$\int_{p'}^p (V_m - V_{\text{perfect}, m}) dp = RT \left(\ln \frac{f}{f'} - \ln \frac{p}{p'} \right)$$

$$\text{or, } \ln \frac{f}{p} - \ln \frac{f'}{p'} = \frac{1}{RT} \int_{p'}^p (V_m - V_{\text{perfect}, m}) dp$$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

$$\int_{p'}^p V_m dp = G_m(p) - G_m(p') = \left\{ G_m^\ominus + RT \ln \frac{f}{p^\ominus} \right\} - \left\{ G_m^\ominus + RT \ln \frac{f'}{p^\ominus} \right\}$$

$$\int_{p'}^p V_{\text{perfect}, m} dp = RT \int_{p'}^p \frac{dp}{p} = RT \ln \frac{p}{p'}$$

$$\int_{p'}^p (V_m - V_{\text{perfect}, m}) dp = RT \left(\ln \frac{f}{f'} - \ln \frac{p}{p'} \right)$$

$$\text{or, } \ln \frac{f}{p} - \ln \frac{f'}{p'} = \frac{1}{RT} \int_{p'}^p (V_m - V_{\text{perfect}, m}) dp$$

$$\lim_{p \rightarrow 0} \frac{f}{p} \rightarrow 1$$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

$$\int_{p'}^p V_m dp = G_m(p) - G_m(p') = \left\{ G_m^\ominus + RT \ln \frac{f}{p^\ominus} \right\} - \left\{ G_m^\ominus + RT \ln \frac{f'}{p^\ominus} \right\}$$

$$\int_{p'}^p V_{\text{perfect}, m} dp = RT \int_{p'}^p \frac{dp}{p} = RT \ln \frac{p}{p'}$$

$$\int_{p'}^p (V_m - V_{\text{perfect}, m}) dp = RT \left(\ln \frac{f}{f'} - \ln \frac{p}{p'} \right)$$

$$\text{or, } \ln \frac{f}{p} - \ln \frac{f'}{p'} = \frac{1}{RT} \int_{p'}^p (V_m - V_{\text{perfect}, m}) dp$$

$$\lim_{p \rightarrow 0} \frac{f}{p} \rightarrow 1 \implies \ln \frac{f}{p} = \frac{1}{RT} \int_0^p (V_m - V_{\text{perfect}, m}) dp$$

molar Gibbs energy vs p : $G_m(p) = G_m^\ominus + RT \ln \frac{f}{p^\ominus}$; f (fugacity) = $\underbrace{\phi}_{\text{fugacity coeff}} \cdot p$

$$\int_{p'}^p V_m dp = G_m(p) - G_m(p') = \left\{ G_m^\ominus + RT \ln \frac{f}{p^\ominus} \right\} - \left\{ G_m^\ominus + RT \ln \frac{f'}{p^\ominus} \right\}$$


$$\int_{p'}^p V_{\text{perfect}, m} dp = RT \int_{p'}^p \frac{dp}{p} = RT \ln \frac{p}{p'}$$

$$\int_{p'}^p (V_m - V_{\text{perfect}, m}) dp = RT \left(\ln \frac{f}{f'} - \ln \frac{p}{p'} \right)$$

$$\text{or, } \ln \frac{f}{p} - \ln \frac{f'}{p'} = \frac{1}{RT} \int_{p'}^p (V_m - V_{\text{perfect}, m}) dp$$

$$\lim_{p' \rightarrow 0} f \rightarrow p \implies \ln \frac{f}{p} = \frac{1}{RT} \int_0^p (V_m - V_{\text{perfect}, m}) dp$$


$$\text{or, } \ln \phi = \int_0^p \frac{Z-1}{p} dp$$



phase : form of matter that is uniform throughout in chemical composition and physical state

$P = \#$ phases in a system

Phase equilibria - triple point (T_3) and critical point (T_c)




Along each curve, $P=2$
at triple point, $P=3$

phase : form of matter that is uniform throughout in chemical composition and physical state

$P = \#$ phases in a system

Phase equilibria - triple point (T_3) and critical point (T_c)



For water : $T_3 = 273.16$ K and 611 Pa (6.11 mbar)

- ▶ Triple point marks the lowest pressure at which a liquid phase can exist
- ▶ If the slope of the solid–liquid phase boundary is positive
 - ▶ then triple point also marks the lowest temperature at which the liquid can exist
- ▶ the critical temperature is the upper limit

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

Phase Rule : $F = C - P + 2$

$F = \# \text{degrees of freedom}$, $C = \# \text{components}$, $P = \# \text{phases}$

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

phases α and β in equilibr. at given p , T

$$\mu_{\alpha}(p, T) = \mu_{\beta}(p, T)$$

eqn. relates p and T ,

so only one variable is independent

$$\text{Phase Rule : } F = C - P + 2$$

F = #degrees of freedom, C = #components, P = #phases

single-component

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

phases α and β in equilibr. at given p , T

$$\mu_{\alpha}(p, T) = \mu_{\beta}(p, T)$$

eqn. relates p and T ,

so only one variable is independent

$$\text{Phase Rule : } F = C - P + 2$$

$F = \# \text{degrees of freedom}$, $C = \# \text{components}$, $P = \# \text{phases}$

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

phases α and β in equilibr. at given p , T

$$\mu_{\alpha}(p, T) = \mu_{\beta}(p, T)$$

eqn. relates p and T ,

so only one variable is independent

single-component
three-phases

$$\text{Phase Rule : } F = C - P + 2$$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

phases α and β in equilibr. at given p , T

$$\mu_{\alpha}(p, T) = \mu_{\beta}(p, T)$$

eqn. relates p and T ,

so only one variable is independent

single-component

three-phases

$$F = 1 - 3 + 2 = 0$$

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

phases α and β in equilibr. at given p , T

$$\mu_{\alpha}(p, T) = \mu_{\beta}(p, T)$$

eqn. relates p and T ,

so only one variable is independent

single-component

three-phases

$$F = 1 - 3 + 2 = 0$$

phases α , β and γ in equilibr. at
given p , T ,

$$\begin{aligned}\mu(\alpha, p, T) &= \mu(\beta, p, T) \\ &= \mu(\gamma, p, T)\end{aligned}$$

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

phases α and β in equilibr. at given p , T

$$\mu_{\alpha}(p, T) = \mu_{\beta}(p, T)$$

eqn. relates p and T ,

so only one variable is independent

single-component

three-phases

$$F = 1 - 3 + 2 = 0$$

phases α , β and γ in equilibr. at
given p , T ,

$$\begin{aligned}\mu(\alpha, p, T) &= \mu(\beta, p, T) \\ &= \mu(\gamma, p, T)\end{aligned}$$

two eqns. for two unknowns p
and T

Phase Rule : $F = C - P + 2$

F = #degrees of freedom, C = #components, P = #phases

single-component, single-phase

$$F = 1 - 1 + 2 = 2$$

single-component, two-phases

$$F = 1 - 2 + 2 = 1$$

phases α and β in equilibr. at given p , T

$$\mu_{\alpha}(p, T) = \mu_{\beta}(p, T)$$

eqn. relates p and T ,

so only one variable is independent

single-component

three-phases

$$F = 1 - 3 + 2 = 0$$

phases α , β and γ in equilibr. at
given p , T ,


$$\begin{aligned}\mu(\alpha, p, T) &= \mu(\beta, p, T) \\ &= \mu(\gamma, p, T)\end{aligned}$$

two eqns. for two unknowns p
and T

fixed soln.

so no variation possible

Typical phase diagram




Experimental phase diagrams



H₂O

CO₂

He






Experimental phase diagrams




He


Experimental phase diagrams



Phase transition : Thermal analysis and Cooling curve



Phase transition : Thermal analysis and Cooling curve



metastable phases : Kinetic barriers



Response of melting to applied pressure

Response of melting to applied pressure

pressure-dependence of chemical potential : $\left(\frac{\partial \mu}{\partial p}\right)_T = V_m$

usual liquids : $V_m(l) > V_m(s)$

water : $V_m(l) < V_m(s)$



$$G_m(p) = G_m^\ominus + RT \ln \frac{p}{p^\ominus}$$

$$\Rightarrow \mu(p) = \mu^\ominus + RT \ln \frac{p}{p^\ominus} \stackrel{p^\ominus=1}{=} \mu^\ominus + RT \ln p$$

location of phase boundaries : $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T = \frac{\Delta_{trs} S}{\Delta_{trs} V}$

location of phase boundaries : $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T = \frac{\Delta_{\text{trs}}S}{\Delta_{\text{trs}}V}$

solid/liq. : $\frac{dp}{dT} = \frac{\Delta_{\text{fus}}H}{T\Delta_{\text{fus}}V}$

location of phase boundaries : $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T = \frac{\Delta_{trs}S}{\Delta_{trs}V}$

solid/liq. : $\frac{dp}{dT} = \frac{\Delta_{fus}H}{T\Delta_{fus}V}$

When pressure is applied to a system with two phases in equilibrium (at a), the equilibrium is disturbed

It can be restored by changing T , moving state of system to b

location of phase boundaries : $\left(\frac{\partial p}{\partial T}\right)_V = \left(\frac{\partial S}{\partial V}\right)_T = \frac{\Delta_{trs}S}{\Delta_{trs}V}$

solid/liq. : $\frac{dp}{dT} = \frac{\Delta_{fus}H}{T\Delta_{fus}V}$

When pressure is applied to a system with two phases in equilibrium (at a), the equilibrium is disturbed

It can be restored by changing T , moving state of system to b

relation between dp and dT ensures that system remains in equilibrium as either variable is changed

