

# A practical evaluation of recent methods in high-dimensional inference

Charles Zheng

Stanford University

May 3, 2015

# Theory and Practice

$$\underbrace{Y \sim N(\beta'X, \sigma^2 I)}_{\text{Theory}} \quad \Bigg| \quad \underbrace{Y, X_1, \dots, X_p}_{\text{Practice}} \text{ unknown relationship}$$

$$\begin{array}{l} \text{Theory} \\ Y \sim N(\beta'X, \sigma^2 I) \\ X_i = \begin{cases} \text{non-null} & \beta_i \neq 0 \\ \text{null} & \beta_i = 0 \end{cases} \end{array}$$

$$\begin{array}{l} \text{Practice} \\ Y, X_1, \dots, X_p \text{ unknown relationship} \\ X_i = \begin{cases} \text{interesting} \\ \text{uninteresting} \\ ??? \end{cases} \end{array}$$

	Control	$p \leq n$	$p > n$
Classical inference (Pearson 1930)	Marginal	Yes	
Covariance test (Lockhart et al. 2014)	?	Yes	Yes
Debiased lasso (Javanmard et al. 2014)	Marginal		Yes
Knockoffs (Barber et al. 2014)	FDR	Yes	?

But what's actually used in practice?

	Control	$p \leq n$	$p > n$
Classical inference (Pearson 1930)	Marginal	Yes	
Covariance test (Lockhart et al. 2014)	?	Yes	Yes
Debiased lasso (Javanmard et al. 2014)	Marginal		Yes
Knockoffs (Barber et al. 2014)	FDR	Yes	?
<b>Marginal screening</b>	?	Yes	Yes

# Regression vs Marginal Screening


- Why not just test  $H_i : \text{Cov}(X_i, Y) \neq 0$ ?
- *Even if  $Y = X\beta + \epsilon$ , most non-null  $X_i$  are probably also correlated*

# Regression vs Marginal Screening

- Why not just test  $H_i : \text{Cov}(X_i, Y) \neq 0$ ?
- Even if  $Y = X\beta + \epsilon$ , most non-null  $X_i$  are probably also correlated
- In “big data” many  $X_i$  are correlated to  $Y$ , but *redundant*

# Regression vs Marginal Screening

## Genome-wide association study



(Adapted from *Mourad 2012*)



# From theory to practice

## *Theory*

- Theory of inference in linear model

## *Practice*

# From theory to practice

## *Theory*

- Theory of inference in linear model

- Validation of given procedure in real data with ground truth

## *Practice*

# From theory to practice

## *Theory*

- Theory of inference in linear model
- Theory of robust inference

- Validation of given procedure in real data with ground truth

## *Practice*

# From theory to practice

## *Theory*

- Theory of inference in linear model
  - Theory of robust inference
  - Simulation studies
- 
- Validation of given procedure in real data with ground truth

## *Practice*

# From theory to practice

## *Theory*

- Theory of inference in linear model
- Theory of robust inference
- Simulation studies
- Validation on real data + **synthetic negative controls**
- Validation of given procedure in real data with ground truth

## *Practice*

- Difficult to validate inference procedures, because we would need to know the 'true'  $\beta$
- What is the 'true'  $\beta$  when the linear model is incorrect? We take


$$\beta = \mathbf{E}[\mathbf{x}\mathbf{x}^T]^{-1}\mathbf{E}[\mathbf{y}\mathbf{x}]$$

(the 'superpopulation' model)

- We don't know the ground truth in real data... what's the next best thing?

# Idea

I give you real data *mixed in* with noise variables



- Can you identify the original columns from the noise columns?
- I can test your procedure this way, because I know the ground truth!

# Synthetic Negative Controls

- Given random vector  $x \in \mathbb{R}^p$ , *define*  $\tilde{x} \in \mathbb{R}^{p+q}$  by

$$\tilde{x} = \begin{pmatrix} I \\ \Gamma \end{pmatrix} x + e$$

where  $\Gamma$  is a fixed matrix and  $e \perp x, y$ .



# Synthetic Negative Controls

- Given random vector  $x \in \mathbb{R}^p$ , *define*  $\tilde{x} \in \mathbb{R}^{p+q}$  by

$$\tilde{x} = \begin{pmatrix} I \\ \Gamma \end{pmatrix} x + e$$

where  $\Gamma$  is a fixed matrix and  $e \perp x, y$ .

- Let

$$\beta = \mathbf{E}[xx^T]^{-1}\mathbf{E}[yx], \quad \tilde{\beta} = \mathbf{E}[\tilde{x}\tilde{x}^T]^{-1}\mathbf{E}[y\tilde{x}]$$

# Synthetic Negative Controls

- Given random vector  $x \in \mathbb{R}^p$ , *define*  $\tilde{x} \in \mathbb{R}^{p+q}$  by

$$\tilde{x} = \begin{pmatrix} I \\ \Gamma \end{pmatrix} x + e$$

where  $\Gamma$  is a fixed matrix and  $e \perp x, y$ .

- Let

$$\beta = \mathbf{E}[xx^T]^{-1}\mathbf{E}[yx], \quad \tilde{\beta} = \mathbf{E}[\tilde{x}\tilde{x}^T]^{-1}\mathbf{E}[y\tilde{x}]$$

- Then

$$\forall i \in \{1, \dots, p\} : \beta_i = \tilde{\beta}_i$$

$$\forall i \in \{p+1, \dots, p+q\} : \tilde{\beta}_i = 0$$

# Synthetic Negative Controls

- Given random vector  $x \in \mathbb{R}^p$ , *define*  $\tilde{x} \in \mathbb{R}^{p+q}$  by

$$\tilde{x} = \begin{pmatrix} I \\ \Gamma \end{pmatrix} x + e$$

where  $\Gamma$  is a fixed matrix and  $e \perp x, y$ .

- Let

$$\beta = \mathbf{E}[xx^T]^{-1}\mathbf{E}[yx], \quad \tilde{\beta} = \mathbf{E}[\tilde{x}\tilde{x}^T]^{-1}\mathbf{E}[y\tilde{x}]$$

- Then

$$\forall i \in \{1, \dots, p\} : \beta_i = \tilde{\beta}_i$$

$$\forall i \in \{p+1, \dots, p+q\} : \tilde{\beta}_i = 0$$

- Special case.*  $X_{p+1}, \dots, X_{p+q}$  are pure noise: this is when  $\Gamma = 0$

# Using SNCs to investigate robustness

- All methods considered depend on strong assumptions (e.g. linearity, Gaussian iid errors, sparsity)
- How well do these methods work on real data where assumptions are most likely violated?

# Using SNCs to investigate robustness

- All methods considered depend on strong assumptions (e.g. linearity, Gaussian iid errors, sparsity)
- How well do these methods work on real data where assumptions are most likely violated?
- Take low-dimensional real data mixed with SNCs (synthetic negative controls): can we identify the real data while controlling Type I error (measured by rejections of SNCs)?

# What can we conclude?

- Experiments using SNCs shows that we can do well on the *hypothesis testing problem* in realistic settings, where assumptions are violated
- However, these experiments cannot tell us if we are solving the right problem!
- Is the *hypothesis testing problem* even relevant for the application?  
The only way to tell is validation on real, high-dimensional data with application-specific ground truth.

*“Statistics is a science in my opinion... for if its methods fail the test of experience – not the test of logic – they are discarded.”*

*“ Both the statistician and the client must learn to confront the uncertainties of the world more explicitly, ... never to avoid responsibility for an ever-present understanding that all assumptions underlying data analysis are always approximations. Above all, they must base their thinking on a recognition that their assumptions will always require review and reappraisal... ”*

– John Tukey

- Barber, R., and Candès, E. (2014). Controlling the False Discovery Rate via Knockoffs. arXiv Preprint arXiv:1404.5609, 127. Retrieved from <http://arxiv.org/abs/1404.5609>
- Javanmard, A., and Montanari, A. (2014). Confidence intervals and hypothesis testing for high-dimensional regression. The Journal of Machine Learning Research, 15, 2869-2909. Retrieved from <http://dl.acm.org/citation.cfm?id=2697057>
- Lockhart, R., Taylor, J., Tibshirani, R. J., and Tibshirani, R. (2014). A Significance Test for the Lasso. Annals of Statistics, 42(2), 413-468. doi:10.1214/13-AOS1175



# Acknowledgements

Thanks to Will Fithian for useful discussions.