

A practical evaluation of recent methods in high-dimensional inference

Charles Zheng

Stanford University

May 6, 2015


Problem and motivation

- $x \in \mathbb{R}^p, y \in \mathbb{R}$ have a joint distribution P where $y|x \sim N(x^T \beta, \sigma^2)$
- Observe $X = (x_1, \dots, x_n)^T$, $Y = (y_1, \dots, y_n)$ iid
- Problem: test $H_i : \beta_0 = i$ for $i = 1, \dots, p$
- Motivation: x are SNPs (mutations), y is phenotype

	Control	$p > n$
Classical inference (Pearson 1930)	Marginal	No
Covariance test (Lockhart et al. 2014)	FWER?	Yes
Debiased lasso (Javanmard et al. 2014)	Marginal	Yes
Knockoffs (Barber et al. 2014)	FDR	?

The LASSO path

$$\hat{\beta}_\lambda = \operatorname{argmin}_\beta \frac{1}{2} \|X\beta - Y\|^2 + \lambda \|\beta\|_1$$



(Image credit: ??)


Covariance test

- (2014) Lockhart, Taylor, Tibshirani ($\times 2$)
- Standard assumptions $Y \sim N(X\beta, \sigma^2 I) + \text{large } p \text{ asymptotics}$
- See *also* non-asymptotic exact test (Lee, Sun $\times 2$, Taylor 2015)

Step	Predictor entered	Forward stepwise	Lasso
1	lcavol	0.000	0.000
2	lweight	0.000	0.052
3	svi	0.041	0.174
4	lbph	0.045	0.929
5	pgg45	0.226	0.353
6	age	0.191	0.650
7	lcp	0.065	0.051
8	gleason	0.883	0.978


Debiased regularized M-estimators

- (2014) Javanmard and Montanari
- Standard assumptions + sparsity condition on β + large n and p asymptotics



Knockoff filter

- (2014) Barber and Candés
- *Finite sample* $Y \sim N(X\beta, \sigma^2 I)$, $n \leq p$, control FDR
- Extension to $p > n$, FWER control, etc. forthcoming...




lweight	22.5652
lcavol	20.5199
svi	4.4871
lbph	1.1865
age	0.0829
gleason	0.0387
lcp	-0.2359
pgg45	-3.3742

But what's actually used in practice?

	Control	$p > n$
Classical inference (Pearson 1930)	Marginal	No
Covariance test (Lockhart et al. 2014)	FWER?	Yes
Debiased lasso (Javanmard et al. 2014)	Marginal	Yes
Knockoffs (Barber et al. 2014)	FDR	?
Marginal screening	???	Yes

Regression vs Marginal Screening

Testing $H_i : \beta_i = 0$ is better than testing $H_i : \text{Cov}(X_i, Y) = 0$ when you are looking for X_i *directly* linked to Y



(Adapted from *Mourad 2012*)

Statistical Validation

- These procedures are derived under strong assumptions (linearity, gaussianity, homoscedasticity)
- How well do they work in real data where these assumptions are violated?
- We could validate inference procedures in real data if only we knew the '*true*' β , defined as

$$\beta = \mathbf{E}[\mathbf{x}\mathbf{x}^T]^{-1}\mathbf{E}[\mathbf{y}\mathbf{x}]$$

Statistical Validation

- These procedures are derived under strong assumptions (linearity, gaussianity, homoscedasticity)
- How well do they work in real data where these assumptions are violated?
- We could validate inference procedures in real data if only we knew the 'true' β , defined as

$$\beta = \mathbf{E}[\mathbf{xx}^T]^{-1} \mathbf{E}[\mathbf{yx}]$$


- Possibility: take a dataset with large p and *humongous* n , so we can get an extremely precise estimate of β using OLS. Then test the high-dimensional inference procedures on subsamples of size $n_0 \leq p < n$ of the data

Example: personality data

- Data with $p = 163$ survey questions from an online personality test, $n = 49086$ (after processing)
- Predict self-reported age of respondent, y , from their responses
- Is n large enough for us to confidently say which $\beta_i = 0$ (for use as ground truth?)

Example: personality data


Coefficient estimates ± 3 sd



Consider declaring all variables whose intervals cross 0 to be null. Then $p_1 = 105$ (out of 163)


Example: personality data

- If n were large enough, then for the selected model S we should have $\hat{y} = \sum_{i=1}^p X_i \hat{\beta}_i$ close to $\hat{y}_S = \sum_{i \in S} X_i \hat{\beta}_i$
- But...



Example: personality data

- Here n is not large enough for $p = 163$
- If we reduce the dimensionality to 15 by subsampling columns, it looks more convincing that we selected the correct 10 variables



- It is by no means *impossible* to get large enough data to estimate high-dimensional β , with say, $p > 100$
- But if were *easy* to get such large n data... we wouldn't need these new inference techniques in the first place!

Why not use simulations?

- Simulations can be used to test robustness of the procedure
- In simulations, we can add all the nonlinearities, nongaussianity, etc. that we want

Why not use simulations?


- Advantage: In simulations, we not only know β , but exactly how the data is generated
- Advantage: We can vary simulation parameters and get a lot of insight about the procedure being tested

Why not use simulations?

- Advantage: In simulations, we not only know β , but exactly how the data is generated
- Advantage: We can vary simulation parameters and get a lot of insight about the procedure being tested
- **Disadvantage:** Are these simulations relevant? How can we tell the simulated models are realistic?

Idea


I give you real data *mixed in* with noise variables



- Can you identify the original columns from the noise columns?


Idea

I give you real data *mixed in* with noise variables




- Can you identify the original columns from the noise columns?
- I can test your procedure this way, because I know the ground truth!

I give you real data *mixed in* with noise variables



- Can you identify the original columns from the noise columns?
- I can test your procedure this way, because I know the ground truth!
- **Caveat:** this test is unrealistically 'easy' (due to lack of correlations)

Synthetic Negative Controls



- Synthetic negative controls (SNCs) are artificial columns *which are correlated* to X , yet still have zero (population) regression coefficients
- Suppose I give you real data + SNCs, then you apply high-dimensional inference. If you reject any SNCs, we know these are errors!
- This gives us some measure of performance on “real” data (maybe?)

Synthetic Negative Controls

- Given random vector $x \in \mathbb{R}^p$, let e be noise in \mathbb{R}^p independent of x .
- Let Γ be a fixed $p \times q$ matrix. *Define* synthetic negative controls $z \in \mathbb{R}^q$ by

$$z = x'\Gamma + e$$

and let $\tilde{x} = (x, z)$, so that

$$\tilde{x}_1 = x_1, \dots, \tilde{x}_p = x_p$$

$$\tilde{x}_{p+1} = z_1, \dots, \tilde{x}_{p+q} = z_q$$

Synthetic Negative Controls

- Given random vector $x \in \mathbb{R}^p$, let e be noise in \mathbb{R}^p independent of x .
- Let Γ be a fixed $p \times q$ matrix. Define synthetic negative controls $z \in \mathbb{R}^q$ by

$$z = x'\Gamma + e$$

and let $\tilde{x} = (x, z)$, so that

$$\tilde{x}_1 = x_1, \dots, \tilde{x}_p = x_p$$

$$\tilde{x}_{p+1} = z_1, \dots, \tilde{x}_{p+q} = z_q$$

- Let

$$\beta = \mathbf{E}[xx^T]^{-1}\mathbf{E}[yx], \quad \tilde{\beta} = \mathbf{E}[\tilde{x}\tilde{x}^T]^{-1}\mathbf{E}[y\tilde{x}]$$

Synthetic Negative Controls

- Given random vector $x \in \mathbb{R}^p$, let e be noise in \mathbb{R}^p independent of x .
- Let Γ be a fixed $p \times q$ matrix. Define synthetic negative controls $z \in \mathbb{R}^q$ by

$$z = x'\Gamma + e$$

and let $\tilde{x} = (x, z)$, so that

$$\tilde{x}_1 = x_1, \dots, \tilde{x}_p = x_p$$

$$\tilde{x}_{p+1} = z_1, \dots, \tilde{x}_{p+q} = z_q$$

- Let

$$\beta = \mathbf{E}[xx^T]^{-1}\mathbf{E}[yx], \quad \tilde{\beta} = \mathbf{E}[\tilde{x}\tilde{x}^T]^{-1}\mathbf{E}[y\tilde{x}]$$

- Then

$$\forall i \in \{1, \dots, p\} : \beta_i = \tilde{\beta}_i$$

$$\forall i \in \{p+1, \dots, p+q\} : \tilde{\beta}_i = 0$$

Why is this...?

- Recall that $\hat{\beta}_i$ is the *univariate regression* coefficient of Y on $X_{i|-i}$, where $X_{i|-i}$ is the *residual* of X_i after X_i is regressed on the other columns..
- Population version: $\beta_i = 0$ if the projection of X_i on the null space of the other covariates is uncorrelated with Y

Why is this...?

- Population version: $\beta_i = 0$ if the projection of X_i on the null space of the other covariates is uncorrelated with Y
- For $i = 1, \dots, q$, we have

$$\tilde{X}_{p+i} = x' \Gamma_i + E_i$$

where here \tilde{X}_{p+1} denotes the random variable (not the column of the design matrix)

Why is this...?

- Population version: $\beta_i = 0$ if the projection of X_i on the null space of the other covariates is uncorrelated with Y
- For $i = 1, \dots, q$, we have

$$\tilde{X}_{p+i} = x' \Gamma_i + E_i$$

where here \tilde{X}_{p+1} denotes the random variable (not the column of the design matrix)

- The orthogonal projection P_X^\perp of \tilde{X}_{p+1} is

$$P_X^\perp \tilde{X} = P_X^\perp X \Gamma_i + P_X^\perp E_i = 0 + E_i$$

since $P_X^\perp X = 0$; meanwhile since $E_i \perp X$, $P_X^\perp E_i = E_i$.

Why is this...?

- Population version: $\beta_i = 0$ if the projection of X_i on the null space of the other covariates is uncorrelated with Y

- The orthogonal projection P_X^\perp of \tilde{X}_{p+1} is

$$P_X^\perp \tilde{X} = P_X^\perp X \Gamma_i + P_X^\perp E_i = 0 + E_i$$

since $P_X^\perp X = 0$; meanwhile since $E_i \perp X$, $P_X^\perp E_i = E_i$.

- Since $E_i \perp y$, we have $\text{Cor}(P_X^\perp \tilde{X}_{p+i}, y) = 0$, hence $\tilde{\beta}_{p+i} = 0$

Why is this...?

- Population version: $\beta_i = 0$ if the projection of X_i on the null space of the other covariates is uncorrelated with Y
- Since $E_i \perp y$, we have $\text{Cor}(P_X^\perp \tilde{X}_{p+i}, y) = 0$, hence $\tilde{\beta}_{p+i} = 0$
- And since $\tilde{\beta}_j = 0$ for all the added variables $j = p+1, \dots, p+q$, it follows that $\tilde{\beta}_i$ is unchanged for $i = 1, \dots, p$.

Using SNCs to evaluate procedures

- Take low-dimensional real data mixed with SNCs (synthetic negative controls), apply inference procedure
- *Proxy for Type I error*: Rejected SNCs
- *Proxy for Power*: Rejected original variables

A step-by-step tutorial (in R)

1. Take the prostate data

```
> data(prostate)
> x <- prostate[, 1:8]
> y <- prostate[, 9]
> colnames(x)
[1] "lcavol" "lweight" "age" "lbph" "svi"
    "lcp" "gleason" "pgg45"
> dim(x)
[1] 97 8
```

A step-by-step tutorial

2. Construct 20 synthetic negative controls

```
> GAMMA <- matrix(rnorm(8 * 20), 8, 20)
> E <- matrix(rnorm(97 * 20), 97, 20)
> sncs <- as.matrix(x) %*% GAMMA + 2 * E
> sncs <- data.frame(sncs)
> colnames(sncs)
[1] "X1"  "X2"  "X3"  "X4"  "X5"  "X6"  ...
[19] "X19" "X20"
```

3. Create combined design matrix

```
> x2 <- cbind(x, sncs)
```

A step-by-step tutorial

4. Try marginal screening

```
> cors <- cor(x2, y)
> cors[order(-abs(cors)), , drop = F]
      [,1]
lcavol  0.7344603
svi      0.5662182
lcp      0.5488132
X6       -0.4591506
X16      0.4482263
lweight  0.4333194
X4       -0.4326898
```

A step-by-step tutorial

5. Try covariance test

```
> library(covTest)
> covTest(lars(as.matrix(x2), y), as.matrix(x2), y)
$results
```

Predictor_Number	Drop_in_covariance	P-value
1	69.0292	0.0000
5	1.5390	0.2219
2	6.8094	0.0020
11	0.8559	0.4294

(Numbers 1, 5, 2 are original, 11 is a SNC)

A step-by-step tutorial

6. Try debiased lasso (code at <http://web.stanford.edu/~montanar/sslasso/>)

```
> res <- SSLasso(as.matrix(x2), y)
[1] "10% done"
...
[1] "90% done"
> rej <- (res$up < 0) | (res$low > 0)
> names(x2)[rej]
[1] "lcavol" "lweight" "svi"
```

A step-by-step tutorial

7. Try knockoffs

```
> library(knockoff)
```

```
> knockoff.filter(x2, y)
```

Call:

```
knockoff.filter(X = x2, y = y)
```

Selected variables:

lweight	X7
2	15

Disclaimer!

- I am *not* proposing SNCs as a methodology for *inference*
- There is a danger of inferring *that Type I error has been controlled* from lack of rejection of SNCs. There are no formal guarantees of this!
- One should interpret results from experiments with SNCs in the same way one interprets simulation results with purely synthetic data


More Experiments!

Data	n	p_1	Linear?	Gaussian?	Constant σ^2 ?
Personality	49k	163	No	No	No
fMRI	1750	44	No	OK	No
HIV	842	207	No	Yes?	OK?
Galaxy	323	4	No	OK	No


- We add $n/2 - p_1$ synthetic negative controls
- X is scaled, Γ is a gaussian matrix, $Var(E)$ is chosen to yield 'interesting' results
- Multiple trials averaging over the randomness of generating SNCs

Marginal Screening


Personality




fMRI



HIV




Galaxy



Legend: 0 = False positives, 1 = True positives


Ordinary Least Squares

“Rel. power” = $TP / (\text{max number of TPs at } \alpha = 0.5 \text{ for any method})$



Legend: **P** = Personality, **F** = fMRI, **H** = HIV, **G** = Galaxy



Debiased Lasso



Legend: **P** = Personality, **F** = fMRI, **H** = HIV, **G** = Galaxy



Covariance Test

Procedure: continue rejecting until $p > \alpha$, then accept all other variables



Legend: P = Personality, F = fMRI, H = HIV, G = Galaxy

Note: $FDR^* = \mathbf{E}[FP/(FP + TP + 1/\alpha)]$




Legend: **P** = Personality, **F** = fMRI, **H** = HIV, **G** = Galaxy

Variable Ranking Criteria


- Forget about Type I error for a second...
- Use procedures to *rank* variables by p-value
- Easy to compare procedures with different Type I criteria and also non-inference variable selection
- (Optional) score by Area Under Curve (AUC), etc.

Variable Ranking: Personality




Legend: o = OLS, c = covariance test, k = knockoff, s = debiased lasso, (line) = marginal screening, (line) = lasso path

Variable Ranking: fMRI




Legend: o = OLS, c = covariance test, k = knockoff, s = debiased lasso, (line) = marginal screening, (line) = lasso path

Variable Ranking: HIV



Legend: o = OLS, c = covariance test, k = knockoff, s = debiased lasso, (line) = marginal screening, (line) = lasso path

Variable Ranking: Galaxy



Legend: o = OLS, c = covariance test, k = knockoff, s = debiased lasso, (line) = marginal screening, (line) = lasso path

What are we learning?

- *Claim:* Performance on data with *known active variables* + SNCs are informative of how inference procedures generally perform in similar data with *unknown active variables*

What are we learning?

- *Claim*: Performance on data with *known active variables* + SNCs are informative of how inference procedures generally perform in similar data with *unknown active variables*
- Example (GWAS): There is a new disease (say, liver cancer) where we have little prior info. I have data for a 'similar' disease, (say, stomach cancer) for which I have *partial information* about the relevant causes. In particular, I know *the most important* the genes involved stomach cancer
- Testing your procedure directly on the stomach cancer data is uninformative: if you reject a gene which is not *a priori* known to cause stomach cancer... it could still be a new discovery!

What are we learning?

- *Claim*: Performance on data with *known active variables* + SNCs are informative of how inference procedures generally perform in similar data with *unknown active variables*
- Example (GWAS): There is a new disease (say, liver cancer) where we have little prior info. I have data for a 'similar' disease, (say, stomach cancer) for which I have *partial information* about the relevant causes. In particular, I know *the most important* the genes involved stomach cancer
- Testing your procedure directly on the stomach cancer data is uninformative: if you reject a gene which is not *a priori* known to cause stomach cancer... it could still be a new discovery!
- Rather than test your inference procedure on the full stomach cancer data, test it on the subset of the *known genes* plus SNCs
- Tricky part: choosing the size of the noise for the SNCs. To be safe, try several noise levels.

How could this be useful?

- There are no formal guarantees... so what could we gain from using SNC experiments?

How could this be useful?

- There are no formal guarantees... so what could we gain from using SNC experiments?
- We can complement *theoretical guarantees* with *application-specific benchmarks*

How could this be useful?

- There are no formal guarantees... so what could we gain from using SNC experiments?
- We can complement *theoretical guarantees* with *application-specific benchmarks*
- Poor performance on benchmarks would tell us where our methods need improvement
 - Failure to control Type I error on benchmarks indicates a need for methods derived under weaker assumptions
 - Overly conservative Type I error control indicates a need for methods which are more adaptive to 'easy' cases

How could this be useful?

- There are no formal guarantees... so what could we gain from using SNC experiments?
- We can complement *theoretical guarantees* with *application-specific benchmarks*
- Poor performance on benchmarks would tell us where our methods need improvement
 - Failure to control Type I error on benchmarks indicates a need for methods derived under weaker assumptions
 - Overly conservative Type I error control indicates a need for methods which are more adaptive to 'easy' cases
- Possible to run a Kaggle-style competition for *inference* rather than prediction

How could this be useful?

- There are no formal guarantees... so what could we gain from using SNC experiments?
- We can complement *theoretical guarantees* with *application-specific benchmarks*
- Poor performance on benchmarks would tell us where our methods need improvement
 - Failure to control Type I error on benchmarks indicates a need for methods derived under weaker assumptions
 - Overly conservative Type I error control indicates a need for methods which are more adaptive to 'easy' cases
- Possible to run a Kaggle-style competition for *inference* rather than prediction
- Recognizing that different procedures can have differing strengths creates room for a diversity of approaches

Do SNCs reduce the need for scientific validation?

- The gold standard of validation: obtaining *scientific confirmation* of the *scientific hypotheses* suggested by the inference procedure
- The answer is no: scientific validation is as important as ever.

Do SNCs reduce the need for scientific validation?

- The gold standard of validation: obtaining *scientific confirmation* of the *scientific hypotheses* suggested by the inference procedure
- The answer is no: scientific validation is as important as ever.
- SNCs can be used to get an idea of our performance on the *regression hypothesis testing problem* in realistic settings
- However, how can we tell if the *regression hypothesis testing problem* is right framework for the real-world problem we are trying to solve? There may be better statistical frameworks, like graphical models, which are a better fit to the problem.

Do SNCs reduce the need for scientific validation?

- The gold standard of validation: obtaining *scientific confirmation* of the *scientific hypotheses* suggested by the inference procedure
- The answer is no: scientific validation is as important as ever.
- SNCs can be used to get an idea of our performance on the *regression hypothesis testing problem* in realistic settings
- However, how can we tell if the *regression hypothesis testing problem* is right framework for the real-world problem we are trying to solve? There may be better statistical frameworks, like graphical models, which are a better fit to the problem.
- Ultimately, we need the practitioner to measure the value of the results we obtain from the inference procedure.

“ Both the client and the statistician... must base their thinking on a recognition that their assumptions will always require review and reappraisal... ”

– John Tukey

Acknowledgements

Thanks to Will Fithian for useful discussions.

- Barber, R., and Candès, E. (2014). Controlling the False Discovery Rate via Knockoffs. arXiv Preprint arXiv:1404.5609, 127. Retrieved from <http://arxiv.org/abs/1404.5609>
- Javanmard, A., and Montanari, A. (2014). Confidence intervals and hypothesis testing for high-dimensional regression. The Journal of Machine Learning Research, 15, 2869-2909. Retrieved from <http://dl.acm.org/citation.cfm?id=2697057>
- Lockhart, R., Taylor, J., Tibshirani, R. J., and Tibshirani, R. (2014). A Significance Test for the Lasso. Annals of Statistics, 42(2), 413-468. doi:10.1214/13-AOS1175