

deeplearning.ai

One hidden layer Neural Network

Gradient descent for neural networks

Gradient descent for neural networks

Parameters:
$$(J^{(1)}, b^{(1)}, b^{(1)$$

Andrew Ng

Formulas for computing derivatives

Formal propagation!

$$Z^{(1)} = U_{(1)} \times U_{(1)}$$

$$A^{(2)} = g^{(1)} (Z^{(1)}) \leftarrow$$

$$Z^{(2)} = U_{(2)} A^{(2)} + U_{(2)}$$

$$A^{(2)} = g^{(2)} (Z^{(2)}) = G(Z^{(2)})$$

$$A^{(2)} = g^{(2)} (Z^{(2)}) = G(Z^{(2)})$$

Back propagation:

$$\begin{aligned}
&\mathcal{L}^{[2]} = \mathcal{L}^{[2]} - \mathcal{L}^{[1]} \\
&\mathcal{L}^{[1]} = \mathcal{L}^{[2]} - \mathcal{L}^{[1]} \\
&\mathcal{L}^{[1]} = \mathcal{L}^{[2]} - \mathcal{L}^{[1]} \\
&\mathcal{L}^{[1]} = \mathcal{L}^{[2]} - \mathcal{L}^{[1]} \\
&\mathcal{L}^{[2]} = \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} = \mathcal{L}^{[2]} - \mathcal{L}^{[2]} + \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} = \mathcal{L}^{[2]} - \mathcal{L}^{[2]} + \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} - \mathcal{L}^{[2]} - - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{[2]} \\
&\mathcal{L}^{[2]} - \mathcal{L}^{[2]} - \mathcal{L}^{$$