

- 1) How do β blockers improve cardiac function in HF ?
 - a- **By decreasing cardiac remodeling.**
 - a- By increasing heart rate.
 - b- By increasing renin release.
 - c- By activating norepinephrine.

- 2) K supplements should not be given if the following drug is prescribed:
 - a) **Enalapril**
 - b) Verapamil
 - c) Thiazide
 - d) Digoxin

- 3) First line of treatment of patients with elevated risk atherosclerotic CV diseases is:
 - a) **Statins**
 - b) Dabigatran
 - c) Alteplase
 - d) Ezetimibe

- 4) Early signs of digitalis toxicity is:
 - a) **Nausea & vomiting**
 - b) Yellow and green vision
 - c) Atrial arrhythmias
 - d) Heart block

- 5) Which is important to monitor in patients taking digoxin ?
 - a- Chloride.
 - b- **Potassium.**
 - c- Sodium.
 - d- Zinc.

- 6) Adverse effect associated with morphine includes:
 - a) Diarrhea
 - b) **Bronchoconstriction**
 - c) Lacrimation
 - d) Vasoconstriction

- 7) What is the most common adverse effect associated with fixed dose hydralazine/isosorbide dinitrate ?
 - a- Diarrhea.
 - b- Drug-induced lupus.
 - c- **Hypotension.**
 - d- Heartburn.

- 8) Variant angina is treated by the following combination:

A. Verapamil & Nitrate	C. Beta blocker & nifedipine
B. Beta blocker & nitrate	D. Nitrate & nifedipine

