

? Potential clinical complication of silicosis is -18

- a. Mycobacterium tuberculosis infection ☒
- b. Rheumatoid arthritis ☐
- c. Scleroderma ☐
- d. Corepulmonale ☐

محو التحديد

:Adverse effect associated with morphine includes .19

- a. Diarrhea ☐
- b. Bronchoconstriction ☒
- c. Lacrimation ☐
- d. Vasoconstriction ☐

محو التحديد

?What does hypoxia mean .20

- a. Increased hemoglobin in the blood. ☐
- b. Decreased oxygen reaching bodily tissues. ☒
- c. Decreased hemoglobin in the blood. ☐
- d. Increased oxygen reaching bodily tissues. ☐

محو التحديد

1. **Pulmonary veins differ from other veins in that they**

- a. Carry blood away from the heart ☐
- b. Carry deoxygenated blood ☐
- c. Carry oxygenated blood ☒
- d. Have cartilaginous walls ☐

محو التحديد

2. **As a risk reduction strategy, which one of the following vaccination should be recommended to all COPD patients beside a yearly influenza ?vaccination**

- a. Herpes zoster vaccine ☐
- b. Haemophilus influenza vaccine ☐
- c. Pneumncocal vaccination ☒
- d. Rubella vaccination ☐

محو التحديد

3- **What are lab test do you order on the sputum sample of a COPD patients**

- a. Culture and sensitivity ☐
- b. Culture and gram stain ☒
- c. Acid fast bacteria ☐
- d. Culture only ☐

محو التحديد

.4 : Thymoma is

- a. Anterior mediastinal lesion ☒
- b. Posterior mediastinal lesion ☐
- c. Middle mediastinal lesion ☐
- d. Superior mediastinal lesion ☐

محو التحديد

.5 A 75-year-old, male presented with dyspnea and chest pain. Chest X-ray showed unilateral pleural effusion. Aspiration of pleural fluid revealed exudative hemorrhagic pleural effusion. Which of the following findings ?coincident with mesothelioma

- a. Low glucose and Ph ☒
- b. High glucose and pH. ☐
- c. Low glucose and high pH. ☐
- d. High glucose and low pH. ☐

محو التحديد

.6 Pleural effusions that occur secondarily to ----- are most often .treated with pleurodesis

- a. Ascites. ☐
- b. Congestive heart failure. ☐
- c. Nephrotic syndrome. ☐
- d. Malignancy. ☒

محو التحديد

?Horner's syndrome is a complication of which of the following lesion .7

- a. Bronchial carcinoid ☐
- b. Right apical carcinoma ☒
- c. Apical lung abscess ☐
- d. Right basal carcinoma ☐

محو التحديد

?Which one of the followings lung cancer has a worse prognosis .8

- a. Adenocarcinoma. ☐
- b. Bronchioalveolar carcinoma. ☐
- c. Squamous cell carcinoma. ☐
- d. Small cell carcinoma. ☒

محو التحديد

**A 45-year-old, male, presented with fever, dyspnea and productive .9
cough. His chest X- ray detected lobar pneumonia. Ultrasound was
:recommended, which of the following is expected ultrasound findings**

- a. Absent pleural sliding/gliding ☐
- b. Absent comet tail artifacts ☐
- c. Horizontal reverberation artifacts ☐
- d. Air Bronchogram ☒

محو التحديد

A 15-year-old, male, basketball player presented at emergency department by severe stitching left sided chest pain and dyspnea. There was no available chest X-ray. Ultrasound was done and revealed absent left lung sliding with horizontal reverberation artifacts. The most probable :diagnosis is

- a. Pleural effusion ☐
- b. Pneumothorax ☒
- c. Pulmonary embolism ☐
- d. Pneumonia ☐

محو التحديد

:Forced Vital Capacity FVC is defined as -11

- a. Maximal amount of air that the patient can forcibly inhale after a maximal exhalation ☐
- b. Maximal amount of air that the patient can forcibly exhale after taking a maximal inhalation ☒
- c. Maximal amount of air that the patient can forcibly exhale after taking a maximal exhalation ☐
- d. Maximal amount of air that the patient can forcibly inhale after taking a maximal inhalation ☐
- Option 5 ☐

محو التحديد

:Forced Expiratory Volume in 1st second FEV1 is defined .12

- a. Volume exhaled in the first second ☒
- b. Volume inhaled in the first second ☐
- c. Volume inhaled and exhaled in the first second ☐
- d. Volume maximally inhaled in the first second ☐

محو التحديد

inhaled corticosteroids for the long-term treatment of asthma exhibit **.13**
:the following desired feature

- a. Enhance cellular response to inflammatory mediators. ☐
- b. Increase production of leukotriene ☐
- c. Mast cell stabilization ☒
- d. Relaxation of skeletal muscle ☐

محو التحديد

Regarding to the drug interaction of theophylline. The half-life of **.14**
?theophylline is decrease by which of the following

- a- Erythromycin ☐
- b. Phenobarbitone ☒
- c. Quinolones ☐
- d. Cimetidine ☐

محو التحديد

?Highest prevalence of sarcoidosis is reported in **-15**

- a. Nordic population ☒
- b. Hispanic population ☐
- c. Aborigines in Australia ☐
- d. Red Indian tribes ☐

محو التحديد

? Löfgren's syndrome is characterized by -16

- a. Arthritis ☐
- b. Erythema nodosum ☐
- c. Bilateral hilar adenopathy ☐
- d. All of the above ☒

محو التحديد

? Major health effect from exposure to asbestos is -17

- a. Pleural fibrosis ☐
- b. Diffuse interstitial pulmonary fibrosis ☐
- c. Cancers of respiratory tract, pleura & peritoneum ☐
- d. All of the above ☒

محو التحديد

? Potential clinical complication of silicosis is -18

- a. Mycobacterium tuberculosis infection ☒
- b. Rheumatoid arthritis ☐
- c. Scleroderma ☐
- d. Corepulmonale ☐

محو التحديد