

نقطتان (2)

* ?What does hypoxia mean (1

- A) Increased hemoglobin in the blood. ☐
- B) Decreased oxygen reaching bodily tissues. ☒
- C) Decreased hemoglobin in the blood. ☐
- D) Increased oxygen reaching bodily tissues. ☐

نقطتان (2)

* ?Shunt occurs when (2

- A) Arterial blood mixes with venous blood (left to right). ☐
- B) Venous blood mixes with arterial blood(right to left). ☒
- C)A combination of both conditions above. ☐
- D) Ventilation and perfusion are equal. ☐

نقطتان (2) *

Non- invasive ventilation is (3
most useful with which one of the
?following condition

docs.google.com

* نقطتان (2) Non- invasive ventilation is (3
most useful with which one of the
?following condition

- A) Acute exacerbation of COPD with CO₂ retention. ☒
- B) Pulmonary embolism with CO₂ retention. ☐
- C) Left ventricular failure with CO₂ retention. ☐
- D) Pneumonia with CO₂ retention. ☐

* نقطتان (2) Which of the following is not (4
considered a routine
investigation in patient with
.suspected respiratory failure

- A) Chest x ray. ☐
- B) Arterial blood gas analysis. ☐
- C) Right sided heart catheterization. ☒
- D) Electrocardiograph. ☐

* نقطتان (2)

A 63 years old man has had (5 progressively worsen dyspnea over the past 10 years. He has noticed a 5 Kg weight loss in the past 2 years. He has chronic cough with minimal sputum production and no chest pain. On physical examination, he is afebrile and normotensive. A chest radiograph shows extensive interstitial disease. Pulmonary function tests show diminished capacity, and restrictive dysfunction. Increased exposure to which of the following pollutants is most likely ?to produce these findings

A) Carbon monoxide. ☐

B) Ozone. ☐

C) Asbestosis. ☒

D) Tobacco smoke. ☐

* نقطتان (2)

Pulmonary hypertension is (6
present in which type of
?respiratory failure

- A) Hypoxemic respiratory failure. ☐
- B) Hypercapnic respiratory failure. ☐
- Acute respiratory failure. ☐
- D) Chronic respiratory failure. ☒

* نقطتان (2)

A 60 years old, constructor (7
worker smoker presented with
progressive dyspnea and a chest
x- ray showed left sided
moderate pleural effusion. Which
of the following is the next step in
?diagnostic approach

- A) Aspiration of pleural fluid sample for analysis. ☒
- B) Thoracoscopic biopsy. ☐
- C) CT chest. ☐
- D) PET scan. ☐



* نقطتان (2)

What is the most accurate (8
enzyme to rule out TB in case of
exudative lymphocytic pleural
?effusion

A) Amylase. ☐

B) Adenosine deaminase. ☒

C) Pepsin. ☐

Lipase. ☐

* نقطتان (2)

Which diagnostic method that (9
differentiate between
mesothelioma and
?adenocarcinoma

A) Pleural fluid chemical analysis. ☐

B) Computed tomography scan of chest ☐

C) Chest x-ray. ☐

Immunohistochemistry. ☒

* نقطتان (2)

Which of the following (10 spirometry parameters) is more sensitive marker of small airway airflow limitation

A) FVC. ☐

B) FEV1. ☐

C) FEV1\FVC. ☐

D) FEF 25-75. ☒