

★ نقطة واحدة

**The primary underlying cause of most cases of Ischemic Heart -1
:Disease (IHD) is**

- A. Coronary artery vasospasm ☐
- B. Atherosclerosis of the coronary arteries ☒
- C. Valvular heart disease ☐
- D. Congenital heart defects ☐

★ نقطة واحدة

**The imbalance in Ischemic Heart Disease refers to a mismatch -2
:between**

- A. Oxygen supply and carbon dioxide removal in the lungs ☐
- B. Myocardial oxygen supply and myocardial oxygen demand ☒
- C. Blood glucose supply and myocardial energy demand ☐
- D. Atrial and ventricular contraction rates ☐

★ نقطة واحدة

:Organic nitrates, like nitroglycerin, primarily alleviate angina by -3

- A. Increasing myocardial contractility ☐
- B. Causing direct coronary artery vasoconstriction ☐
- C. Reducing myocardial oxygen demand through venodilation (reducing preload) and some arteriodilation (reducing afterload) ☒
- D. Blocking beta-adrenergic receptors ☐

★ نقطة واحدة

**Which of the following beta-blockers is cardioselective (beta-1 -4
?selective)**

- A. Propranolol ☐
- B. Metoprolol ☒
- C. Nadolol ☐
- D. Timolol ☐

نقطة واحدة * In patients with stable IHD and a history of myocardial infarction -5 or heart failure, which class of drugs is generally recommended ?indefinitely IF not contraindicated to reduce mortality

- A. Diuretics ☐
- B. Beta-blockers (evidence-based) ☒
- C. Dihydropyridine calcium channel blockers as monotherapy ☐
- D. Ranolazine ☐

نقطة واحدة * A crescendo-decrescendo systolic ejection murmur, radiating to -6 :the carotids, is a hallmark of

- A. Mitral stenosis ☐
- B. Aortic regurgitation ☐
- C. Mitral valve prolapse ☐
- D. Aortic stenosis ☒

نقطة واحدة * Pulsus parvus et tardus' (a weak and delayed carotid pulse) is a -7 :clinical sign associated with severe

- A. Mitral regurgitation ☐
- B. Aortic regurgitation ☐
- C. Aortic stenosis ☒
- D. Tricuspid stenosis ☐

نقطة واحدة * A high-pitched, blowing, early diastolic decrescendo murmur -8 :best heard at the left sternal border is typical of

- A. Aortic regurgitation ☒
- B. Pulmonary stenosis ☐
- C. Tricuspid regurgitation ☐
- D. Mitral stenosis ☐

* نقطة واحدة

**Corrigan's pulse (water-hammer pulse) and a wide pulse -9
:pressure are classic findings in**

- A. Mitral stenosis ☐
- B. Severe aortic regurgitation ☒
- C. Severe aortic stenosis ☐
- D. Cardiac tamponade ☐

* نقطة واحدة

**A 'P mitrale' (broad, notched P wave in lead II) on an ECG is -10
:most indicative of**

- A. Right atrial enlargement ☐
- B. Left ventricular hypertrophy ☐
- C. Left atrial enlargement ☒
- D. Right ventricular hypertrophy ☐