

45. Secondary hypertension is caused commonly by:

A. Necrotizing arteriolitis

B. Hypothyroidism

C. Aortic stenosis

D. Renal ischemia

46. Not true regarding gross picture of the kidney in benign essential hypertension:

A. Kidney is Decreased size

B. Capsule is adherent

C. Smooth surface with areas of hemorrhage

D. On cut section don't differentiate between cortex and medulla

47. In malignant hypertension, the arterioles DO NOT SHOW:

A. Fibrinoid necrosis

B. Thickening of the wall

C. Elastosis

D. Narrowing of the lumen

Thick
Fibrinoid necrosis

Narrow.

25- Gradual incomplete occlusion of the coronary artery Not complicated by:

- A-Angina Pectoris ✓
- B-Chronic heart failure ✓
- C-Myocardial infarction ✓
- D-Cardiac arrhythmias ✓

26- Which of the following is NOT a cause of myocarditis?

- A-Tuberculosis.
- B-Diphtheria.
- C-Atherosclerosis. ✓
- D-Rheumatic fever.

Atherosclerosis

VALVULAR DISEASES AND HEART FAILURE

27- Not considered as a cause of right ventricular hypertrophy.

MYOCARDIAL DISEASES

19- The main sites of myocardial infarction are:

- ☒ A- Left ventricle
- ☐ B- Left atrium
- ☐ C- Right ventricle
- ☐ D- Right atrium

20- Autopsy of a patient who died from myocardial infarction often reveals coronary artery obstruction by:

- ☐ A- Non-complicated atheroma
- ☒ B- A thrombus on top of atheroma
- ☐ C- Thrombotic embolus
- ☐ D- Stenosis at the coronary ostia

21- The catastrophic form of ischemic heart disease is:

- ☐ A- Angina pectoris
- ☐ B- Ischemic cardiomyopathy
- ☒ C- Myocardial infarction
- ☐ D- Toxic myocarditis

MI

22- Common cause of death immediately after the onset of acute myocardial infarction is:

- ☒ A- Arrhythmias
- ☐ B- Left ventricular rupture
- ☐ C- Congestive Heart failure
- ☐ D- Shock

Arrhythmia

23- Presence of intermittent chest pain caused by reversible myocardial ischemia is characteristic finding in:

- ☐ A- Myocardial infarction
- ☒ B- Angina Pectoris
- ☐ C- Pericarditis
- ☐ D- Myocarditis

Angina Pectoris

24- A 53-year-old man is found to have a hemorrhagic pericarditis (cardiac tamponade).

This should make you most strongly suggest:

- ☐ A- Autoimmune disease
- ☐ B- Rheumatic fever
- ☐ C- Malignant hypertension
- ☒ D- Acute myocardial infarction

36- Which one of the following is the most common congenital heart defect to cause an initial left-to-right shunt?

- A- Tetralogy of Fallot. ✓
- B- Coarctation of the aorta. ✗
- C- Ventricular septal defect. ✓
- D- Atrial septal defect. ✗

ventricular septal defect

37- False regarding Fallot's triology

- A- Atrial septal defect ✓
- B- Pulmonary stenosis ✓
- C- Mitral stenosis ✗
- D- Right ventricular Hypertrophy ✓

Mitral

38- The most common primary tumor of the heart in adults is usually located in the left atrium and is called a ?

- A- Chordoma
- B- Rhabdomyoma
- C- Leiomyoma
- D- Myxoma

Myxoma

CONGENITAL HEART DISEASES & CARDIAC TUMORS

35- The combination of a ventricular septal defect with an aorta that overrides the septal defect, stenosis of the pulmonic valve, and increased thickness of the right ventricle is diagnostic of?

A- Tetralogy of Fallot

B- Eisenmenger's syndrome.

C- Löffler's syndrome.

D- Chagas' disease.

Tetralogy of Fallot

13- The characteristic histologic feature of rheumatic fever is:

A-Rheumatic nodule.

B-Aschoff nodule ✓

C-MacCallum's patch

D-Peyer's patch

Aschoff nodule

14- MacCallum's patch is feature of:

A-Rheumatic pericarditis

B-Rheumatic myocarditis

C-Rheumatic endocarditis ✓

D-Rheumatic valvulitis

✓ mural endocarditis

15- Not considered as complication of Subacute infective endocarditis:

A-Clubbing of the fingers

B-Septic emboli cause pyemia. ✓

C-Osler's nodes

D-Mycotic aneurysm.

✓ Septic

★ 16- Vegetations in acute infective endocarditis Not characterized by?

A-Single ✓

B-Globular

C-Large

D-Firmly adherent to the valve ✓

17- Libman-Sacks endocarditis occurs in:

A-Rheumatic fever ✓

B-Rheumatoid fever ✓

C-Systemic lupus erythematosus ✓

D-Systemic sclerosis ✓

SLE

18- The following is not a feature of rheumatic vegetation:

A-Multiple ✓

B-Beaded. ✓

C-Easily detached ✓

D-Pin-head size.

Firmly attached (adherent)

- major
- Arthritis
 - Erythema
 - Carditis
 - Subcutaneous nodule
 - Chorea

- minor
- Fever
 - ↑ ESR
 - ↑ CRP
 - Arthralgia
 - Glomerulonephritis

7- Not considered as major manifestation (Jones criteria) for acute rheumatic fever?

A-Carditis

B-Glomerulonephritis

C-Migratory polyarthritis

D-Sydenham's chorea

GN

8- Lesions due to Rheumatic fever in the basal ganglion of brain causes the following?

A-Meningitis

B-Encephalitis

C-Chorea

D-Degeneration of brain

Chorea

MCQ

9- Not considered as modified Jones criteria for Diagnosis of Rheumatic Fever?

A-Presence of two major criteria

B-One major & two minor criteria

C-Two major & one minor criteria

D-Only one minor criteria

10- The following type of vegetations never give emboli:

Rheumatic v

A-Rheumatic vegetations

B-Vegetation in acute infective endocarditis

C-Vegetations in subacute infective endocarditis

D-Vegetations in non-bacterial endocarditis

11- Bread and butter appearance is characteristic of:

A-Rheumatic pericarditis

Pericarditis

B-Rheumatic myocarditis

C-Rheumatic endocarditis

D-Rheumatic valvulitis

12- Not included in structure of Aschoff nodule:

A-Epithelioid cell

B-Macrophages

C-Lymphocytes

D-Giant cell

Epithelioid cells

HP
IS
GK

31- Paroxysmal nocturnal dyspnea is a manifestation of:

A-Right-heart failure

✓ B-Left sided heart failure

C-Bronchial asthma

D- Angina pectoris.

left s. see heart failure

32- A ball thrombus is found inside the heart in a case of?

A-Aortic stenosis

B-Aortic incompetence

✓ C-Mitral stenosis

D-Pulmonary stenosis

mitral stenosis

33- A 45 -years- old woman has had worsening shortness of breath for 3 years. She now has to sleep sitting up on two pillows. She has no history of chest pain. A chest radiograph reveals a near-normal left ventricular size with a prominent left atrial border. Which of the following conditions is most likely to account for these findings?

A- Essential hypertension

B- Cardiomyopathy

✓ C-Mitral valve stenosis

✓ D- Aortic coarctation

mitral stenosis

34- A 42-year-old woman has increasing congestive heart failure. As a child she suffered recurrent bouts of pharyngitis with group A beta hemolytic streptococcal infections. The cardiac valves most likely to be affected are:

A-Mitral and pulmonic

B-Aortic and pulmonic

C-Tricuspid and pulmonic

✓ D-Mitral and aortic

mitral + aortic

left side

CONGENITAL HEART DISEASES & CARDIAC TUMORS

35- The combination of a ventricular septal defect with an aorta that overrides the septal defect, stenosis of the pulmonic valve, and increased thickness of the right ventricle is diagnostic of?

✓ A-Tetralogy of Fallot

B- Eisenmenger's syndrome.

C-Löffler's syndrome.

D-Chagas' disease

tetralogy of Fallot

MCQs

RHEUMATIC FEVER & ENDOCARDITIS

1- Commonest Age group involved in Rheumatic Fever is:

- A-Old age
- B-Young age from 21-30 years.
- C-Childhood 5-15 yrs.
- D-Above 60 Years of age.

5-15 ✓

2- The type of inflammation in rheumatic pericarditis is:

- A-Catarrhal.
- B-Fibrinous.
- C-Membranous.
- D-Suppurative.

Serofibrinous ✓

3- Acute rheumatic fever classically develops in children 1 to 4 weeks after a group A β -hemolytic streptococcal (*Streptococcus pyogenes*) infection of the?

- A-Aorta
- B-Heart
- C-Lungs
- D-Pharynx

Pharynx

4- Which of the following cells not included in Aschoff body?

- A-Langhans giant cells.
- B-Antischkow myocytes.
- C-Lymphocytes
- D-Aschoff giant cells

1- Antischkow Myocytes.
2- Aschoff giant cells
3- Lymphocytes.

5- MacCallum's patch is usually located in the:

- A-Pericardium.
- B-Posterior wall of right atrium
- C-Interventricular septum.
- D-Posterior wall of left atrium.

→ mural endocardium

6- Type of necrosis accompanied with Aschoff nodule is called:

- A-Coagulative necrosis
- B-Liquefactive necrosis
- C-Fibrinoid necrosis
- D-Fat necrosis

Fibrinoid necrosis