

0/1*

A nurse is taking .2 ✕
pulmonary artery catheter
measurements of a male
client with acute respiratory
distress syndrome. The
pulmonary capillary wedge
pressure reading is 18mm
Hg. The nurse interprets that
this readings is



High and expected



Low and unexpected



Normal and expected



Uncertain and unexpected



الإجابة الصحيحة

Normal and expected



1/1*

Which of the following .3 ✓
drug classes are not
considered primary therapy
of asthma and are more
often used as alternatives

1/1* Which of the following .3 ✓
 drug classes are not
 considered primary therapy
 of asthma and are more
 often used as alternatives
 when control of the asthma
 cannot be achieved by first-
 ?line regimens

SABA. ☐

Leukotriene modifiers ☐

Theophylline/methylxanthines. ☐

✓ ICSs and Leukotriene ☒
 modifiers

1/1* Regarding to the drug .4 ✓
 interaction of theophylline.
 The half-life of theophylline
 is decrease by which of the
 ?following

Erythromycin ☐

✓ Phenobarbitone ☒



Leukotriene modifiers ☐

Theophylline/methylxanthines. ☐

☒ ICSs and Leukotriene modifiers ☒

1/1* Regarding to the drug .4 ☒
 interaction of theophylline.
 The half-life of theophylline
 is decrease by which of the
 ?following

Erythromycin ☐

☒ Phenobarbitone ☒

Quinolones ☐

Cimetidine ☐

1/1* A 30-year-old patient .5 ☒
 with a history of mild
 persistent asthma (baseline
 peak expiratory flow rate of
 85%) presents to the
 emergency department with
 shortness of breath and

1/1*

A 30-year-old patient .5 ✓
 with a history of mild
 persistent asthma (baseline
 peak expiratory flow rate of
 85%) presents to the
 emergency department with
 shortness of breath and
 wheezing that has not
 relieved by her albuterol
 inhaler for the past 12 hours.
 She was able to tolerate
 pulmonary function tests
 and a set was performed.
 Which of the following is the
 ?most likely test result

✓ Decreased FEV1,
 normal/increased FVC,
 decreased FEV1: FVC ratio,
 with post bronchodilator. ☒

FEV1 increased by 13%. ☐

Decreased residual volume and
total lung capacity ☐

Increased FEV1 increased FVC,
normal FEV1: FVC ratio ☐

1/1*

Which one of the following, with Formoterol, is a long acting anti-asthmatic ?drugs



Albuterol ☐



Salmeterol ☒

Terbutaline ☐

Salbutamol ☐

1/1*

Which of the following is not a common symptom of ?mesothelioma of pleura



Chest pain. ☐

Dyspnea. ☐

Weight loss ☐



Diarrhea. ☒

0/1

Which one of the following statements is true



0/1

Which one of the following statements is true of mesothelioma .8 X

Incidence is decreasing, reflecting the ban on use of asbestos ☐

The disease is more common in females. ☐

All deaths due to mesothelioma should be discussed with the coroner. ☐

X Radical surgery is the treatment of choice ☒

أخرى: ☐

الإجابة الصحيحة

All deaths due to mesothelioma should be discussed with the coroner. ☒

0/1*

MRI in patient with mediastinal lesion used for diagnosis of .9 X



0/1*

MRI in patient with .9 ✕
mediastinal lesion used for
diagnosis of



Vascular lesion ☒

Cardiac lesion ☐

Esophageal lesion ☐

Tracheal lesion ☐

الإجابة الصحيحة

Esophageal lesion ☒

1/1*

The most common .10 ✓
clinical picture of
mediastinal lesion is



Asymptomatic ☒

Hornor syndrome ☐

Horsness of voice ☐

dysphagia ☐

1/1* The right bronchial .11 ✓
 ._____ vein drain into



Azygos vein ☒

Hemi-azygos vein ☐

Axillary vein ☐

Left superior intercostal vein ☐

0/1* Which of the following .12 ✗
 is a nonspecific finding
 common to interstitial lung
 ?disease

Raised LDH. ☐

Antinuclear antibodies. ☐

Anti-immunoglobulin antibodies
 (rheumatoid factors). ☐

✗ Circulating immune
 complexes. ☒

الإجابة الصحيحة

Raised LDH. ☒



1/1* The most common .13 ✓
form of DPLD is

- ✓ Occupational lung disease ☒
- Sarcoidosis ☐
- Idiopathic interstitial pneumonitis ☐
- Drug induced ☐

0/1* Which of the following .14 ✗
?is not a fibrosing disease

- Anthracosis. ☐
- ✗ Asbestosis ☒
- Pneumoconiosis ☐
- Hypersensitivity pneumonitis ☐

الإجابة الصحيحة

Pneumoconiosis ☒

0/1* Which of the following .15 ✗



Hypersensitivity pneumonitis ☐

الإجابة الصحيحة

Pneumoconiosis ☒

0/1* Which of the following .15 X
chest x-ray findings is
indicative of a poor
prognosis in a case of
?interstitial lung disease

X Bibasilar reticular pattern ☒

Honeycombing ☐

Nodular pattern of alveolar filling ☐

Mixed pattern of alveolar filling ☐

الإجابة الصحيحة

Honeycombing ☒

0/1 A 66 years old lady .16 X
presented to ER with sudden
acute chest pain and
shortness of breath. Her



0/1

A 66 years old lady .16 
presented to ER with sudden
acute chest pain and
shortness of breath. Her
blood pressure was 45\30
mmHg, pulse 132, respiratory
rate 35, SpO2 was 81%. There
was redness and hotness in
left lower limb. In this case
the heparin therapy for
pulmonary embolism must be
:started

On clinical suspicion. ☐

 After confirmation of
diagnosis ☒

After confirmation of RV
dysfunction. ☐

After confirmation of DVT. ☐

الإجابة الصحيحة

On clinical suspicion. ☒

0/1

Acute sub massive .17 

On clinical suspicion. ☒

0/1 Acute sub massive .17 ✗
pulmonary embolism is
defined as

Cardiac arrest ☐

Normal systolic blood pressure
with right ventricular strain ☐

✗ Systolic blood pressure less
than 100mmHg ☒

Hypotension with no right
ventricular strain ☐

Normal systolic blood pressure
with right ventricular strain ☒

1/1* Which of the following .18 ✓
echocardiographic findings
is not a feature of acute
massive pulmonary
?embolism

ventricular strain

الإجابة الصحيحة

Normal systolic blood pressure
with right ventricular strain



1/1* Which of the following .18 ✓
echocardiographic findings
is not a feature of acute
massive pulmonary
?embolism

RV/LV ratio >1. ☐

Can detect thrombus floating in
the right side of the heart before
reaching lungs. ☐

✓ Identification of LV
dysfunction ☒

Identification of RV dysfunction ☐

0/1* A 70-year-old, female, .19 ✗
his husband is a constructor
presented with dyspnea and
chest pain. Chest X-ray
showed small RT sided

0/1* A 70-year-old, female, 19 X
 his husband is a constructor
 presented with dyspnea and
 chest pain. Chest X-ray
 showed small RT sided
 pleural effusion and pleural
 nodules. Aspiration of
 pleural fluid revealed
 exudative hemorrhagic
 pleural effusion. Which of
 the following is the most
 ?diagnostic next step

Transthoracic ultrasound guided
 pleural biopsy ☐

X Thoracoscopic guided pleural
 biopsy ☒

Abrams pleural biopsy ☐

CT guided pleural biopsy ☐

الإجابة الصحيحة

Transthoracic ultrasound guided
 pleural biopsy ☒

pleural biopsy

✗ Thoracoscopic guided pleural biopsy



Abrams pleural biopsy



CT guided pleural biopsy



الإجابة الصحيحة

Transthoracic ultrasound guided pleural biopsy



1/1

The mediastinum .20 ✓
:demarcated anterior by

Heart



Manubrium sterni



Sternum



Right lung



لم يتم إنشاء هذا المحتوى ولا اعتماده من قبل Google. الإبلاغ عن
إساءة الاستخدام - شروط الخدمة - سياسة الخصوصية

Google نماذج



1/1* A study of persons with a .1 ✓
history of mining
occupational exposure to
inhaled dusts is performed.
Though found in urban air in
small amounts, this dust
consists of 1 to 5micron
particles that are inert and
insoluble. Fibrosis occurs
only with large amounts of
dust accumulation, mainly in
upper lobes, with nodular
opacities larger than 1 cm
seen on chest
radiographs. What is most li
?kely in this dust

Iron ☐



coal dust ☒

Carbon ☐

Sulfur dioxide ☐

أخرى: ☐

1/1

**Neurological and .2 ✓
behavioral findings in
patients with hypoxia
:included**

Headache ☐



Drowsiness ☒

Tremor and asterix. ☐

Verbal language loss ☐

أخرى: ☐

1/1

**Glasgow coma scale is .3 ✓
:used to assess**

Behavioral functions. ☐

Motor functions. ☐

Sensory functions. ☐



Conscious level. ☒

أخرى: ☐

Conscious level. ☒أخرى: ☐

1/1 Which of the following .4 ✓
causes of respiratory failure
is recognized to have the
?highest mortality rate

Chronic obstructive pulmonary
disease (COPD) ☐

Asthma ☐

✓ Acute respiratory distress
syndrome (ARDS) ☒

Obesity ☐

أخرى: ☐

1/... Which of the following is .5 ✗
a common cardiovascular
complication in patients with
?acute respiratory failure

A Hypertension

✓ Acute respiratory distress syndrome (ARDS) ☒

Obesity ☐

أخرى: ☐

1/... Which of the following is .5 ✗
a common cardiovascular
complication in patients with
?acute respiratory failure

A. Hypertension

B. Endocarditis

C. Aortic coarctation

D. Coronary artery spasm

✗

Option 1 ☒

ليس هناك أي إجابات صحيحة

1/1 Which of the following .6 ✓
is *not* considered a routine
investigation in patients with
?suspected respiratory failure

ليس هناك أي إجابات صحيحة

1/1 Which of the following .6 ✓
is *not* considered a routine
investigation in patients with
?suspected respiratory failure

Chest x-ray ☐

Arterial blood gas analysis ☐

✓ Right-sided heart catheterization ☒

Electrocardiography ☐

أخرى: ☐

0/1 Which one of the .7 ✗
following is not true regarding
?bronchogenic carcinomas

✗ Radiation Therapy is effective when the lung cancers are confined to the chest but cannot be removed surgically. ☒

For non-small cell lung cancer

0/1

Which one of the following is not true regarding ?bronchogenic carcinomas

☒ Radiation Therapy is effective when the lung cancers are confined to the chest but cannot be removed surgically.

For non-small cell lung cancer, TNM descriptions lead to a simpler categorization of stages.

Small cell lung cancer has two stages: limited or extensive stages.

Smoking is contraindication for surgery in bronchogenic carcinomas

أخرى:

الإجابة الصحيحة

Smoking is contraindication for surgery in bronchogenic carcinomas

1/1

Which one of the following is true regarding computed tomography for chest scans with intravenous injection of contrast (CT chest) for radiological assessment of lung cancer ✓

CT may be performed on the chest, abdomen, and/or brain for diagnosis of both primary and metastatic lung tumors. ☐

CT may be performed for nodule details as calcification, spiculation etc. ☐

✓ study a particular area as: Lung tumor based on the diaphragm or the uppermost part of the lung ☒

Evaluate extension into adjacent structures: i.e. endobronchial, great vessels, pericardium etc. ☐

أخرى: ☐

1/1

Which one of the .9 ✓

1/1

Which one of the following is true regarding computed tomography for chest scans with intravenous injection of contrast (CT chest) for radiological assessment of lung cancer ✓

CT is superior to MRI Chest for radiological assessment of suspected lung cancer. ☐

CT is often used in the routine work-up of suspected lung cancer. ☐

✓ CT chest with contrast is non-efficient for evaluation of adenopathy. ☒

Upper abdominal pathology as metastatic lesions in liver, adrenals, & kidneys. ☐

أخرى: ☐

1/1

Which one of the following is true regarding magnetic resonance imaging (MRI) for radiological ✓

1/1

Which one of the following is true regarding magnetic resonance imaging (MRI) for radiological assessment of ?bronchogenic carcinomas ✓

MRI is superior to CT Chest for radiological assessment of suspected lung cancer. ☐

MRI is often used in the routine work-up of suspected lung cancer. ☐

✓ MRI may be used to study a particular area that may be difficult to be shown on CT scan such as: Lung tumor based on the diaphragm or the uppermost part of the lung ☒

Evaluation of adenopathy ☐

أخرى: ☐

1/1

Which one of the following is a first line drug ?for TB treatment ✓

1/1

Which one of the following is a first line drug for TB treatment ✓



Pyrazinamide ☒

PASA ☐

Kanamycin ☐

Cycloserine ☐

أخرى: ☐

1/1

The diameter of indurations is >15mm for ✓

Children less than 4 years old ☐



Normal individuals ☒

HIV or high risk for HIV ☐

Microbiology lab workers ☐

Option 4 ☐

أخرى: ☐

1/1

The main virulence factor in TB pathogenesis is .13 ✓

Bacteria's ability to survive inside the bronchi ☐

Bacteria's ability to destroy the macrophage ☐

Bacteria's ability to destroy the pulmonary tissue ☐

✓ Bacteria's ability to survive inside the macrophage ☒

أخرى: ☐

1/1

What is the most common cause of simple parapneumonic exudative ?pleural effusion .14 ✓

Viral infection ☐

Fungal infection. ☐

✓ Bacterial infection. ☒

Tb infection ☐

أخرى: ☐

1/1

**What is the most .15 ✓
diagnostic feature in case of
?urinorhox**

Effusion ipsilateral to obstructed
kidney with retroperitoneal urine
leak. ☐

Resolves after treatment of
obstruction ☐

pH usually low. ☐

✓ Pleural fluid creatinine > serum
creatinine is diagnostic ☒

أخرى: ☐

**1/1 The most common site of .16 ✓
origin of malignant
:Mesothelioma is**

Pericardium. ☐

✓ Pleura ☒

Peritoneum. ☐

Lung. ☐

أخرى: ☐

1/1 **Which of the following is true regarding pulmonary circulation?** ✓

✓ It is a low-pressure system that allows for improved gas exchange. ☒

It is the larger of the two circulatory systems. ☐

The system functions with an increased arterial pressure to circulate through the distal parts of the body. ☐

It consists of the left side of the heart, the aorta, and its branches. ☐

أخرى: ☐

1/1 **A 45-year-old, female, 18 presented with dyspnea and dry cough. Chest X-ray showed left sided cavity with air-fluid level, patient started broad spectrum antibiotics with no response for 10 days.** ✓

أخرى: ☐

1/1 **A 45-year-old, female, 18** ✓
presented with dyspnea and
dry cough. Chest X-ray
showed left sided cavity with
air-fluid level, patient started
broad spectrum antibiotics
with no response for 10 days.
?What is next step

✓ Bronchoscope for
microbiological analysis of
bronchoalveolar lavage. ☒

Referral to cardiothoracic surgeon. ☐

Thoracoscopic biopsy. ☐

Transthoracic percutaneous
needle aspiration (CT- or US-
guided). ☐

أخرى: ☐

1/1 **Concerning lung abscess, 19** ✓
?which of the following is true



أخرى: ☐

1/1 **Concerning lung abscess, .19 ✓**
?which of the following is true

✓ Lung abscess is a type of
liquefactive necrosis of the
lung tissue and formation of
cavities. ☒

Lung abscess is a type of
liquefactive necrosis of the lung
tissue without formation of
cavities. ☐

Lung abscess is a type of fibrosis
of the lung tissue. ☐

Lung abscess is a type of
inflammation of the lung tissue. ☐

أخرى: ☐

1/1 **the cause of localized .20 ✓**
bronchiectasis is

Recurrent chest infection ☐

cavities.

Lung abscess is a type of fibrosis of the lung tissue.



Lung abscess is a type of inflammation of the lung tissue.



أخرى:



1/1

20. the cause of localized bronchiectasis is



Recurrent chest infection



Foreign body aspiration



Cystic fibrosis



Allergic bronchopulmonary aspergillosis



أخرى:



لم يتم إنشاء هذا المحتوى ولا اعتماده من قبل Google. الإبلاغ عن
إساءة الاستخدام - شروط الخدمة - سياسة الخصوصية

