microCT imaging of threespine stickleback

This manuscript (permalink) was automatically generated from habi/sticklebacks-manuscript@6016dd9 on February 25, 2026.

Authors

- **David Haberthür**
  0000-0003-3388-9187 · habi · @habi@mastodon.social
  microCT research group, Institute of Anatomy, University of Bern, Baltzerstrasse 2, 3012 Bern, Switzerland

- **Ben Sulser**
  0000-0002-8750-0942 · sulserrb
  Evolutionary Ecology Group, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland · Funded by Bern Burgergemeinde

- **Sheila Christen**

- **Catherine L. Peichel**
  0000-0002-7731-8944 · cpeichel
  Division of Evolutionary Ecology, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland · Funded by Swiss National Science Foundation (TMAG-3_209309/1)

- **Ruslan Hlushchuk** ✉
  0000-0001-2345-6789 · RuslanHlushchuk
  microCT research group, Institute of Anatomy, University of Bern, Baltzerstrasse 2, 3012 Bern, Switzerland

✉ — Correspondence possible via GitHub Issues or email to Ruslan Hlushchuk <ruslan.hlushchuk@unibe.ch>.

Abstract

Can we predict evolution? The three-spined stickleback (Gasterosteus aculeatus) is a well-recognized system for understanding adaptation to divergent habitats. Populations of benthic and limnetic stickleback differ in a number of phenotypic traits that are associated with shifts in dietary specialization. However, analyses of the structures required for feeding – especially the jaws and complex internal branchial anatomy – requires considerable time and expertise, with destructive sampling and fine dissection skills needed for quantitative analysis. The advent of μCT and 3D-scanning technology affords non-destructive sampling and an increase the resolution of data available for study, but at the substantial cost of increasing complexity and processing time for each specimen. To address these concerns, we developed a rapid and semi-automated segmentation and analysis pipeline based around the Biomedisa Image Segmentation platform for investigating three-dimensional morphological adaptation within the three-spined stickleback. The pipeline includes splitting a multi-specimen scan into regions of interest for each specimen, reconstruction of targeted anatomy, and morphometric analyses. We then applied this pipeline to a sampling effort encompassing hundreds of samples of divergent benthic and limnetic stickleback populations, showcasing the possibility of using high-throughput scanning data to provide tests of ecological and evolutionary hypotheses.

Introduction

- Embedded into Alaska Stickleback Restoration Project, Genomics axis where Katie Peichel, Ben Sulser and Sheila Christen are affiliated.
- Why are we studying what we are studying in these fish?

Micro-computed tomography

X-ray microtomography is an indispensable tool to gain non-destructive insights into the inner structure of highly diverse samples, namely for specimens studied in the biomedical sciences [1]. Microtomographic imaging is ideally suited to nondestructively assess the morphology of different fish species, large and small [2].

Materials & Methods

Sample procurement and preparation

Due to their inherent contrast difference to the surrounding tissue, the structures of interest we touch upon in this manuscript (teeth and bones, i.e. jaws and skull) are well visualized in unstained samples, hence no further preparation of the fish was necessary.

microtomographic imaging

In a small pilot study we determined the optimal scanning parameters to fit the constraints on total scanning time, resolution and sample handling. To optimize for these constraints, we scanned all the sticklebacks in batches of six fish in a custom-made 3D printed sample holder in a single scan. This holder was generated with OpenSCAD and is available online, either directly as STL file for printing or as (parametrized) OpenSCAD file for adaptation to other classes of samples. Both files are part of a library of 3D-printable sample holders for tomographic imaging [3].

Tomographic imaging was performed on a Bruker SkyScan 2214 at the Institute of Anatomy, University of Bern, Switzerland. In total we performed 44 scans, each of the scan usually containing 6 fish in the sample holder.

The relevant details of each scan are collated in a table in the Supplementary Materials; a short overview of the scanning parameters is given below. The X-ray source was set to a voltage of 60 kV and a current of around 110 µA for all but one scan where we used a source voltage of 49 kV and 159 µA due to operator error. For each sample, we recorded a set of 3601 projections of approximately 3000 x 2000 pixels at every 0.1° over a 360° sample rotation. Every single projection was exposed for about a second (depending on the sample). Due to the lenght of the fish, we had to acquire so-called stacked scans, on average we scanned 3 fields of view along the rotation axis of the sample holder. This resulted in scan times between 3 to 5 hours hours. The projection images were then subsequently reconstructed into a 3D stack of images with NRecon (Bruker microCT, Kontich Belgium, Version: 2.1.0.1 or 2.2.0.6). The isometric voxel sized in the resulting datasets vary from 15 to 19 µm.

Data analysis

Preparation and handling of tomographic datasets

After acquisition, a simple script was used to copy the relevant data to both archival storage and storage accessible by all co-authors at the same time.

Further processing of the tomographic dataset was performed with a set of Jupyter [4] notebooks [5].

- Efficiently loading data from disk with `dask` [6]
- Extract position of single fish (all scanned together), based on the MIP of the scan (see Figure 2).
- Crop out each fish (with a buffer) and write to cropped dataset (see Figure 4).
- Cropped datasets are saved to discrete folders for easy handling. In both original gray-scale plus as thresholded dataset, e.g. binarized into bone and “not bone”. These are saved out as `zarr` [7] and `nrrd` files.



Figure 1: Maximum intensity projections of one acquired dataset along all three cardinal axes.



Figure 2: Automatically detected regions based on maximum intensity projection along the rotation axis of the tomographic scan. The regions are numbered consecutively from the top left to the bottom right. These numbers are mapped to the correct fish ID in the next step.



Figure 3: Mapping lab book notes, photos and detected regions to fish ID.



Figure 4: Doublechecking crop extent and fish ID.

Extraction of features of interest

- Biomedisa [8]

Results

- 3D shape variation on internal pharyngobranchial bone. Only possible to get this information in 3D.

Discussion

- Both *repeatable* and *reproducible* research
- Automated cropping out of single fish from combined scan very efficiently uses machine time. Several fish can be scanned together, splitting is performed after the fact, reproducibly and without manual input.
- Combination of methods cuts down on time **a lot**.
- Biomedisa makes more “extraction” possible. Other biological questions can be answered, too.

Conclusion

Author Contributions

Contributor Roles Taxonomy (CRediT), as defined in [9]:

- Conceptualization: Ben Sulser, Ruslan Hlushchuk
- Data curation: David Haberthür, Ben Sulser
- Formal analysis: David Haberthür, Ben Sulser
- Funding acquisition: Ben Sulser, Catherine L. Peichel, Ruslan Hlushchuk
- Investigation: David Haberthür, Ben Sulser
- Methodology: David Haberthür, Ben Sulser, Ruslan Hlushchuk
- Project administration: David Haberthür, Ben Sulser, Catherine L. Peichel, Ruslan Hlushchuk
- Resources: Ben Sulser, Ruslan Hlushchuk
- Software: David Haberthür, Ben Sulser
- Supervision: Ben Sulser, Catherine L. Peichel, Ruslan Hlushchuk
- Validation: David Haberthür, Ben Sulser
- Visualization: David Haberthür, Ben Sulser
- Writing – original draft: David Haberthür, Ben Sulser
- Writing – review & editing: David Haberthür, Ben Sulser, Catherine L. Peichel, Ruslan Hlushchuk

Competing Interest

Author	Competing Interests	Last Reviewed
David Haberthür	None	2026-01-14
Ben Sulser	Nothing to Declare	
Sheila Christen		
Catherine L. Peichel	none	19.01.2026
Ruslan Hlushchuk	None	2026-01-19

Acknowledgments

We are grateful to the Microscopy Imaging Center of the University of Bern for their infrastructural support. We also thank the `manubot` project [10] for facilitating collaborative writing of this manuscript.

Supplementary Materials

Parameters of tomographic scans of all the fishes

The CSV file ScanningDetails.csv gives a tabular overview of all the (relevant) parameters of all the scans we performed. This file was generated with the data processing notebook and collates the relevant data read from *all* the log files of *all* the scans we performed. A copy of each log file containing *all* scanning parameters is available in a folder in the data processing repository.

References

1. **X-ray computed tomography in life sciences**
Shelley D Rawson, Jekaterina Maksimcuka, Philip J Withers, Sarah H Cartmell
BMC Biology (2020-02-27) https://doi.org/ggrx9x
DOI: 10.1186/s12915-020-0753-2 · PMID: 32103752 · PMCID: PMC7045626

2. **A New Era of Morphological Investigations: Reviewing Methods for Comparative Anatomical Studies**
KL Ford, JS Albert, AP Summers, BP Hedrick, ER Schachner, AS Jones, K Evans, P Chakrabarty
Integrative Organismal Biology (2023) https://doi.org/grx5rb
DOI: 10.1093/iob/obad008 · PMID: 37035037 · PMCID: PMC10081917

3. **TomoGraphics/Hol3Drs: A release**
David Haberthür
Zenodo (2019-03-08) https://doi.org/gg9fxh
DOI: 10.5281/zenodo.2587555

4. **Jupyter Notebooks – a publishing format for reproducible computational workflows**
Kluyver Thomas, Ragan-Kelley Benjamin, Pérez Fernando, Granger Brian, Bussonnier Matthias, Frederic Jonathan, Kelley Kyle, Hamrick Jessica, Grout Jason, Corlay Sylvain, … Jupyter Development Team
Positioning and Power in Academic Publishing: Players, Agents and Agendas (2016) https://doi.org/gf48c9
DOI: 10.3233/978-1-61499-649-1-87

5. **habi/sticklebacks: Maintenance release**
David Haberthür
Zenodo (2026-01-15) https://doi.org/hbj4g5
DOI: 10.5281/zenodo.18257528

6. **Dask: Library for dynamic task scheduling**
Dask Development Team
(2016) https://dask.org

7. **Zarr** https://www.wikidata.org/wiki/Q130377195

8. **Introducing Biomedisa as an open-source online platform for biomedical image segmentation**
Philipp D Lösel, Thomas van de Kamp, Alejandra Jayme, Alexey Ershov, Tomáš Faragó, Olaf Pichler, Nicholas Tan Jerome, Narendar Aadepu, Sabine Bremer, Suren A Chilingaryan, … Vincent Heuveline
Nature Communications (2020-11-04) https://doi.org/gh58qn
DOI: 10.1038/s41467-020-19303-w · PMID: 33149150 · PMCID: PMC7642381

9. **ANSI/NISO Z39.104-2022, CRediT, Contributor Roles Taxonomy** *NISO* https://doi.org/gqx265
DOI: 10.3789/ansi.niso.z39.104-2022

10. **Open collaborative writing with Manubot**
Daniel S Himmelstein, Vincent Rubinetti, David R Slochower, Dongbo Hu, Venkat S Malladi, Casey S Greene, Anthony Gitter
PLOS Computational Biology (2019-06-24) https://doi.org/c7np
DOI: 10.1371/journal.pcbi.1007128 · PMID: 31233491 · PMCID: PMC6611653