

Advanced Machine learning Mastering Course

Introduced by

George Samuel

Master in computer science
Cairo University

Innovisionray.com

2024



Agenda

1 Supervised Learning

2 Introduction to Regression

3 More Regression

4 Regression in Sklearn

5 Gradient decent-Normal equation

6 Regularization-Logistic Regression

7 Logistic Regression

8 Decision Tree

10 Neural Networks

11 Neural Nets Mini-Project

11 Math behind SVMs

11 SVMs in Practice

12 Instance Based Learning

13 Regression in Sklearn

14 Naive Bayes

15 Bayesian Inference

16 Ensemble B&B

17 Finding donors for CharityML

Three Types of machine learning models

ML Models

Geometric models

Linear Regression

Logical models

Decision Tree

Probabilistic Models

Logistic Regression

CLASSIFICATION

The classification problem is just like the regression problem, except that the values y we now want to predict take on only a **small number of discrete values**.

Some Example of Classification problem


- Email : Spam / Not spam
- Tumor: Malignant/ Benign

$$y \in \{0, 1\}$$

0: "Negative Class" (e.g., benign tumor)

1: "Positive Class" (e.g., malignant tumor)

Logistic Regression



Threshold classifier output $h_{\theta}(x)$ at 0.5:

If $h_{\theta}(x) \geq 0.5$, predict "y = 1"

If $h_{\theta}(x) < 0.5$, predict "y = 0"

■ Binary Logistic Regression

- We have a set of feature vectors X with corresponding binary outputs

$$X = \{x_1, x_2, \dots, x_n\}^T$$

$$Y = \{y_1, y_2, \dots, y_n\}^T, \text{ where } y_i \in \{0, 1\}$$

- We want to model $p(y|x)$

$$p(y_i = 1 | x_i, \theta) = \sum_j \theta_j x_{ij} = x_i \theta$$

By definition $p(y_i = 1 | x_i, \theta) \in \{0, 1\}$. We want to transform the probability to remove the range restrictions, as $x_i \theta$ can take any real value.

- **Hypothesis** $h_{\theta}(x) = g(\theta^T x)$

$$= \frac{1}{1 + e^{-\theta^T x}}$$

- **Interpretation**

$$h_{\theta}(x) = p(y = 1|x, \theta)$$

- **Because probabilities should sum to 1, define**

$$p(y = 0|x, \theta) := 1 - p(y = 1|x, \theta)$$

- **If $h_{\theta}(x) = 0.7$ interpret as 70% chance data point belongs to class**
- **If $h_{\theta}(x) \geq 0.5$ classify as *positive sentiment, malignant tumor, ...***

- Odds

p : probability of an event occurring

$1 - p$: probability of the event not occurring

The odds for event i are then defined as

$$odds_i = \frac{p_i}{1 - p_i}$$

Taking the *log* of the odds removes the range restrictions.

$$\log_e \left(\frac{p_i}{1 - p_i} \right) = \sum_j \theta_j x_{ij} = X_i \theta$$


This way we map the probabilities from the $[0; 1]$ range to the entire number line (real value).

Logistic Regression

HYPOTHESIS FUNCTION

We need to calculate P_i

$$\log\left(\frac{p_i}{1-p_i}\right) = x_i\theta$$
$$\frac{p_i}{1-p_i} = e^{x_i\theta}$$
$$p_i = \frac{e^{x_i\theta}}{1+e^{x_i\theta}} = \frac{1}{1+e^{-x_i\theta}}$$



Standard logistic sigmoid function

$$h_{\theta}(x) = \theta^T x$$

$$h_{\theta}(x) = g(\theta^T x)$$

$$p_i = g(\theta x) = \frac{1}{1 + e^{-\theta x}}$$

$$\log_2 8 = 3$$

$$\log_4 16 = 2$$

LOGISTIC REGRESSION MODEL

Linear Regression

$$h_{\theta}(x) = \theta^t x$$

Logistic Regression


$$g(\theta^t x) = \begin{cases} 1, & \frac{1}{1+e^{-\theta^t x}} \geq 0.5 \\ 0, & 1 - \frac{1}{1+e^{-\theta^t x}} < 0.5 \end{cases}$$

$$p(y_i = 1 | x_i, \theta) = \frac{1}{1 + e^{-\theta^t x}}$$

$$p(y_i = 0 | x_i, \theta) = 1 - \frac{1}{1 + e^{-\theta^t x}}$$

$$p(y_i | x_i : \theta) = \left(\frac{1}{1 + e^{-\theta^t x}} \right)^{y_i} \left(1 - \frac{1}{1 + e^{-\theta^t x}} \right)^{1-y_i}$$

Logistic Regression



- **If** $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$

and $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$

- **Prediction $y = 1$ whenever**

$$\begin{aligned} \theta^T x &\geq 0 \\ \Leftrightarrow -3 + x_1 + x_2 &\geq 0 \\ \Leftrightarrow x_1 + x_2 &\geq 3 \end{aligned}$$

Logistic Regression

$$l(\theta) = \prod_{i=1}^m \left(\frac{1}{1 + e^{-\theta^T x}} \right)^{y_i} \left(1 - \frac{1}{1 + e^{-\theta^T x}} \right)^{1-y_i}$$

We can simplify $L(\theta)$ by taking its **log** and then differentiate to get the gradient.

$$j(\theta) = l(\theta) = \sum_1^m \left[y_i \log \left(\frac{1}{1 + e^{-\theta^T x}} \right) + (1 - y_i) \log \left(1 - \frac{1}{1 + e^{-\theta^T x}} \right) \right]$$

$$\begin{aligned} \frac{d}{d\theta} j(\theta) &= \frac{d}{d\theta} \sum_1^m \left[y_i \log \left(\frac{1}{1 + e^{-\theta^T x}} \right) + (1 - y_i) \log \left(1 - \frac{1}{1 + e^{-\theta^T x}} \right) \right] \\ &= \sum_{i=1}^m \left(y_i - \frac{1}{1 + e^{-\theta^T x_i}} \right) x_i \end{aligned}$$

Gradient Descent for logistic regression

- **Gradient Descent to minimize logistic regression cost function**

$$J(\theta) = -\frac{1}{m} \left(\sum_{i=1}^m y^{(i)} \log(h_{\theta}(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)})) \right)$$

with identical algorithm as for linear regression

while not converged:

for all j :

$$tmp_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

$$\theta := \begin{bmatrix} tmp_0 \\ \vdots \\ tmp_n \end{bmatrix}$$

Thank You