

CodeChef Discussion

Search Here...

● questions ○ tags ○ users

ask a question about faq

algorithm to find inverse modulo m

i have two questions how to find inverse of a number modulo m and n! modulo m

2

inverse modulo number theory

asked 26 Jun '12, 13:07

vamsi2708 31-2-3-5 accept rate: 0%

oldest newest most voted 2 Answers:

.....

I hope I understood well, I asked same question (inverse modulo) in OLYMPIC tutorial.

10 $\frac{a}{b}\%P = (a\%P * (b^{P-2})\%P)\%P$

Note: P is prime number

n! is simply

n! = (n%MOD * (n-1)%MOD) % MOD

link

edited 09 Dec '12, 20:03

answered 26 Jun '12, 14:18



betlista ◆◆ 16.5k • 49 • 113 • 220 accept rate: 11%

Suppose we need to calculate nCr, in these cases, n > P. how to handle these cases?

sandipan (29 Dec '13, 20:08)

Firstly, the inverse of an element a in the residue classes modulo m exists if and only if: gcd(a,m) = 1 i.e. they are relatively prime

For finding the inverse, use the extended euclidean algorithm

(http://en.wikipedia.org/wiki/Extended_Euclidean_algorithm)

It finds the solution(x,y) to the following equation:

ax + by = gcd(a,b)

Taking b = m, the equation becomes:

ax + my = gcd(a,m)

since gcd(a,m) = 1

ax + my = 1

If we use the modulo m operation on both sides:

ax(mod m) + my(mod m) = 1(mod m)

ax(mod m) = 1(mod m)

=> x is the inverse of a modulo m

You are not logged in. Please login at www.codechef.com to post yo algorithm:

```
questions!
void EE(int a, int b, int& x, int& y)
   if(a%b == 0)
   {
       x=0;
       y=1;
```

Follow this question By Email:

Once you sign in you will be able to subscribe for any updates here

By RSS:

Answers

Answers and Comments

Tags:

modulo ×94

number ×24

inverse ×13

theory ×5

Asked: 26 Jun '12, 13:07

Seen: 4,929 times

Last updated: 29 Dec '13, 20:08

Related questions

 $X^a = b \pmod{2k + 1}$

Editorial: Wilson's theorem

Modulo inverse calculation verification

inverse modulo for non primes

inverse modulo

is this a property of inverse modulo operation?

finding Inverse Modulo of a range of numbers under modulo m

Decimal to binary conversion.

just a simple sum

modulo calculate

```
return;
         }
         EE(b,a%b,x,y);
          int temp = x;
          x = y;
          y = temp - y*(a/b);
     Using this function and the explanation above, the inverse function can be implemented as follows:
      int inverse(int a, int m)
      {
          int x,y;
         EE(a,m,x,y);
         if(x<0) x += m;
         return x;
      link
                                                                                         answered 10 Dec '12, 15:55
                                                                                             nihalb
                                                                                               171•3•6
                                                                                                accept rate: 0%
[hide preview]
                                                                                                     community wiki
Post Your Answer
```

About CodeChef | About Directi | CEO's Corner CodeChef Campus Chapters | CodeChef For Schools | Contact Us

