

Cell wall dynamics in yeast

Gertien J Smits, Johan C Kapteyn, Herman van den Ende and Frans M Klis*

The yeast *Saccharomyces cerevisiae* is the first fungus for which the structure of the cell wall is known at the molecular level. It is a dynamic and highly regulated structure. This is vividly illustrated when the cell wall is damaged and a salvage pathway becomes active, resulting in compensatory changes in the wall.

Addresses

BioCentrum Amsterdam, University of Amsterdam, Institute of Molecular Cell Biology, Developmental Biology of Fungi, Kruislaan 318, 1098 SM Amsterdam, The Netherlands
*e-mail: klis@bio.uva.nl

Current Opinion in Microbiology 1999, 2:348–352

<http://biomednet.com/elecref/1369527400200348>

© Elsevier Science Ltd ISSN 1369-5274

Abbreviations

CWP cell wall protein
GPI glycosyl phosphatidylinositol
Pir protein with internal repeats
PKC protein kinase C

Introduction

Fungi devote a considerable amount of metabolic energy to building a cell wall, which accounts for 20–30% of the cell dry weight. Not only does the fungal cell wall have a skeletal function, it also plays a key role in morphogenesis and cell–cell recognition. We present a tentative molecular model of the cell wall of *Saccharomyces cerevisiae*. We further show that the composition and structure of the cell wall are strictly regulated and vary in response to a wide range of environmental conditions. We also touch upon the relation between the cell wall and morphogenesis. Finally, we present evidence for the existence of a salvage pathway, designated as the cell wall integrity pathway, allowing the cell to compensate for various forms of cell wall damage. Although the cell wall of *S. cerevisiae* is certainly not representative of all fungi, we believe that our model has a strong predictive value for studying the cell wall of the Ascomycotina, and especially *Candida albicans*.

A molecular model of the cell wall

On the basis of on recent work by various groups ([1–3,4**]; see [5*,6*] for reviews) we present a tentative model of the cell wall of yeast at a molecular level (Figure 1). The main features of this model are discussed below.

Firstly, an internal skeletal framework, formed by a three-dimensional network of β 1,3-glucan molecules, surrounds the entire cell and is largely responsible for the mechanical strength of the wall. Because mature β 1,3-glucan molecules are branched [7], they have multiple nonreducing ends. These may function as attachment sites for the other components of the cell wall [1,2].

Secondly, the skeletal framework is strengthened by chitin chains [1], which are mainly found close to the plasma membrane. Some chitin chains, however, become linked to short side-chains of β 1,6-glucan [8].

Thirdly, mature β 1,6-glucan molecules are mainly found at the outside of the skeletal framework and interconnect a particular class of cell wall proteins (CWPs), glycosyl phosphatidylinositol (GPI)-CWPs, with the framework [2].

Finally, two classes of covalently linked CWPs are known, the already mentioned GPI-CWPs [9,10*,11**] and the protein with internal repeats (Pir)-CWPs [3,4**,12]. They differ from each other in that Pir-CWPs seem to be directly linked to β 1,3-glucan molecules without an interconnecting β 1,6-glucan moiety [4**] and can be released from the cell wall by mild alkali [3]. In contrast to the structural complex GPI-CWP→ β 1,6-glucan→ β 1,3-glucan, which has been extensively investigated [2], the Pir-CWP→ β 1,3-glucan complex is as yet ill defined.

Glucan remodeling and cell wall assembly take place outside the plasma membrane. Likely candidates for glucan remodeling enzymes are Gas1, a GPI-anchored plasma membrane protein (reviewed in [13*]; see also [14**]), and the proteins belonging to the Bgl2 family of endo-beta-1,3-glucanases and/or transglucosylases [15]. Interestingly, homologs of Gas1 have been found in various other (mycelial) fungi [13*,16,17*].


The question arises in how far our model has predictive value for other fungi. The cell wall of *C. albicans*, recently reviewed in [18*], is a good test case. First, there is strong evidence for the existence of a family of GPI-CWPs in *C. albicans* [19**,20]. Although it has not yet been shown that the proteins in this family are linked through an interconnecting β 1,6-glucan moiety to β 1,3-glucan, there is strong evidence for the presence of the structural complex GPI-CWP→ β 1,6-glucan→ β 1,3-glucan [21]. Second, the cell wall of *C. albicans* contains several proteins that like the Pir-CWPs in *S. cerevisiae* can be released by mild alkali [22], suggesting that also in *C. albicans* a Pir-CWP→ β 1,3-glucan-like complex might exist. Another interesting finding concerns a putative GPI-CWP at the cell surface of the mycelial fungus *Penicillium marneffei* [23*]. Taken together, these data strongly indicate that our model has predictive value for other Ascomycotina.

Functions of cell wall proteins

Following a genomic approach, we were able to predict the existence of approximately 40 different GPI-CWPs in yeast [9], and this has been confirmed experimentally

Figure 1

A molecular model of the cell wall of *Saccharomyces cerevisiae*. The internal skeletal layer consists of β 1,3-glucan molecules that form a three-dimensional network surrounding the entire cell. This network is kept together by local alignments between segments of β 1,3-glucan molecules, allowing the formation of multiple hydrogen bridges. At the outside of the skeletal layer, cell wall proteins are linked to the nonreducing ends of β 1,3-glucan molecules either directly (Pir-CWPs) or indirectly through an interconnecting β 1,6-glucan moiety (GPI-CWPs). Some GPI-CWPs, such as Cwp1, may be linked both ways. After cytokinesis, the skeletal layer becomes strengthened by the coupling of chitin chains to nonreducing ends of β 1,3-glucan chains. This takes place mainly at the inside of the skeletal layer. β 1,6-Glucan is much more branched than β 1,3-glucan [7,8], probably explaining why the mature β 1,6-glucan molecule is water-soluble. Thus, β 1,6-glucan probably functions as a flexible tether for GPI-CWPs. Note that branched polysaccharides such as β 1,3-glucan and β 1,6-glucan, in principle,



have a single reducing end and multiple nonreducing ends. For reasons of clarity, non-covalently bound proteins and proteins linked through disulfide bridges to other cell wall

proteins have been omitted. PM, plasma membrane. This model is based on data from [1–3,4**,5*,6*,52,53].

[10*]. Although it has been shown that CWPs, collectively, limit cell wall permeability [24], the function of most individual CWPs remains a mystery. Some are clearly involved in cell–cell adhesion such as the flocculins Flo1, Flo5, Flo9, and Flo10 [25] and the sexual agglutinins Aga1,


Aga2, and Sag1 [26]. Flo11 forms a special case [27,28**,29]. Discovered first as another flocculin, Flo11 appears to be required for invasive growth into agar, and possibly also pseudohyphal growth in response to nitrogen starvation [28**,29,30*]. This indicates that Flo11 may play

Table 1

Overview of CWP-encoding genes that vary in expression levels during the cell cycle and under various growth conditions.

Signal	Regulated genes	References
Cell cycle		
G1	<i>PRY3, YGR189c, YNL300w</i>	[32,33**,34,35]
G2	<i>CIS3, CWP1, CWP2, TIR1</i>	
M	<i>SED1, YIP1, YOR383c</i>	
M/G1	<i>AGA1, AGA2, EGT2, HSP150, PIR1, PIR3, SAG1, TIP1, TIR5, UTR2, YER150c, YHR126c, YLR194c</i>	
Pheromone	<i>AGA1, CWP1, FIG2, SAG1</i>	[26,31**]
Nutrients		
Carbon source	<i>CIS3, CWP1, EGT2, FLO1, FLO5, FLO9, SED1, SUN4, TIP1, TIR1, TIR6, UTR2, YER150w, YOR383c</i>	[25,36,38*,39]*
Nitrogen	<i>FLO11, HSP150</i>	[28**,41]
Sporulation	<i>AGA2, BAR1, CIS3, CWP1, CWP2, EGT2, FIG2, FLO11, HSP150, PIR1, PIR3, SED1, TIP1, TIR2, TIR5, UTR2, YDR134c, YER150w, YGR189c, YIB1, YLR110c, YLR194c, YOL155c, YOR214c, YOR382w, YOR383c</i>	[37**]
Stress		
Weakened wall	<i>CIS3, CWP1, HSP150, PIR3, SED1, SSR1</i>	[4**,14**]†
Temperature	<i>HSP150, TIP1, TIR1, TIR2</i>	
Hypoxia	<i>DAN1, TIP1, TIR1</i>	[39,40,54]
Aluminum	<i>SED1, HSP150</i>	[55]

*T Fujii, H Shimoi, I Fujishige, T Ohba, abstract 246, Yeast Genetics and Molecular Biology Meeting, College Park, Maryland, July 28 to August 2 1998. †JC Kapteyn, unpublished data.

Figure 2

Model for the cell wall integrity pathway. Cell wall stress is sensed at the cell surface by sensor membrane proteins, for which the Wsc and Mid proteins are likely candidates. The signal from Slg1/Wsc1 is relayed through the phosphatidylinositol-4 kinase Tor2 to the exchange factor Rom2, which activates Rho1. Rho1 can directly activate the glucan synthases Fks1 and Fks2, as well as the protein kinase C (PKC) pathway, and thus leads to increased β 1,3-glucan synthesis and, through the Slt2/Mpk1 MAP kinase cascade and an as yet unidentified transcription factor (represented as Txn factor), to increased expression of cell wall biosynthetic enzymes and CWPs. Whether alterations in chitin synthesis, and possibly also in β 1,6-glucan synthesis, and the altered cross-linking of proteins to the cell wall matrix are achieved through the same signal transduction pathway remains to be clarified.

a role in determining cellular morphology. Another GPI-CWP, Fig2, also seems to affect cell morphology. When *fig2Δ* haploid cells mate, they form a narrow mating projection and fusion bridge, which interferes with nuclear fusion and migration [31^{••}]. Finally, Egt2, which is also a GPI-CWP, seems to be required for cell separation [32].

Several genes encoding CWPs show homology to the glucoamylase gene *STA1*, which is specific for the variant strain *S. cerevisiae* var. *diastaticus*. *STA1*, however, is probably the result of a recombination event between *FLO11* and *SGA1*, the normal glucoamylase. As the homology is limited to the *FLO11* domain, this probably excludes a glucoamylase-like function for the gene products involved.

Regulation of cell wall protein expression

The recent transcript profiling studies strongly suggest that expression of many CWPs is cell cycle regulated

[33^{••},34,35] and is affected by nutrient availability [36,37^{••}]. It is further known that pheromones and various other environmental conditions affect the expression of CWP-encoding genes (Table 1; see also below).

Nutrient availability and environmental conditions

Batch-cultured cells growing on glucose rapidly consume the glucose by fermentation, thereby producing ethanol. Subsequently, they switch to respiratory growth and use up the ethanol before entering stationary phase. The transcript levels of many CWP-encoding genes change when the cells switch to respiratory growth or enter stationary phase [36] (Table 1). For example, *Sed1*, a GPI-CWP [10[•]], becomes a major CWP in stationary phase cells [38[•]]. Fermentative growth can also be triggered by hypoxic conditions, and a similar set of CWPs is then induced. Other wall proteins, however, are specifically induced under hypoxic conditions, indicating that they are regulated differently [39,40] (Table 1).

When diploid yeast cells are starved for nitrogen, they switch to pseudohyphal growth. Interestingly, this is accompanied by increased expression of *FLO11* [28^{••},30[•]] and presumably also of *HSP150/PIR2* [41], raising the question whether still more CWPs are preferentially used by the cell during pseudohyphal growth. Without both suitable carbon and nitrogen sources, cells activate the sporulation program. This is accompanied by up- and down-regulation of various CWP-encoding genes [37^{••}] (Table 1).

The cell cycle

Approximately 13% of the genes of *S. cerevisiae* are regulated in a cell-cycle-dependent manner [33^{••}]. Intriguingly, more than half of all CWP-encoding genes (22 out of 43) are cell cycle regulated, including *PIR1*, the most strictly cell-cycle-regulated gene in yeast. Although specific CWP-encoding genes seem to be expressed during each phase of the cell cycle [33^{••},34,35], most of them are active in late M and early G1 phase, around the time of cell separation and the subsequent period of isotropic growth by the daughter cell (Table 1). Finally, consistent with the extensive cell-cycle-dependent expression of CWPs, some CWPs are indeed known to be localized to specific regions of the cell wall [42,43[•]].

When haploid yeast cells sense the mating pheromone of the opposite mating type, they arrest in G1 and form a mating projection [44^{••}]. The density of sexual agglutinins in the wall of the mating projection dramatically increases [26]. In addition, as discussed above, expression of *FIG2* [31^{••}], which encodes a putative GPI-CWP, is upregulated. Also more chitin is deposited in the wall of the mating projection [26]. These observations clearly indicate that the wall of the mating projection differs from normal walls.

The cell wall integrity pathway

There is increasing evidence that weakening of the cell wall results in activation of a salvage pathway, leading to compensatory changes in the wall. We propose to call it the

cell wall integrity pathway. The existence of such a pathway might explain why so many cell wall mutants show hypersensitivity to caffeine [45]. Caffeine activates protein kinase A, which represses many stress responses [46]. Thus, mutant cells might be hypersensitive to caffeine because they depend on the cell wall integrity pathway for their survival.

A speculative scheme of the cell wall integrity pathway is presented in Figure 2. The sensing of cell wall weakening, possibly through membrane stretch, is thought to occur by the Wsc and Mid families of membrane proteins [44**]. The signal is believed to be relayed to Rho1, a small GTP-binding protein, through Tor2, a phosphatidylinositol-4 kinase, and the exchange factor Rom2 [47,48**]. Rho1 modulates the protein kinase C (PKC) pathway [49*] as well as the β 1,3-glucan synthase, resulting in increased synthesis of chitin and glucan [8,14**,50*], altered cross-linking of glucan and proteins [4**,8], and increased expression of Cwp1 and Pir-CWPs [4**,8,14**]. Other stress conditions that are also relayed through the PKC pathway, such as high temperatures and low osmolarity, might be sensed by the same or similar membrane sensors and lead to similar effects. This might be relevant for several CWPs (*HSP150/PIR2*, *TIP1* and *TIR2*), which are more strongly expressed at high temperatures [41,51].

Perspectives

Although many questions concerning cell wall biogenesis remain unanswered, our model of the cell wall allows to formulate them in molecular terms. Its predictive value concerning mycelial fungi is an important question that needs to be further addressed. It is also clear that the wall is highly dynamic and forms an integral part of cell metabolism, raising fascinating questions about the control mechanisms involved.

Acknowledgements

We thank our colleagues for many stimulating discussions. This work has received support from the University of Amsterdam, NWO, STW, Unilever, and the European Union (EUROFAN II).

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Kollár R, Petraková E, Ashwell G, Robbins PW, Cabib E: **Architecture of the yeast cell wall: the linkage between chitin and β (1 \rightarrow 3)-glucan.** *J Biol Chem* 1995, **270**:1170-1178.
 2. Kollár R, Reinhold BB, Petraková E, Yeh HJC, Ashwell G, Drgonová J, Kapteyn JC, Klis FM, Cabib E: **Architecture of the yeast cell wall: β (1 \rightarrow 6)-glucan interconnects mannoprotein, β (1 \rightarrow 3)-glucan, and chitin.** *J Biol Chem* 1997, **272**:17762-17788.
 3. Mrsa V, Seidl T, Gentzsch M, Tanner W: **Specific labelling of cell wall proteins by biotinylation. Identification of four covalently linked O-mannosylated proteins of *Saccharomyces cerevisiae*.** *Yeast* 1997, **13**:1145-1154.
 4. Kapteyn JC, Van Egmond P, Sievi E, Van den Ende H, Makarow M, Klis FM: **The contribution of the O-glycosylated protein Pir2p/Hsp-150 to the construction of the yeast cell wall in wild-type cells and β 1,6-glucan-deficient mutants.** *Mol Microbiol* 1999, **31**:1835-1844.
- Pir-cell wall proteins (CWPs) are attached to β 1,3-glucan through a base-labile linkage. The cell compensates for decreased levels of β 1,6-glucan in

its wall by increased chitin deposition in the lateral walls. In addition, the transcript levels of Pir-CWP encoding genes and *FKS2* are up-regulated.

5. Lipke PN, Ovalle R: **Cell wall architecture in yeast: new structure and new challenges.** *J Bacteriol* 1998, **180**:3735-3740.
This review describes the cell wall as a latticework of structural units, each being composed of a branched β 1,3-glucan molecule, presenting multiple attachment sites for both β 1,6-glucan and chitin chains.
6. Kapteyn JC, Van Den Ende H, Klis FM: **The contribution of cell wall proteins to the organization of the yeast cell wall.** *Biochim Biophys Acta* 1999, **1426**:373-383.
This review proposes that the cell wall contains two structural units, one having the core structure GPI-CWP \rightarrow β 1,6-glucan \rightarrow β 1,3-glucan, and the other having the structure Pir-CWP- β 1,3-glucan.
7. Manners DJ, Masson AJ, Patterson JC: **The structure of a β -(1-3)-D-glucan from yeast cell walls.** *Biochem J* 1973, **135**:19-30.
8. Kapteyn JC, Ram AF, Groos EM, Kollar R, Montijn RC, Van Den Ende H, Llobell A, Cabib E, Klis FM: **Altered extent of cross-linking of β 1,6-glucosylated mannoproteins to chitin in *Saccharomyces cerevisiae* mutants with reduced cell wall β 1,3-glucan content.** *J Bacteriol* 1997, **179**:6279-6284.
9. Caro LHP, Tettelin H, Vossen JH, Ram AF, van den Ende H, Klis FM: **In silico identification of glycosyl phosphatidylinositol-anchored plasma-membrane and cell wall proteins of *Saccharomyces cerevisiae*.** *Yeast* 1997, **13**:1477-1489.
10. Hamada K, Fukuchi S, Arisawa M, Baba M, Kitada K: **Screening for glycosylphosphatidylinositol (GPI)-dependent cell wall proteins in *Saccharomyces cerevisiae*.** *Mol Gen Genet* 1998, **258**:53-59.
The identification of potential GPI-cell wall proteins using a reporter construct is described.
11. Hamada K, Terashima H, Arisawa M, Kitada K: **Amino acid sequence requirement for efficient incorporation of glycosylphosphatidylinositol-associated proteins into the cell wall of *Saccharomyces cerevisiae*.** *J Biol Chem* 1998, **273**:26946-26953.
A short amino acid region upstream of the ω site for glycosyl phosphatidylinositol (GPI) attachment directs GPI-cell wall proteins to the cell wall.
12. Yun DJ, Zhao Y, Pardo JM, Narasimhan ML, Damsz B, Lee H, Abad LR, D'Urzo MP, Hasegawa PM, Bressan RA: **Stress proteins on the yeast cell surface determine resistance to osmotic stress, a plant antifungal protein.** *Proc Natl Acad Sci USA* 1997, **94**:7082-7087.
13. Popolo L, Vai M: **The Gas1 glycoprotein, a putative wall polymer cross-linker.** *Biochim Biophys Acta* 1999, **1426**:385-400.
This review discusses the evidence that Gas1 and homologs of Gas1 in other fungi function as transglucosylases involved in maturation of β 1,3-glucan.
14. Ram AFJ, Kapteyn JC, Montijn RC, Caro LHP, Douwes JE, Baginsky W, Mazur P, Van Den Ende H, Klis FM: **Loss of the plasma membrane-bound protein Gas1p in *Saccharomyces cerevisiae* results in the release of β 1,3-glucan into the medium and induces a compensation mechanism to ensure cell wall integrity.** *J Bacteriol* 1998, **180**:1418-1424.
The cell compensates for decreased levels of β 1,3-glucan in its wall by enhancing the levels of chitin and mannan, and increasing expression of *CWP1* and *FKS2*.
15. Capparello C, Mrsa V, Tanner W: **New potential cell wall glucanases in *Saccharomyces cerevisiae* and their involvement in mating.** *J Bacteriol* 1998, **180**:5030-5037.
16. Popolo L, Vai M: **Defects in assembly of the extracellular matrix are responsible for altered morphogenesis of a *Candida albicans* *phr1* mutant.** *J Bacteriol* 1998, **180**:163-166.
17. Nakazawa T, Horiuchi H, Ohta A, Takagi M: **Isolation and characterization of *EPD1*, an essential gene for pseudohyphal growth of a dimorphic yeast *Candida maltosa*.** *J Bacteriol* 1998, **180**:2079-2086.
The GAS1 homolog *EPD1* in *Candida maltosa* is essential for pseudohyphal growth.
18. Chaffin WL, Lopez-Ribot JL, Casanova M, Gozalbo D, Martinez JP: **Cell wall and secreted proteins of *Candida albicans*: identification, function, and expression.** *Microbiol Mol Biol Rev* 1998, **62**:130-180.
Extensive overview of cell wall-associated proteins in *Candida albicans*.
19. Hoyer LL, Payne TL, Hecht JE: **Identification of *Candida albicans* *ALS2* and *ALS4* and localization of Als proteins to the fungal cell surface.** *J Bacteriol* 1998, **180**:5334-5343.
The cell wall of *Candida albicans* contains a family of glycosyl phosphatidylinositol linked cell wall proteins, encoded by the *ALS* genes.
20. Hoyer LL, Payne TL, Bell M, Myers AM, Scherer S: ***Candida albicans* *ALS3* and insights into the nature of the *ALS* gene family.** *Curr Genet* 1998, **33**:451-459.

21. Kapteyn JC, Montijn RC, Dijkgraaf GJP, van den Ende H, Klis FM: **Covalent association of β -1,3-glucan with β -1,6-glucosylated mannoproteins in cell walls of *Candida albicans*.** *J Bacteriol* 1995, **177**:3788-3792.
 22. Mormeneo S, Marcilla A, Irazo M, Sentandreu R: **Structural mannoproteins released by β -elimination from *Candida albicans* cell walls.** *FEMS Microbiol Lett* 1994, **123**:131-136.
 23. Cao L, Chan C-M, Lee C, Wong SS-Y, Yuen K-Y: **MP1 encodes an abundant and highly antigenic cell wall mannoprotein in the pathogenic fungus *Penicillium marneffei*.** *Infect Immun* 1998, **66**:966-973.
- The cell wall of the mycelial fungus *Penicillium marneffei* contains a glycosyl phosphatidylinositol linked cell wall protein.
24. De Nobel JG, Klis FM, Priem J, Munnik T, van den Ende H: **The glucanase-soluble mannoproteins limit cell wall porosity in *Saccharomyces cerevisiae*.** *Yeast* 1990, **6**:491-499.
 25. Teunissen AW, Steensma HY: **The dominant flocculation genes of *Saccharomyces cerevisiae* constitute a new subtelomeric gene family.** *Yeast* 1995, **11**:1001-1013.
 26. Lipke PN, Kurjan J: **Sexual agglutination in budding yeasts: structure, function, and regulation of adhesion glycoproteins.** *Microbiol Rev* 1992, **56**:180-194.
 27. Lo WS, Dranginis AM: **FLO11, a yeast gene related to the STA genes, encodes a novel cell surface flocculin.** *J Bacteriol* 1996, **178**:7144-7151.
 28. Lo WS, Dranginis AM: **The cell surface flocculin Flo11 is required for pseudohyphae formation and invasion by *Saccharomyces cerevisiae*.** *Mol Biol Cell* 1998, **9**:161-171.
- FLO11 encodes a glycosyl phosphatidylinositol linked cell wall protein essential for invasive growth.
29. Robertson LS, Fink GR: **The three yeast A kinases have specific signaling functions in pseudohyphal growth.** *Proc Natl Acad Sci USA* 1998, **95**:13783-13787.
 30. Rupp S, Summers E, Lo HJ, Madhani H, Fink G: **MAP kinase and cAMP filamentation signaling pathways converge on the unusually large promoter of the yeast FLO11 gene.** *EMBO J* 1999, **18**:1257-1269.
- Two signal transduction pathways mediate FLO11 gene expression, namely the cAMP/PKA pathway and the Kss1 mitogen activated protein kinase (MAPK) cascade.
31. Erdman S, Lin L, Malczynski M, Snyder M: **Pheromone-regulated genes required for yeast mating differentiation.** *J Cell Biol* 1998, **140**:461-483.
- In the absence of the glycosyl phosphatidylinositol linked cell wall protein (GPI-CWP) Fig2, an abnormally narrow conjugation bridge is formed between two mating cells.
32. Kovacech B, Nasmyth K, Schuster T: **EGT2 gene transcription is induced predominantly by Swi5 in early G1.** *Mol Cell Biol* 1996, **16**:3264-3274.
 33. Spellman PT, Sherlock G, Zhang MQ, Iyer VR, Anders K, Eisen MB, Brown PO, Botstein D, Futcher B: **Comprehensive identification of cell cycle-regulated genes of the yeast *Saccharomyces cerevisiae* by microarray hybridization.** *Mol Biol Cell* 1998, **9**:3273-3297.
- The expression of many Pir-CWPs, GPI-CWPs, and proteins involved in cell wall construction is cell cycle regulated.
34. Cho RJ, Campbell MJ, Winzeler EA, Steinmetz L, Conway A, Wodicka L, Wolfsberg TG, Gabrielian AE, Landsman D, Lockhart DJ, Davis RW: **A genome-wide transcriptional analysis of the mitotic cell cycle.** *Mol Cell* 1998, **2**:65-73.
 35. Caro LHP, Smits GJ, van Egmond P, Chapman JW, Klis FM: **Transcription of multiple cell wall protein-encoding genes in *Saccharomyces cerevisiae* is differentially regulated during the cell cycle.** *FEMS Microbiol Lett* 1998, **161**:345-349.
 36. DeRisi JL, Iyer VR, Brown PO: **Exploring the metabolic and genetic control of gene expression on a genomic scale.** *Science* 1997, **278**:680-686.
 37. Chu S, DeRisi J, Eisen M, Mulholland J, Botstein D, Brown PO, Herskowitz I: **The transcriptional program of sporulation in budding yeast.** *Science* 1998, **282**:699-705.
- During sporulation the transcription of a large group of genes encoding cell wall proteins is up-regulated.
38. Shimoi H, Kitagaki H, Ohmori H, Iimura Y, Ito K: **Sed1p is a major cell wall protein of *Saccharomyces cerevisiae* in the stationary phase and is involved in lytic enzyme resistance.** *J Bacteriol* 1998, **180**:3381-3387.
- The glycosyl phosphatidylinositol linked cell wall protein Sed1 is highly expressed in stationary phase cells, giving protection against lytic enzymes.
39. Donzeau M, Bourdineaud JP, Lauquin GJ: **Regulation by low temperatures and anaerobiosis of a yeast gene specifying a putative GPI-anchored plasma membrane protein.** *Mol Microbiol* 1996, **20**:449-459.
 40. Sertil O, Cohen BD, Davies KJ, Lowry CV: **The DAN1 gene of *S. cerevisiae* is regulated in parallel with the hypoxic genes, but by a different mechanism.** *Gene* 1997, **192**:199-205.
 41. Russo P, Simonen M, Uimari A, Teesalu T, Makarow M: **Dual regulation by heat and nutrient stress of the yeast HSP150 gene encoding a secretory glycoprotein.** *Mol Gen Genet* 1993, **239**:273-280.
 42. Bony M, Barre P, Blondin B: **Distribution of the flocculation protein, Flop, at the cell surface during yeast growth: the availability of Flop determines the flocculation level.** *Yeast* 1998, **14**:25-35.
 43. Ram AF, Van den Ende H, Klis FM: **Green fluorescent protein-cell wall fusion proteins are covalently incorporated into the cell wall of *Saccharomyces cerevisiae*.** *FEMS Microbiol Lett* 1998, **162**:249-255.
- Green fluorescent protein-tagging of the glycosyl phosphatidylinositol linked cell wall protein Cwp1 and Cwp2 shows that they are targeted to specific areas of the cell wall.
44. Gustin MC, Albertyn J, Alexander M, Davenport K: **MAP kinase pathways in the yeast *Saccharomyces cerevisiae*.** *Microbiol Mol Biol Rev* 1998, **62**:1264-1300.
- Superb review of the different mitogen activated protein (MAP) kinase pathways in yeast, and their role in controlling cell morphology, and the composition and molecular organization of the cell wall.
45. Lussier M, White AM, Sheraton J, di Paolo T, Treadwell J, Southard SB, Horenstein CI, Chen-Weiner J, Ram AF, Kapteyn JC *et al.*: **Large scale identification of genes involved in cell surface biosynthesis and architecture in *Saccharomyces cerevisiae*.** *Genetics* 1997, **147**:435-450.
 46. Kronstad J, De Maria AD, Funnell D, Laidlaw RD, Lee N, de Sa MM, Ramesh M: **Signaling via cAMP in fungi: interconnections with mitogen-activated protein kinase pathways.** *Arch Microbiol* 1998, **170**:395-404.
 47. Schmidt A, Bickle M, Beck T, Hall MN: **The yeast phosphatidylinositol kinase homolog TOR2 activates RHO1 and RHO2 via the exchange factor ROM2.** *Cell* 1997, **88**:531-542.
 48. Bickle M, Delley PA, Schmidt A, Hall MN: **Cell wall integrity modulates RHO1 activity via the exchange factor ROM2.** *EMBO J* 1998, **17**:2235-2245.
- Destabilization of the cell wall activates Rho1, the regulatory subunit of β 1,3-glucan synthase and activator of the Pkc1 MAP kinase cascade, through the exchange factor Rom2.
49. Cabib E, Drgonová J, Drgon T: **Role of small G proteins in yeast cell polarization and wall biosynthesis.** *Annu Rev Biochem* 1998, **67**:307-333.
- Lucid review of the role of Rho1 in β 1,3-glucan synthesis and the cell wall integrity pathway (Pkc1 MAP kinase pathway).
50. Dallies N, François J, Paquet V: **A new method for quantitative determination of polysaccharides in the yeast cell wall. Application to the cell wall defective mutants of *Saccharomyces cerevisiae*.** *Yeast* 1998, **14**:1297-1306.
- Presents an improved method for determining the sugar composition of yeast cell walls. Various cell wall defective mutants show strongly increased levels of chitin.
51. Kowalski LRZ, Kondo K, Inouye M: **Cold-shock induction of a family of TIP1-related proteins associated with the membrane in *Saccharomyces cerevisiae*.** *Mol Microbiol* 1995, **15**:341-353.
 52. Manners DJ, Masson AJ, Patterson JC, Bjorndal H, Lindberg B: **The structure of a β -(1-6)-D-glucan from yeast cell walls.** *Biochem J* 1973, **135**:31-36.
 53. Rees DA: *Polysaccharide Shapes*. London: Chapman and Hall; 1977.
 54. Kitagaki H, Shimoi H, Ito K: **Identification and analysis of a static culture-specific cell wall protein, Tir1p/Srp1p in *Saccharomyces cerevisiae*.** *Eur J Biochem* 1997, **249**:343-349.
 55. Ezaki B, Gardner RC, Ezaki Y, Kondo H, Matsumoto H: **Protective roles of two aluminum (Al)-induced genes, HSP150 and SED1 of *Saccharomyces cerevisiae*, in Al and oxidative stresses.** *FEMS Microbiol Letts* 1998, **159**:99-105.