


A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion



INSTITUTE OF ECONOMIC STUDIES, FACULTY OF SOCIAL SCIENCES  
CHARLES UNIVERSITY IN PRAGUE (*established 1348*)

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion



# *ROBUST STATISTICS AND ECONOMETRICS*

INSTITUTE OF ECONOMIC STUDIES  
FACULTY OF SOCIAL SCIENCES  
CHARLES UNIVERSITY IN PRAGUE

JAN ÁMOS VÍŠEK

Week 1

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion


## First of all - the (technical) framework of the course



## First of all - the (technical) framework of the course

### Technicalities:

- 1 The text of lecture (both - the full lecture and the handout) will be available on web (at least) from Monday (preceding the lecture), 2 p.m.



## First of all - the (technical) framework of the course

### Technicalities:

- 1 The text of lecture (both - the full lecture and the handout)  
will be available on web (at least) from Monday  
(preceding the lecture), 2 p.m.
- 2 An active presence on the seminars will be a necessary  
and (nearly) sufficient condition for passing the course -  
- the test at the end will be oriented on “ideas”  
not on “formulas”.

## First of all - the (technical) framework of the course

### Technicalities:

- ① The text of lecture (both - the full lecture and the handout)  
will be available on web (at least) from Monday  
(preceding the lecture), 2 p.m.
- ② An active presence on the seminars will be a necessary  
and (nearly) sufficient condition for passing the course -  
- the test at the end will be oriented on “ideas”  
not on “formulas”.
- ③ Any comment will be welcomed  
(to content, on an explanation, correction of ..., etc..)

## First of all - the (technical) framework of the course

### Technicalities:

- ① The text of lecture (both - the full lecture and the handout)  
will be available on web (at least) from Monday  
(preceding the lecture), 2 p.m.
- ② An active presence on the seminars will be a necessary  
and (nearly) sufficient condition for passing the course -  
- the test at the end will be oriented on “ideas”  
not on “formulas”.
- ③ Any comment will be welcomed  
(to content, on an explanation, correction of ..., etc..)
- ④ I encourage You to be active also on lecture -  
- don't let me escape from any topic without understanding it.

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

## The character of lectures and of seminars






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion


## The character of lectures and of seminars

- 1 The lectures will be oriented on ideas -  
- there will be pictures explaining them or creating an inspiration.



## The character of lectures and of seminars

- 1 The lectures will be oriented on ideas -
  - there will be pictures explaining them or creating an inspiration.
- 2 Not to disappoint (completely) those
  - who came for an exact mathematics
  - some pattern of mathematics will be included.



## The character of lectures and of seminars

- 1 The lectures will be oriented on ideas -
  - there will be pictures explaining them or creating an inspiration.
- 2 Not to disappoint (completely) those
  - who came for an exact mathematics
  - some pattern of mathematics will be included.
- 3 Some excursions to mathematics will be of general interest -
  - e. g. You can learn how it is with infinity,  
what is countable and uncountable.

## The character of lectures and of seminars


- 1 The lectures will be oriented on ideas -
  - there will be pictures explaining them or creating an inspiration.
- 2 Not to disappoint (completely) those
  - who came for an exact mathematics
  - some pattern of mathematics will be included.
- 3 Some excursions to mathematics will be of general interest -
  - e. g. You can learn how it is with infinity,  
what is countable and uncountable.
- 4 There will be some quick reminder(s)
  - of something from statistics and econometrics, of history, etc.
  - and at the end of term - if we will have some time
  - a drop of philosophy.

## The character of lectures and of seminars

- 1 The lectures will be oriented on ideas -
  - there will be pictures explaining them or creating an inspiration.
- 2 Not to disappoint (completely) those
  - who came for an exact mathematics
  - some pattern of mathematics will be included.
- 3 Some excursions to mathematics will be of general interest -
  - e. g. You can learn how it is with infinity,  
what is countable and uncountable.
- 4 There will be some quick reminder(s)
  - of something from statistics and econometrics, of history, etc.
  - and at the end of term - if we will have some time
  - a drop of philosophy.
- 5 On seminars - which will be completely under governance
  - of Tomáš Křehlík, we assume mainly some exercises with software
  - but also Your active role with creating them
  - just to fulfill the word “seminar”.


# Content of lecture

- 1 A very first insight into robustness
  - An attractive application as foreword




## Content of lecture

- 1 A very first insight into robustness
  - An attractive application as foreword
- 2 A bit more modest motivation




# Content of lecture


- 
- 1 A very first insight into robustness
    - An attractive application as foreword
  - 2 A bit more modest motivation
  - 3 Let's start more serious discussion
    - Vulnerability of classical procedures to contamination
    - The classical requirements on estimators



# Content


- 
- 1 A very first insight into robustness
    - An attractive application as foreword
  - 2 A bit more modest motivation
  - 3 Let's start more serious discussion
    - Vulnerability of classical procedures to contamination
    - The classical requirements on estimators

# Content

- 
- 1 A very first insight into robustness
    - An attractive application as foreword
  - 2 A bit more modest motivation
  - 3 Let's start more serious discussion
    - Vulnerability of classical procedures to contamination
    - The classical requirements on estimators


## ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994

### *Number of industries 91*

- 
- $X_\ell$  - export from  $i$ -th industry,
  - $US_\ell$  - number of university-passed employees in the  $i$ -th industry,
  - $HS_\ell$  - number of high school-passed employees in the  $i$ -th industry,
  - $VA_\ell$  - value added in the  $i$ -th industry,
  - $K_\ell$  - capital in the  $i$ -th industry,
  - $CR_\ell$  - percentage of market occupied by 3 largest producers,
  - $TFPW_\ell$  - by wages normed productivity in the  $i$ -th industry,
  - $Bal_\ell$  - Balasa index in the  $i$ -th industry,
  - $DP_\ell$  - cost discontinuity in 1993 in the  $i$ -th industry
  - etc., about 20 explanatory variables

## ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994

### *Number of industries 91*



$X_\ell$	-	<i>export from <math>i</math>-th industry,</i>
$US_\ell$	-	<i>number of university-passed employees in the <math>i</math>-th industry,</i>
$HS_\ell$	-	<i>nuber of high school-passed employees in the <math>i</math>-th industry,</i>
$VA_\ell$	-	<i>value added in the <math>i</math>-th industry,</i>
$K_\ell$	-	<i>capital in the <math>i</math>-th industry,</i>
$CR_\ell$	-	<i>percentage of market occupied by 3 largest producers,</i>
$TFPW_\ell$	-	<i>by wages normed productivity in the <math>i</math>-th industry,</i>
$Bal_\ell$	-	<i>Balasa index in the <math>i</math>-th industry,</i>
$DP_\ell$	-	<i>cost discontinuity in 1993 in the <math>i</math>-th industry</i>
	-	<i>etc., about 20 explanatory variables</i>

NO REASONABLE MODEL BY OLS - COEFFICIENT OF DETERMINATION 0.28

ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994  
BY MEANS OF THE *Least Trimmed Squares*

*has found:*

MAIN SUBGROUP

*with number of industries 54 and the model*

$$\frac{X_\ell}{S_\ell} = 4.64 - 0.032 \cdot \frac{US_\ell}{VA_\ell} - 0.022 \cdot \frac{HS_\ell}{VA_\ell} - 0.124 \cdot \frac{K_\ell}{VA_\ell} + 1.035 \cdot CR_\ell \\ - 3.199 \cdot TFPW_\ell + 1.048 \cdot BAL_\ell + 0.452 \cdot DP_\ell + \varepsilon_\ell$$

- $X_\ell$  - export from  $i$ -th industry,
- $US_\ell$  - number of university-passed employees in the  $i$ -th industry,
- $HS_\ell$  - number of high school-passed employees in the  $i$ -th industry,
- $VA_\ell$  - value added in the  $i$ -th industry,
- $K_\ell$  - capital in the  $i$ -th industry,
- $CR_\ell$  - percentage of market occupied by 3 largest producers,
- $TFPW_\ell$  - by wages normed productivity in the  $i$ -th industry,
- $Bal_\ell$  - Balasa index in the  $i$ -th industry,
- $DP_\ell$  - cost discontinuity in 1993 in the  $i$ -th industry

with coefficient of determination 0.97 and stable submodels

ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994  
BY MEANS OF THE *Least Trimmed Squares*

*has found:*

COMPLEMENTARY SUBGROUP

*with number of industries 33 and the model*

$$\frac{X_\ell}{S_\ell} = -0.634 + 0.089 \cdot \frac{US_\ell}{VA_\ell} + 0.235 \cdot \frac{HS_\ell}{VA_\ell} + 0.249 \cdot \frac{K_\ell}{VA_\ell} + 1.174 \cdot CR_\ell \\ + 0.690 \cdot TFPW_\ell + 2.691 \cdot BAL_\ell - 0.051 \cdot DP_\ell + \varepsilon_\ell$$


- $X_\ell$  - export from  $i$ -th industry,
- $US_\ell$  - number of university-passed employees in the  $i$ -th industry,
- $HS_\ell$  - number of high school-passed employees in the  $i$ -th industry,
- $VA_\ell$  - value added in the  $i$ -th industry,
- $K_\ell$  - capital in the  $i$ -th industry,
- $CR_\ell$  - percentage of market occupied by 3 largest producers,
- $TFPW_\ell$  - by wages normed productivity in the  $i$ -th industry,
- $Bal_\ell$  - Balasa index in the  $i$ -th industry,
- $DP_\ell$  - cost discontinuity in 1993 in the  $i$ -th industry

*with coefficient of determination 0.93 and stable submodels*

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

An attractive application as foreword

ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994  
BY MEANS OF THE Least Trimmed Squares.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

An attractive application as foreword

ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994  
BY MEANS OF THE Least Trimmed Squares.



RELATION BETWEEN  $K/W$  AND  $L/S$  FOR THE WHOLE DATA.






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

An attractive application as foreword

ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994  
BY MEANS OF THE Least Trimmed Squares.



RELATION BETWEEN  $K/W$  AND  $L/S$  FOR THE Main subpopulation

(LEFT PICTURE)

AND FOR THE Complementary subpopulation


(RIGHT PICTURE).

ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994  
BY MEANS OF THE Least Trimmed Squares.



Cobb, C., Douglas, P.H. (1928): A Theory of Production.

*American Economic Review*, 18, 139-165.



RELATION BETWEEN  $K/W$  AND  $L/S$  FOR THE Main subpopulation

(LEFT PICTURE)

AND FOR THE Complementary subpopulation


(RIGHT PICTURE).

*It seems we have at hand a miraculous method*


WARNING !!!

We haven't reached something which is  
*"BOMB und IDIOTEN SICHER"*  
but which is the powerful tool, if used with a care.

# Content


- 
- 1 A very first insight into robustness
    - An attractive application as foreword
  - 2 A bit more modest motivation
  - 3 Let's start more serious discussion
    - Vulnerability of classical procedures to contamination
    - The classical requirements on estimators

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

## Hertzsprung-Russell diagram




## Hertzsprung-Russell diagram



Humphreys, R. M. (1978): Studies of luminous stars in nearby galaxies. Supergiant and O stars in the milky way.  
*Astrophysical Journal Supplement Ser.*, 38, 309 - 350.


A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

*A model - we can expect - can't we?*



*A model - we can expect - can't we?*



6.5

## REGRESSION MODEL

$$\begin{aligned} Y_i &= X_i' \beta^0 + e_i \\ &= X_{i1} \beta_1^0 + X_{i2} \beta_2^0 + \dots + X_{ip} \beta_p^0 + e_i, \end{aligned}$$

$i = 1, 2, \dots, n$

$Y_i$	-	RESPONSE VARIABLE	(for $i$ -th object, known)
$X_i \in R^p$	-	EXPLANATORY VARIABLES	(for $i$ -th object, known)
$\beta^0$	-	<u>REGRESSION COEFFICIENTS</u>	("true", unknown)
$e_i$	-	DISTURBANCES, ERROR TERM	(for $i$ -th object, unknown)

3.5

4.6

4.4

4.2

4

3.8

3.6

3.4

SURFACE TEMPERATURE

6.5

## REGRESSION MODEL

$$Y_i = X_i' \beta^0 + e_i$$

$$= X_{i1}\beta_1^0 + X_{i2}\beta_2^0 + \dots + X_{ip}\beta_p^0 + e_i,$$

$i = 1, 2, \dots, n$

$Y_i$

-

RESPONSE VARIABLE

(for  $i$ -th object, known)

Galton, F. (1886): Regression towards mediocrity in hereditary stature.  
*Journal of the Anthropological Institute* vol. 15,. 246–263.

3.5

4.6

4.4

4.2

4

3.8

3.6

3.4

SURFACE TEMPERATURE

6.5

## REGRESSION MODEL

$$Y_i = X_i' \beta^0 + e_i$$

$$= X_{i1}\beta_1^0 + X_{i2}\beta_2^0 + \dots + X_{ip}\beta_p^0 + e_i,$$

$i = 1, 2, \dots, n$

$Y_i$

-

RESPONSE VARIABLE

(for  $i$ -th object, known)

Galton, F. (1886): Regression towards mediocrity in hereditary stature.  
*Journal of the Anthropological Institute* vol. 15,. 246–263.


3.5

4.6


4.4


THE TASK IS TO ESTIMATE UNKNOWN  
REGRESSION COEFFICIENTS

CONTROL TEMPERATURE




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion








A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

## The Least Squares


LUMINOUS  
OUTPUT



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

## Let's turn to economic data - investments in various industries and their profits


industry	investment	profit
1	9.39	165.2
2	1.22	9.4
3	0.62	-42.2
4	1.22	289.4
5	1.35	200.4
6	1.21	-179.0
7	1.35	160.4
8	0.07	49.0
9	0.03	-156.1
10	0.07	88.0
11	10.03	103.9
12	0.62	181.7
13	1.23	-93.7
14	1.23	-123.7
15	0.73	-294.4
16	1.10	-131.5
17	0.98	216.3
18	0.61	-165.7



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Let's turn to economic data - investments in various industries and their profits

industry	investment	profit
1	9.39	165.2
2	1.22	9.4
3	0.62	-42.2
4	1.22	289.4
5	1.35	200.4
6	1.21	-179.0
7	1.35	160.4
8	0.07	49.0
9	0.03	-156.1
10	0.07	88.0
11	10.03	103.9
12	1.23	-123.7
13	0.73	-294.4
14	1.10	-131.5
15	0.98	216.3
16	0.61	-165.7




*Are two leverage points suspicious ?*

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

## Let's turn to economic data


industry	investment	profit
1	9.39	165.2
2	1.22	9.4
3	0.62	-42.2
4	1.22	289.4
5	1.35	200.4
6	1.21	-179.0
7	1.35	160.4
8	0.07	49.0
9	0.03	-156.1
10	0.07	88.0
11	10.03	103.9
12	0.62	181.7
13	1.23	-93.7
14	1.23	-123.7
15	0.73	-294.4
16	1.10	-131.5
17	0.98	216.3
18	0.61	-165.7



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

## Let's turn to economic data - data without leverage points

industry	investment	profit
2	1.22	9.4
3	0.62	-42.2
4	1.22	289.4
5	1.35	200.4
6	1.21	-179.0
7	1.35	160.4
8	0.07	49.0
9	0.03	-156.1
10	0.07	88.0
12	0.62	181.7
13	1.23	-93.7
14	1.23	-123.7
15	0.73	-294.4
16	1.10	-131.5
17	0.98	216.3
18	0.61	-165.7




## Let's turn to economic data - data without leverage points

industry	investment	profit
6	1.21	-179.0
7	1.35	160.4
8	0.07	49.0
9	0.03	-156.1
10	0.07	88.0
12	0.62	181.7
13	1.23	-93.7
14	1.23	-123.7
15	0.73	-294.4
16	1.10	-131.5
17	0.98	216.3
18	0.61	-165.7

A philosophical question<sub>(put by statistical fundamentalist)</sub>:

Who gave us a justification to delete some observation(s) ?



## Let's turn to economic data - data without leverage points

industry	investment	profit
----------	------------	--------

A philosophical question *(put by statistical fundamentalist)*:

Who gave us a justification to delete some observation(s) ?

6	1.21	-179.0
7	1.35	160.4

100+

The question can be inverted *(by a critical preview)*:

Who can force us to employ all observations  
when some of them are (evidently) wrong ?

13	1.23	-93.7
14	1.23	-123.7
15	0.73	-294.4
16	1.10	-131.5
17	0.98	216.3
18	0.61	-165.7

100+

-200


-300

0 0.2 0.4 0.6 0.8 1 1.2 1.4




*The method of the least squares is seen to be our best course when we have thrown overboard a certain portion of our data - a sort of sacrifice which has often to be made by those who sail the stormy seas of Probability.*

*Francis Ysidro Edgeworth (1887)*



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

industry	investment	profit
2	1.22	9.4
3	0.62	-42.2
4	1.22	289.4
5	1.35	200.4
6	1.21	-179.0
7	1.35	160.4
8	0.07	49.0
9	0.03	-156.1
10	0.07	88.0
12	0.62	181.7
13	1.23	-93.7
14	1.23	-123.7
15	0.73	-294.4
16	1.10	-131.5
17	0.98	216.3
18	0.61	-165.7




response variable = profit, explanatory variable = investment

$$y_i = \beta_0 + \beta_1 \cdot x_i + u_i \quad i = \text{industry} = 2, 3, \dots, 10, 12, \dots, 18$$

## Graphical analysis

Drawing the data on the screen can help a lot  
- but it has one, very significant restriction (limitation).



## Graphical analysis

Drawing the data on the screen can help a lot  
- but it has one, very significant restriction (limitation).

*Could You guess which one it is ?*

## Graphical analysis

Drawing the data on the screen can help a lot  
- but it has one, very significant restriction (limitation).

*Could You guess which one it is ?*

If no idea,

THE ANSWER WILL BE CLEAR AFTER TRYING TO EMPLOY IT.

ANALYSIS OF THE EXPORT FROM THE CZECH REPUBLIC TO EU IN 1994  
BY MEANS OF THE *Least Trimmed Squares*

*has found:*

MAIN SUBGROUP

*with number of industries 54 and the model*

$$\frac{X_\ell}{S_\ell} = 4.64 - 0.032 \cdot \frac{US_\ell}{VA_\ell} - 0.022 \cdot \frac{HS_\ell}{VA_\ell} - 0.124 \cdot \frac{K_\ell}{VA_\ell} + 1.035 \cdot CR_\ell \\ - 3.199 \cdot TFPW_\ell + 1.048 \cdot BAL_\ell + 0.452 \cdot DP_\ell + \varepsilon_\ell$$

- $X_\ell$  - export from  $i$ -th industry,
- $US_\ell$  - number of university-passed employees in the  $i$ -th industry,
- $HS_\ell$  - number of high school-passed employees in the  $i$ -th industry,
- $VA_\ell$  - value added in the  $i$ -th industry,
- $K_\ell$  - capital in the  $i$ -th industry,
- $CR_\ell$  - percentage of market occupied by 3 largest producers,
- $TFPW_\ell$  - by wages normed productivity in the  $i$ -th industry,
- $Bal_\ell$  - Balasa index in the  $i$ -th industry,
- $DP_\ell$  - cost discontinuity in 1993 in the  $i$ -th industry

with coefficient of determination 0.97 and stable submodels

HOW IS IT WITH THE INFLUENCE OF THE INDIVIDUAL EXPLANATORY VAR?  
POSITIVE SIGN  $\implies$  POSITIVE INFLUENCE?

MAIN SUBGROUP


*with number of industries 54 and the model*

$$\frac{X_\ell}{S_\ell} = 4.64 - 0.032 \cdot \frac{US_\ell}{VA_\ell} - 0.022 \cdot \frac{HS_\ell}{VA_\ell} - 0.124 \cdot \frac{K_\ell}{VA_\ell} + 1.035 \cdot CR_\ell \\ - 3.199 \cdot TFPW_\ell + 1.048 \cdot BAL_\ell + 0.452 \cdot DP_\ell + \varepsilon_\ell$$

- $X_\ell$  - export from  $i$ -th industry,
- $US_\ell$  - number of university-passed employees in the  $i$ -th industry,
- $HS_\ell$  - number of high school-passed employees in the  $i$ -th industry,
- $VA_\ell$  - value added in the  $i$ -th industry,
- $K_\ell$  - capital in the  $i$ -th industry,
- $CR_\ell$  - percentage of market occupied by 3 largest producers,
- $TFPW_\ell$  - by wages normed productivity in the  $i$ -th industry,
- $BAL_\ell$  - Balasa index in the  $i$ -th industry,
- $DP_\ell$  - cost discontinuity in 1993 in the  $i$ -th industry

with coefficient of determination 0.97 and stable submodels

# Content

- 
- 1 A very first insight into robustness
    - An attractive application as foreword
  - 2 A bit more modest motivation
  - 3 Let's start more serious discussion
    - Vulnerability of classical procedures to contamination
    - The classical requirements on estimators



# Content

- 
- 1 A very first insight into robustness
    - An attractive application as foreword
  - 2 A bit more modest motivation
  - 3 Let's start more serious discussion
    - Vulnerability of classical procedures to contamination
    - The classical requirements on estimators

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Recalling the classical approach to point estimation

Maximum likelihood - solving an extremal problem

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta)$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Recalling the classical approach to point estimation

Maximum likelihood - solving an extremal problem

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta)$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \log \left\{ \prod_{i=1}^n f(x_i, \theta) \right\}$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Recalling the classical approach to point estimation

Maximum likelihood - solving an extremal problem

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta)$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \log \left\{ \prod_{i=1}^n f(x_i, \theta) \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log (f(x_i, \theta))$$

## Recalling the classical approach to point estimation

Maximum likelihood - solving an extremal problem

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta)$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \log \left\{ \prod_{i=1}^n f(x_i, \theta) \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log (f(x_i, \theta))$$

$$\hat{\theta}^{(ML,n)} = \arg \left\{ \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0 \right\}$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Recalling the classical approach to point estimation

Let e.g.  $f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(x-\mu)^2}{2\sigma^2} \right\}$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$



## Recalling the classical approach to point estimation

$$\text{Let e.g. } f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(x-\mu)^2}{2\sigma^2} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

$$\theta = (\mu, \sigma)$$

$$\frac{\partial f(x_i, \theta)}{\partial \mu} = 2 \cdot f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{2\sigma^2} \quad \text{and} \quad \frac{\partial f(x_i, \theta)}{\partial \sigma} = -f(x_i, \mu, \sigma^2) \left\{ \frac{1}{\sigma} - \frac{(x_i - \mu)^2}{\sigma^3} \right\}$$

## Recalling the classical approach to point estimation

$$\text{Let e.g. } f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(x-\mu)^2}{2\sigma^2} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

$$\theta = (\mu, \sigma)$$

$$\frac{\partial f(x_i, \theta)}{\partial \mu} = 2 \cdot f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{2\sigma^2} \quad \text{and} \quad \frac{\partial f(x_i, \theta)}{\partial \sigma} = -f(x_i, \mu, \sigma^2) \left\{ \frac{1}{\sigma} - \frac{(x_i - \mu)^2}{\sigma^3} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \left\{ \sum_{i=1}^n \frac{(x_i - \mu)}{2\sigma^2} = 0 \quad \text{and} \quad \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^3} = \frac{n}{\sigma} \right\}$$



## Recalling the classical approach to point estimation

$$\text{Let e.g. } f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(x-\mu)^2}{2\sigma^2} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

$$\theta = (\mu, \sigma)$$

$$\frac{\partial f(x_i, \theta)}{\partial \mu} = 2 \cdot f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{2\sigma^2} \quad \text{and} \quad \frac{\partial f(x_i, \theta)}{\partial \sigma} = -f(x_i, \mu, \sigma^2) \left\{ \frac{1}{\sigma} - \frac{(x_i - \mu)^2}{\sigma^3} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \left\{ \sum_{i=1}^n \frac{(x_i - \mu)}{2\sigma^2} = 0 \quad \text{and} \quad \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^3} = \frac{n}{\sigma} \right\}$$

$$\hat{\theta}^{(ML,n)} = \left( \hat{\mu}^{(ML,n)} = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad \hat{\sigma}^{(ML,n)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \hat{\mu}^{(ML,n)})^2} \right)$$

## Recalling the classical approach to point estimation

$$\text{Let e.g. } f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(x-\mu)^2}{2\sigma^2} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

$$\theta = (\mu, \sigma)$$

$$\frac{\partial f(x_i, \theta)}{\partial \mu} = 2 \cdot f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{2\sigma^2} \quad \text{and} \quad \frac{\partial f(x_i, \theta)}{\partial \sigma} = -f(x_i, \mu, \sigma^2) \left\{ \frac{1}{\sigma} - \frac{(x_i - \mu)^2}{\sigma^3} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \left\{ \sum_{i=1}^n \frac{(x_i - \mu)}{2\sigma^2} = 0 \quad \text{and} \quad \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^3} = \frac{n}{\sigma} \right\}$$

$$\hat{\theta}^{(ML,n)} = \left( \hat{\mu}^{(ML,n)} = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad \hat{\sigma}^{(ML,n)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \hat{\theta}^{(ML,n)})^2} \right)$$

$$\hat{\mu}^{(ML,n)} = \bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{unbiased, consistent}$$

## Recalling the classical approach to point estimation

$$\text{Let e.g. } f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(x-\mu)^2}{2\sigma^2} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \sum_{\theta \in \Theta}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

$$\theta = (\mu, \sigma)$$

$$\frac{\partial f(x_i, \theta)}{\partial \mu} = 2 \cdot f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{2\sigma^2} \quad \text{and} \quad \frac{\partial f(x_i, \theta)}{\partial \sigma} = -f(x_i, \mu, \sigma^2) \left\{ \frac{1}{\sigma} - \frac{(x_i - \mu)^2}{\sigma^3} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \sum_{\theta \in \Theta}^n \left\{ \sum_{i=1}^n \frac{(x_i - \mu)}{2\sigma^2} = 0 \quad \text{and} \quad \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^3} = \frac{n}{\sigma} \right\}$$

$$\hat{\theta}^{(ML,n)} = \left( \hat{\mu}^{(ML,n)} = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad \hat{\sigma}^{(ML,n)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \hat{\theta}^{(ML,n)})^2} \right)$$

$$\hat{\mu}^{(ML,n)} = \bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{unbiased, consistent}$$

$$\hat{\sigma}^{(ML,n)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2} \quad \text{biased, consistent}$$

## Recalling the classical approach to point estimation

$$\text{Let e.g. } f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(x-\mu)^2}{2\sigma^2} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \sum_{\theta \in \Theta}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

$$\theta = (\mu, \sigma)$$

$$\frac{\partial f(x_i, \theta)}{\partial \mu} = 2 \cdot f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{2\sigma^2} \quad \text{and} \quad \frac{\partial f(x_i, \theta)}{\partial \sigma} = -f(x_i, \mu, \sigma^2) \left\{ \frac{1}{\sigma} - \frac{(x_i - \mu)^2}{\sigma^3} \right\}$$

$$\hat{\theta}^{(ML,n)} = \arg \sum_{\theta \in \Theta}^n \left\{ \sum_{i=1}^n \frac{(x_i - \mu)}{2\sigma^2} = 0 \quad \text{and} \quad \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^3} = \frac{n}{\sigma} \right\}$$

$$\hat{\theta}^{(ML,n)} = \left( \hat{\mu}^{(ML,n)} = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad \hat{\sigma}^{(ML,n)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \hat{\theta}^{(ML,n)})^2} \right)$$

$$\hat{\mu}^{(ML,n)} = \bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{unbiased, consistent}$$

$$\hat{\sigma}^{(ML,n)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2} \quad \text{biased, consistent}$$

$$\rightarrow s_n^2 = \frac{1}{(n-1)} \sum_{i=1}^n (x_i - \bar{x}_n)^2 \quad \text{unbiased, consistent}$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

What we have observed on the previous slide ?

Typical features of the classical estimators




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## What we have observed on the previous slide ?

Typical features of the classical estimators

Let's consider only estimators which are as  $\hat{\mu}^{(ML,n)}$ , then:



## What we have observed on the previous slide ?

### Typical features of the classical estimators

Let's consider only estimators which are as  $\hat{\mu}^{(ML,n)}$ , then:

Pros:

- 1 The estimators are defined as solution of extremal problem.

## What we have observed on the previous slide ?

### Typical features of the classical estimators

Let's consider only estimators which are as  $\hat{\mu}^{(ML,n)}$ , then:

#### Pros:

- 1 The estimators are defined as solution of extremal problem.
- 2 The extremal problem is (usually) invertible,  
i. e. we have a formula for the estimator,  
hence we can (more or less) easy implement it.



## What we have observed on the previous slide ?

### Typical features of the classical estimators

Let's consider only estimators which are as  $\hat{\mu}^{(ML,n)}$ , then:

#### Pros:

- 1 The estimators are defined as solution of extremal problem.
- 2 The extremal problem is (usually) invertible,  
i. e. we have a formula for the estimator,  
hence we can (more or less) easy implement it.
- 3 They are (mostly) unbiased, consistent, asymptotically normal, etc.

## What we have observed on the previous slide ?

### Typical features of the classical estimators

Let's consider only estimators which are as  $\hat{\mu}^{(ML,n)}$ , then:

#### Pros:

- 1 The estimators are defined as solution of extremal problem.
- 2 The extremal problem is (usually) invertible,  
i. e. we have a formula for the estimator,  
hence we can (more or less) easy implement it.
- 3 They are (mostly) unbiased, consistent, asymptotically normal, etc.
- 4 If exponential family, usually efficient.

## What we have observed on the previous slide ?

### Typical features of the classical estimators

Let's consider only estimators which are as  $\hat{\mu}^{(ML,n)}$ , then:

Pros:

- 1 The estimators are defined as solution of extremal problem.
- 2 The extremal problem is (usually) invertible,  
i. e. we have a formula for the estimator,  
hence we can (more or less) easy implement it.
- 3 They are (mostly) unbiased, consistent, asymptotically normal, etc.
- 4 If exponential family, usually efficient.

## What we have observed on the previous slide ?

### Typical features of the classical estimators

Let's consider only estimators which are as  $\hat{\mu}^{(ML,n)}$ , then:

Pros:

- 1 The estimators are defined as solution of extremal problem.
- 2 The extremal problem is (usually) invertible,  
i. e. we have a formula for the estimator,  
hence we can (more or less) easy implement it.
- 3 They are (mostly) unbiased, consistent, asymptotically normal, etc.
- 4 If exponential family, usually efficient.


Cons: ??? see the next slide !!!

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice the location of mean

The data generated as standard normal, mean denoted by ●.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice the location of mean

The data generated as standard normal, mean denoted by ●.



Contamination at 1.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice the location of mean

The data generated as standard normal, mean denoted by ●.



Contamination at 2.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice the location of mean

The data generated as standard normal, mean denoted by ●.



Contamination at 3.






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice the location of mean

The data generated as standard normal, mean denoted by ●.



Contamination at 4.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice the location of mean

The data generated as standard normal, mean denoted by ●.



Contamination at 8.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion


Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice the location of mean

The data generated as standard normal, mean denoted by ●.




Contamination at 12.




## Notice the location of mean

The data generated as standard normal, mean denoted by ●.



Contamination at 12.



Conclusion - the classical estimators are (frequently)  
vulnerable to contamination.

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion


Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Let's study general reasons causing it - returning a few slides back.

Maximum likelihood - solving an extremal problem

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta)$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log(f(x_i, \theta))$$



Let's study general reasons causing it - returning a few slides back.

Maximum likelihood - solving an extremal problem

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta)$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log(f(x_i, \theta))$$

Let again  $f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\}$  and consider only  $\mu$

$$\Rightarrow \hat{\mu}^{(ML,n)} = \arg \min_{\mu \in R} \left\{ \sum_{i=1}^n (x_i - \mu)^2 \right\}$$

Let's study general reasons causing it - returning a few slides back.

Maximum likelihood - solving an extremal problem

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta)$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log(f(x_i, \theta))$$

Let again  $f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\}$  and consider only  $\mu$

$$\Rightarrow \hat{\mu}^{(ML,n)} = \arg \min_{\mu \in R} \left\{ \sum_{i=1}^n (x_i - \mu)^2 \right\}$$


The observations with large  $(x_i - \mu)^2$   
have a large influence on solution.

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Evidently, low robustness is consequence of quadratic objective function

We have such objective function.






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Evidently, low robustness is consequence of quadratic objective function

We should depress influence of large residuals.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Evidently, low robustness is consequence of quadratic objective function

We should employ such **objective function**.



## Evidently, low robustness is consequence of quadratic objective function




Conclusion - instead of solving

$$\hat{\mu}^{(ML,n)} = \arg \min_{\mu \in R} \sum_{i=1}^n (x_i - \mu)^2$$

we should solve

$$\hat{\mu}^{(ML,n)} = \arg \min_{\mu \in R} \sum_{i=1}^n \rho(x_i - \mu).$$



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Let's study general reasons causing it - an alternative way.

Maximum likelihood - solving the normal equations

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta) = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log(f(x_i, \theta))$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

## Let's study general reasons causing it - an alternative way.

### Maximum likelihood - solving the normal equations

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta) = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log(f(x_i, \theta))$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

Let again  $f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\}$ , i. e.  $\frac{\partial f(x_i, \theta)}{\partial \mu} = f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{\sigma^2}$

and consider only  $\mu \Rightarrow \hat{\mu}^{(ML,n)} = \arg \max_{\mu \in R} \left\{ \sum_{i=1}^n (x_i - \mu) = 0 \right\}$

## Let's study general reasons causing it - an alternative way.

### Maximum likelihood - solving the normal equations

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \prod_{i=1}^n f(x_i, \theta) = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \log(f(x_i, \theta))$$

$$\hat{\theta}^{(ML,n)} = \arg \max_{\theta \in \Theta} \sum_{i=1}^n \frac{1}{f(x_i, \theta)} \cdot \frac{\partial f(x_i, \theta)}{\partial \theta} = 0$$

Let again  $f(x, \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\}$ , i.e.  $\frac{\partial f(x_i, \theta)}{\partial \mu} = f(x_i, \mu, \sigma^2) \cdot \frac{(x_i - \mu)}{\sigma^2}$

and consider only  $\mu \Rightarrow \hat{\mu}^{(ML,n)} = \arg \max_{\mu \in R} \left\{ \sum_{i=1}^n (x_i - \mu) = 0 \right\}$

The same conclusion:


The observations with large  $|x_i - \mu|$   
have a large influence on solution.

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Equivalently, low robustness is consequence of identity in normal equations

We have such influence function.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Equivalently, low robustness is consequence of identity in normal equations

We should depress influence of large residuals.






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Equivalently, low robustness is consequence of identity in normal equations

We should employ such **influence function**.




## Equivalently, low robustness is consequence of identity in normal equations

Conclusion - instead of solving


$$\sum_{i=1}^n (x_i - \mu) = 0$$

we should solve

$$\sum_{i=1}^n \psi(x_i - \mu) = 0.$$



# Content

- 
- 1 A very first insight into robustness
    - An attractive application as foreword
  - 2 A bit more modest motivation
  - 3 **Let's start more serious discussion**
    - Vulnerability of classical procedures to contamination
    - **The classical requirements on estimators**

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Recalling the classical requirements on estimators

### 1 Unbiasedness



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion


Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Recalling the classical requirements on estimators


- 1 Unbiasedness
- 2 Consistency (weak, strong)




## Recalling the classical requirements on estimators

- 
- 1 Unbiasedness
  - 2 Consistency (weak, strong)
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)

## Recalling the classical requirements on estimators


- 
- 1 Unbiasedness
  - 2 Consistency (weak, strong)
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)
  - 4 Asymptotic normality

## Recalling the classical requirements on estimators


- 
- 1 Unbiasedness
  - 2 Consistency (weak, strong)
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)
  - 4 Asymptotic normality
  - 5 Efficiency




## Recalling the classical requirements on estimators

- 
- 1 Unbiasedness
  - 2 Consistency (weak, strong)
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)
  - 4 Asymptotic normality
  - 5 Efficiency
  - 6 Scale- and regression-equivariance


## Recalling the classical requirements on estimators

- 
- 1 Unbiasedness
  - 2 Consistency (weak, strong)
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)
  - 4 Asymptotic normality
  - 5 Efficiency
  - 6 Scale- and regression-equivariance
  - 7 **Admissibility**

## Recalling the classical requirements on estimators

- 
- 1 Unbiasedness
  - 2 Consistency (weak, strong)
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)
  - 4 Asymptotic normality
  - 5 Efficiency
  - 6 Scale- and regression-equivariance
  - 7 Admissibility


## Recalling the classical requirements on estimators

- 
- 1 Unbiasedness
  - 2 Consistency (weak, strong)
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)
  - 4 Let's discuss them one by one.
  - 5 Efficiency
  - 6 Scale- and regression-equivariance
  - 7 Admissibility

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Unbiasedness

$$E_{\theta} \left[ \hat{\theta}^n(x_1, x_2, \dots, x_n) \right] = \int_{\mathcal{X}^n} \hat{\theta}^n(x_1, x_2, \dots, x_n) f_{\theta}(x_1, x_2, \dots, x_n) dx_1 \cdot dx_2 \cdot \dots \cdot dx_n = \theta$$


A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Unbiasedness

$$E_{\theta} \left[ \hat{\theta}^n(x_1, x_2, \dots, x_n) \right] = \int_{\mathcal{X}^n} \hat{\theta}^n(x_1, x_2, \dots, x_n) f_{\theta}(x_1, x_2, \dots, x_n) dx_1 \cdot dx_2 \cdot \dots \cdot dx_n = \theta$$

Is this requirement justifiable everytime ?

## Unbiasedness

$$E_{\theta} \left[ \hat{\theta}^n(x_1, x_2, \dots, x_n) \right] = \int_{\mathcal{X}^n} \hat{\theta}^n(x_1, x_2, \dots, x_n) f_{\theta}(x_1, x_2, \dots, x_n) dx_1 \cdot dx_2 \cdot \dots \cdot dx_n = \theta$$

Hoerl, A. E., R. W. Kennard (1970): Ridge regression:

Biased estimation for nonorthogonal problems.

*Technometrics* 12, 55 - 68.


$$\hat{\beta}^{(R,n)} = (X'X + \delta \cdot I)^{-1} X'Y$$

## Possible density of unbiased and biased estimator



Now we are going to discuss the following situation:

Unbiased estimator has slowly (if any) decreasing variance,  
while the variance and the bias of other (green) estimator decrease rapidly.






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


## Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Notice decreasing variance and bias






A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators


Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Notice decreasing variance and bias



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Consistency


1 (Weak) consistency - convergence in probability



## Consistency

- 
- 1 (Weak) consistency - convergence in probability
  - 2 Strong consistency - convergence almost surely

## Consistency


- 
- 1 (Weak) consistency - convergence in probability
  - 2 Strong consistency - convergence almost surely
  - 3  $\sqrt{n}$ -consistency (root-n-consistency)

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Convergence in probability (weak convergence)

Let  $X$  and  $\{X_n\}_{n=1}^{\infty}$  be random variable (r. v.) and a sequence of r.v.'s, respectively.



## Convergence in probability (weak convergence)

Let  $X$  and  $\{X_n\}_{n=1}^{\infty}$  be random variable (r. v.) and a sequence of r.v.'s, respectively.

We say that the sequence

$\{X_n\}_{n=1}^{\infty}$  converge in probability (weakly) to  $X$

if:

$$\forall (\varepsilon > 0, \delta > 0) \quad \exists (n_{\varepsilon, \delta} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, \delta})$$

$$P(\{\omega \in \Omega : |X_n(\omega) - X(\omega)| > \delta\}) < \varepsilon$$



## Convergence in probability (weak convergence)

Let  $X$  and  $\{X_n\}_{n=1}^{\infty}$  be random variable (r. v.) and a sequence of r.v.'s, respectively.

We say that the sequence

$\{X_n\}_{n=1}^{\infty}$  converge in probability (weakly) to  $X$

if:

$$\forall (\varepsilon > 0, \delta > 0) \quad \exists (n_{\varepsilon, \delta} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, \delta})$$

$$P(\{\omega \in \Omega : |X_n(\omega) - X(\omega)| > \delta\}) < \varepsilon$$

or alternatively

$$P(\{\omega \in \Omega : |X_n(\omega) - X(\omega)| < \delta\}) > 1 - \varepsilon.$$

## (Weak) consistency - convergence in probability

In the case that we speak about an estimator of “true”  $\beta^0$ ,  
we say that  $\hat{\beta}^{(method,n)}$  is (weakly) consistent if:

$$\forall (\varepsilon > 0, \delta > 0) \quad \exists (n_{\varepsilon, \delta} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, \delta}) \\ P \left( \left\{ \omega \in \Omega : \left\| \hat{\beta}^{(method,n)} - \beta^0 \right\| > \delta \right\} \right) < \varepsilon$$

## (Weak) consistency - convergence in probability

In the case that we speak about an estimator of “true”  $\beta^0$ ,  
we say that  $\hat{\beta}^{(method,n)}$  is (weakly) consistent if:

$$\forall (\varepsilon > 0, \delta > 0) \quad \exists (n_{\varepsilon, \delta} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, \delta})$$

$$P \left( \left\{ \omega \in \Omega : \left\| \hat{\beta}^{(method,n)} - \beta^0 \right\| > \delta \right\} \right) < \varepsilon$$

or alternatively


$$P \left( \left\{ \omega \in \Omega : \left\| \hat{\beta}^{(method,n)} - \beta^0 \right\| < \delta \right\} \right) > 1 - \varepsilon.$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Convergence almost surely (strong convergence)

Let  $X$  and  $\{X_n\}_{n=1}^{\infty}$  be r.v. and a sequence of r.v.'s, respectively.



## Convergence almost surely (strong convergence)

Let  $X$  and  $\{X_n\}_{n=1}^{\infty}$  be r.v. and a sequence of r.v.'s, respectively.  
We say that the sequence

$\{X_n\}_{n=1}^{\infty}$  converges almost surely (strongly) to  $X$

if:

$$\exists (A \in \mathcal{A}, P(A) = 1) \quad \forall (\varepsilon > 0, \omega_0 \in A) \quad \exists (n_{\varepsilon, \omega_0} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, \omega_0}) \\ |X_n(\omega_0) - X(\omega_0)| < \varepsilon.$$

## Strong consistency - convergence almost surely

In the case that we speak about an estimator of “true”  $\beta^0$ ,  
we say that  $\hat{\beta}^{(method,n)}$  is **strongly consistent** if:


$$\exists (A \in \mathcal{A}, P(A) = 1) \quad \forall (\varepsilon > 0, \omega_0 \in A) \quad \exists (n_{\varepsilon, \omega_0} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, \omega_0}) \\ \left\| \hat{\beta}^{(method,n)}(\omega_0) - \beta^0 \right\| < \varepsilon.$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## $\sqrt{n}$ -consistency (root of $n$ consistency)

In this case we typically speak about an estimator of “true”  $\beta^0$ .



## $\sqrt{n}$ -consistency (root of $n$ consistency)

In this case we typically speak about an estimator of “true”  $\beta^0$ .

Then we say that  $\hat{\beta}^{(method,n)}$  is  $\sqrt{n}$ -consistent if:

$$\forall (\varepsilon > 0) \quad \exists (K_\varepsilon < \infty \text{ and } n_{\varepsilon, K_\varepsilon} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, K_\varepsilon})$$

$$P \left( \left\{ \omega \in \Omega : \sqrt{n} \left\| \hat{\beta}^{(method,n)} - \beta^0 \right\| > K_\varepsilon \right\} \right) < \varepsilon.$$



## $\sqrt{n}$ -consistency (root of $n$ consistency)

In this case we typically speak about an estimator of “true”  $\beta^0$ .

Then we say that  $\hat{\beta}^{(method,n)}$  is  $\sqrt{n}$ -consistent if:

$$\forall (\varepsilon > 0) \quad \exists (K_\varepsilon < \infty \text{ and } n_{\varepsilon, K_\varepsilon} \in \mathcal{N}) \quad \forall (n \geq n_{\varepsilon, K_\varepsilon})$$

$$P \left( \left\{ \omega \in \Omega : \sqrt{n} \left\| \hat{\beta}^{(method,n)} - \beta^0 \right\| > K_\varepsilon \right\} \right) < \varepsilon.$$

or alternatively


$$P \left( \left\{ \omega \in \Omega : \sqrt{n} \left\| \hat{\beta}^{(method,n)} - \beta^0 \right\| \leq K_\varepsilon \right\} \right) > 1 - \varepsilon.$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Asymptotic normality

It is again case when we speak about an estimator of “true”  $\beta^0$ .



## Asymptotic normality

It is again case when we speak about an estimator of “true”  $\beta^0$ .

Then we say that  $\hat{\beta}^{(method,n)}$  is asymptotically normal if :

$$\mathcal{L} \left( n^a \left( \hat{\beta}^{(method,n)} - \beta^0 \right) \right) \rightarrow \mathcal{N}(0, \Sigma)$$

where  $a \leq \frac{1}{2}$  and usually reaches the upper bound, i. e. usually  $a = \frac{1}{2}$ .

## Asymptotic normality

It is again case when we speak about an estimator of “true”  $\beta^0$ .

Then we say that  $\hat{\beta}^{(method,n)}$  is asymptotically normal if :

$$\mathcal{L} \left( n^a \left( \hat{\beta}^{(method,n)} - \beta^0 \right) \right) \rightarrow \mathcal{N}(0, \Sigma)$$

where  $a \leq \frac{1}{2}$  and usually reaches the upper bound, i. e. usually  $a = \frac{1}{2}$ .

Asymptotic normality is (was? - in the case of OLS, ML, etc.) employed:

- 1 for constructing (asymptotic) confidence interval  
and

## Asymptotic normality

It is again case when we speak about an estimator of “true”  $\beta^0$ .

Then we say that  $\hat{\beta}^{(method,n)}$  is asymptotically normal if :

$$\mathcal{L} \left( n^a \left( \hat{\beta}^{(method,n)} - \beta^0 \right) \right) \rightarrow \mathcal{N}(0, \Sigma)$$

where  $a \leq \frac{1}{2}$  and usually reaches the upper bound, i. e. usually  $a = \frac{1}{2}$ .

Asymptotic normality is (was? - in the case of OLS, ML, etc.) employed:

- 1 for constructing (asymptotic) confidence interval  
and
- 2 for verification of  $\sqrt{n}$ -consistency.

## Efficiency

We usually say that  $\hat{\beta}^{(method,n)}$  is (asymptotically) efficient, if its covariance matrix reaches (asymptotically) lower Rao-Cramer bound in the sense of ordering the matrices by positive semidefiniteness.

## Efficiency

We usually say that  $\hat{\beta}^{(method,n)}$  is (asymptotically) efficient, if its covariance matrix reaches (asymptotically) lower Rao-Cramer bound in the sense of ordering the matrices by positive semidefiniteness.

Sometimes, we say that  $\hat{\beta}^{(method,n)}$  is (asymptotically) efficient, if its covariance matrix reaches (asymptotically) the minimal possible value in given family of estimators - again in the sense of ordering the matrices by positive semidefiniteness.

## Efficiency

Efficiency is:

- 1 important notion from the pedagogical point view,





## Efficiency

Efficiency is:

- 1 important notion from the pedagogical point view,
- 2 important from abstract theoretical background of statistics
  - how much we could reach if data would be “clear”,

## Efficiency

Efficiency is:

- 1 important notion from the pedagogical point view,
- 2 important from abstract theoretical background of statistics
  - how much we could reach if data would be “clear”,
- 3 it can be destroyed by a small deviation  
from the exponential family - Huber's example

## Efficiency

Efficiency is:

- 1 important notion from the pedagogical point view,
- 2 important from abstract theoretical background of statistics
  - how much we could reach if data would be “clear”,
- 3 it can be destroyed by a small deviation
  - from the exponential family - Huber's example

## Efficiency

Efficiency is:

- 1 important notion from the pedagogical point view,
  - 2 important from abstract theoretical background of statistics
    - how much we could reach if data would be “clear”,
  - 3 it can be destroyed by a small deviation
    - from the exponential family - Huber's example
- and
- 4 need not imply too much - Fisher's example.

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Small deviation from exact model can cause ...

Huber, P. J. (1980): *Robust Statistics*.

New York: J.Wiley and Sons.

$$s_n = \left[ \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2 \right]^{\frac{1}{2}}$$

## Small deviation from exact model can cause ...

Huber, P. J. (1980): *Robust Statistics*.

New York: J.Wiley and Sons.

$$s_n = \left[ \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2 \right]^{\frac{1}{2}} \quad d_n = \frac{\pi}{2n} \sum_{i=1}^n |x_i - \bar{x}_n|$$

## Small deviation from exact model can cause ...

Huber, P. J. (1980): *Robust Statistics*.

New York: J.Wiley and Sons.

$$s_n = \left[ \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2 \right]^{\frac{1}{2}} \quad d_n = \frac{\pi}{2n} \sum_{i=1}^n |x_i - \bar{x}_n|$$

$$F(x) = (1 - \varepsilon)\Phi(x) + \varepsilon\Phi\left(\frac{x}{3}\right)$$

## Small deviation from exact model can cause ...

Huber, P. J. (1980): *Robust Statistics*.

New York: J.Wiley and Sons.

$$s_n = \left[ \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2 \right]^{\frac{1}{2}} \quad d_n = \frac{\pi}{2n} \sum_{i=1}^n |x_i - \bar{x}_n|$$

$$F(x) = (1 - \varepsilon)\Phi(x) + \varepsilon\Phi\left(\frac{x}{3}\right)$$


$$ARE_F(\varepsilon) = \lim_{n \rightarrow \infty} \frac{\text{var}_F s_n / E_F^2 s_n}{\text{var}_F d_n / E_F^2 d_n}$$



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion


Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Small deviation from exact model can cause ...



$\varepsilon$	0	0.001	0.002	0.05
$ARE(\varepsilon)$	0.876	0.948	1.016	2.035

## Small deviation from exact model can cause ...




$\varepsilon$	0	0.001	0.002	0.05
$ARE(\varepsilon)$	0.876	0.948	1.016	2.035

So, 5% of contamination  $\rightarrow d_n$  is two times better than  $s_n$ .

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Is 5% contamination too much or too little?



Hampel, F. R., E. M. Ronchetti, P. J. Rousseeuw, W. A. Stahel. (1986):  
*Robust Statistic - The Approach Based on Influence Curve.*  
New York: J.Wiley and Sons.

E. g. Switzerland has 6% of errors in mortality tables.

## Is the efficiency really important or a bit misleading?

Fisher, R. A. (1922): On the mathematical foundation of theoretical statistics.

*Philos. Trans. Roy. Soc. London Ser. A* 222, 309 - 368.

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(\bar{X}_n)}{\text{var}_{t(\nu)}(\bar{X}_n)} = 1 - \frac{6}{\nu(\nu+1)}$$

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(s_n^2)}{\text{var}_{t(\nu)}(s_n^2)} = 1 - \frac{12}{\nu(\nu+1)}$$

## Is the efficiency really important or a bit misleading?

Fisher, R. A. (1922): On the mathematical foundation of theoretical statistics.

*Philos. Trans. Roy. Soc. London Ser. A* 222, 309 - 368.

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(\bar{X}_n)}{\text{var}_{t(\nu)}(\bar{X}_n)} = 1 - \frac{6}{\nu(\nu+1)}$$

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(S_n^2)}{\text{var}_{t(\nu)}(S_n^2)} = 1 - \frac{12}{\nu(\nu+1)}$$

## Is the efficiency really important or a bit misleading?

Fisher, R. A. (1922): On the mathematical foundation of theoretical statistics.

*Philos. Trans. Roy. Soc. London Ser. A* 222, 309 - 368.

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(\bar{X}_n)}{\text{var}_{t(\nu)}(\bar{X}_n)} = 1 - \frac{6}{\nu(\nu+1)}$$

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(s_n^2)}{\text{var}_{t(\nu)}(s_n^2)} = 1 - \frac{12}{\nu(\nu+1)}$$

## Is the efficiency really important or a bit misleading?

Fisher, R. A. (1922): On the mathematical foundation of theoretical statistics.

*Philos. Trans. Roy. Soc. London Ser. A* 222, 309 - 368.

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(\bar{X}_n)}{\text{var}_{t(\nu)}(\bar{X}_n)} = 1 - \frac{6}{\nu(\nu+1)}$$
$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(S_n^2)}{\text{var}_{t(\nu)}(S_n^2)} = 1 - \frac{12}{\nu(\nu+1)}$$

## Is the efficiency really important or a bit misleading?

Fisher, R. A. (1922): On the mathematical foundation of theoretical statistics.

*Philos. Trans. Roy. Soc. London Ser. A* 222, 309 - 368.

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(\bar{X}_n)}{\text{var}_{t(\nu)}(\bar{X}_n)} = 1 - \frac{6}{\nu(\nu+1)}$$

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(s_n^2)}{\text{var}_{t(\nu)}(s_n^2)} = 1 - \frac{12}{\nu(\nu+1)}$$



## Is the efficiency really important or a bit misleading?

Fisher, R. A. (1922): On the mathematical foundation of theoretical statistics.

*Philos. Trans. Roy. Soc. London Ser. A* 222, 309 - 368.


$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(\bar{X}_n)}{\text{var}_{t(\nu)}(\bar{X}_n)} = 1 - \frac{6}{\nu(\nu+1)}$$

$$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(S_n^2)}{\text{var}_{t(\nu)}(S_n^2)} = 1 - \frac{12}{\nu(\nu+1)}$$

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## Is the efficiency really important or a bit misleading?




$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(T_n)}{\text{var}_{t(\nu)}(T_n)}$	$t_9$	$t_5$	$t_3$
$\bar{X}_n$	0.93	0.80	0.50
$S_n^2$	0.83	0.40	0!

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

Is the efficiency really important or a bit misleading?




$\lim_{n \rightarrow \infty} \frac{\text{var}_{N(0,1)}(T_n)}{\text{var}_{t(\nu)}(T_n)}$	$t_9$	$t_5$	$t_3$
$\bar{X}_n$	0.93	0.80	0.50
$S_n^2$	0.83	0.40	0!

A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## How far is Student density from the normal one ?

THE BLUE CURVE IS STANDARD NORMAL WHILE THE RED ONE IS THE STUDENT'S WITH 3 DEGREES OF FREEDOM.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## How far is Student density from the normal one ?

THE BLUE CURVE IS STANDARD NORMAL WHILE THE RED ONE IS THE  
STUDENT'S WITH 5 DEGREES OF FREEDOM.




A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

## How far is Student density from the normal one ?

THE BLUE CURVE IS STANDARD NORMAL WHILE THE RED ONE IS THE STUDENT'S WITH 9 DEGREES OF FREEDOM.



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators

The scale- and regression equivariance  
and the admissibility will be discussed later on.



A very first insight into robustness  
A bit more modest motivation  
Let's start more serious discussion

Vulnerability of classical procedures to contamination  
The classical requirements on estimators



*THANKS FOR ATTENTION*