

Loestring 23 Fe – Birth Control Pill

IMAGE OF PILL



Jane Smith
15/05/1989





WARNING

Morning 7am-9am	Noon 11am-1pm	Evening 7am-9am	Bedtime 7am-9am
1			

Take one tablet by mouth at the same time every day.
Tablets should not be skipped or taken at intervals exceeding 24 hours.

Drug Info

Loestrin 23 Fe contains female hormones that prevent ovulation (the release of an egg from an ovary). This medication also causes changes in your cervical mucus and uterine lining, making it harder for sperm to reach the uterus and harder for a fertilized egg to attach to the uterus.




If you meet any of these criteria Do Not Use this product.

- if you are pregnant or if you have recently had a baby.
- uncontrolled high blood pressure
- heart disease
- coronary artery disease
- circulation problems (especially with diabetes)
- undiagnosed vaginal bleeding
- liver disease or liver cancer
- severe migraine headaches
- if you also take certain hepatitis C medication
- if you have major surgery
- if you smoke and are over 35
- or if you have ever had a heart attack
- a stroke, a blood clot
- jaundice caused by pregnancy or birth control pills
- or cancer of the breast, uterus/cervix, or vagina.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. You should not take Loestrin 24 Fe if you smoke and are over 35 years old.

Razadyne – Alzheimer’s Medicine

IMAGE OF PILL



John Smith
21/11/1954

WARNING

Morning 7am-9am	Noon 11am-1pm	Evening 7am-9am	Bedtime 7am-9am
1		1	





Twice daily, one in the morning, one at night preferably with food.

Drug Info

Razadyne acts to improve communication between nerve cells, by increasing the levels of the essential neurotransmitter (a chemical messenger that transmits signals between nerve cells) acetylcholine, which is involved in learning, thinking, and memory processes.

The symptoms of AD are believed to be caused by the death of nerve cells in the brain. This loss can include nerve cells producing acetylcholine. As these nerve cells die, less acetylcholine is produced. This results in a disruption of these essential cognitive processes and can cause the symptoms of AD.



To make sure Razadyne is safe for you, tell your doctor if you have:

- urination problems;
- heart disease or a heart rhythm disorder;
- a history of stomach ulcer or bleeding;
- seizures or epilepsy;
- kidney disease;
- liver disease;
- Or a history of asthma or chronic obstructive pulmonary disease (COPD).

It is not known whether Razadyne will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

Common side effects may include:

- nausea, vomiting, diarrhea, loss of appetite;
- headache, dizziness; or
- weight loss.