

MATB44 - Week 5 Tutorial Teaching Notes

Week 5 Topics

Homogeneous Linear ODEs with Constant Coefficients

Definition and Theory

Linear Homogeneous ODE with Constant Coefficients

An n th-order linear homogeneous ODE with constant coefficients has the form:

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 \frac{dy}{dx} + a_0 y = 0, \quad a_n \neq 0,$$

where a_0, \dots, a_n are constants.

Principle of Superposition

- If y_1, \dots, y_k are solutions, then any linear combination $c_1 y_1 + \cdots + c_k y_k$ is also a solution.
- A fundamental solution set consists of n linearly independent solutions; the general solution is their linear combination.

Characteristic Equation

- Seek solutions of the form $y = e^{rx}$.
- Substitution gives the characteristic polynomial:

$$a_n r^n + a_{n-1} r^{n-1} + \cdots + a_1 r + a_0 = 0.$$

- Roots (including multiplicities) determine the shape of the general solution.

Root Cases and General Solution Forms

- Distinct real roots r_1, \dots, r_n :

$$y(x) = C_1 e^{r_1 x} + \cdots + C_n e^{r_n x}.$$

- Repeated real root r of multiplicity m :

$$y(x) = e^{rx} (C_0 + C_1 x + \cdots + C_{m-1} x^{m-1}).$$

- Complex conjugate pair $r = \alpha \pm i\beta$ ($\beta \neq 0$):

$$y(x) = e^{\alpha x} (C_1 \cos(\beta x) + C_2 \sin(\beta x)).$$

- Complex pair with multiplicity m :

$$y(x) = e^{\alpha x} \sum_{k=0}^{m-1} (A_k x^k \cos(\beta x) + B_k x^k \sin(\beta x)).$$

Key Examples

Example 1: Distinct Real Roots

Solve $y'' - 3y' + 2y = 0$ with $y(0) = 2$, $y'(0) = 0$.

Characteristic: $r^2 - 3r + 2 = 0 \Rightarrow r = 1, 2$. General solution: $y = C_1e^x + C_2e^{2x}$.

Apply IVP: - $y(0) = C_1 + C_2 = 2$ - $y'(x) = C_1e^x + 2C_2e^{2x} \Rightarrow y'(0) = C_1 + 2C_2 = 0$

Solve: from $C_1 + 2C_2 = 0$ we get $C_1 = -2C_2$; then $-2C_2 + C_2 = 2 \Rightarrow C_2 = -2$, $C_1 = 4$.

Final: $y(x) = 4e^x - 2e^{2x}$.

Example 2: Repeated Real Root

Solve $y'' - 4y' + 4y = 0$.

Characteristic: $(r - 2)^2 = 0$. General solution: $y(x) = (C_1 + C_2x)e^{2x}$.

Example 3: Complex Conjugate Roots

Solve $y'' + 4y' + 13y = 0$.

Characteristic: $r = -2 \pm 3i$. General solution: $y(x) = e^{-2x}(C_1 \cos 3x + C_2 \sin 3x)$.

Example 4: Higher Order, Mixed Roots

Solve $y''' - y' = 0$.

Characteristic: $r(r^2 - 1) = 0 \Rightarrow r \in \{0, 1, -1\}$. General solution: $y(x) = C_1 + C_2e^x + C_3e^{-x}$.

Example 5: Repeated Complex Roots

Solve $\frac{d^4y}{dx^4} + 8\frac{d^3y}{dx^3} + 42\frac{d^2y}{dx^2} + 104\frac{dy}{dx} + 169y = 0$.

Characteristic: $(r^2 + 4r + 13)^2 = 0$ (i.e., $r = -2 \pm 3i$ with multiplicity 2). General solution: $y(x) = e^{-2x}[(C_1 + C_3x) \cos(3x) + (C_2 + C_4x) \sin(3x)]$.

Quick Reminder

- **Midterm exam:** Next week.
- **Assignment 1 result:** Will be released before Oct. 18 (the day before the midterm).
- **Next tutorial:** Will be a review session for the midterm.