

# MATB44 - Week 10 Tutorial Teaching Notes

## Week 10 Topics

### Reduction of Order for Second-Order Linear Homogeneous ODEs

- Method: finding a second linearly independent solution when one solution is known
- Application to variable-coefficient equations
- Wronskian and linear independence

### Definition and Theory

#### Reduction of Order Method

For a second-order linear homogeneous ODE in standard form:

$$y'' + p(x)y' + q(x)y = 0,$$

if one solution  $y_1(x)$  is known, a second linearly independent solution  $y_2(x)$  can be found using the formula:

$$y_2(x) = y_1(x) \int \frac{e^{-\int p(x) dx}}{[y_1(x)]^2} dx.$$

#### General Form (Non-Standard)

For equations of the form:

$$a_2(x)y'' + a_1(x)y' + a_0(x)y = 0,$$

first convert to standard form by dividing by  $a_2(x)$  (where  $a_2(x) \neq 0$ ):

$$y'' + \frac{a_1(x)}{a_2(x)}y' + \frac{a_0(x)}{a_2(x)}y = 0,$$

so that  $p(x) = \frac{a_1(x)}{a_2(x)}$  and  $q(x) = \frac{a_0(x)}{a_2(x)}$ .

### Key Points

- The method relies on the fact that if  $y_1$  is a solution, then  $y_2 = v(x)y_1$  is also a solution for an appropriate function  $v(x)$ .
- The substitution  $y = vy_1$  reduces the second-order ODE to a first-order ODE in  $v'$ .
- The Wronskian  $W(y_1, y_2) = y_1y_2' - y_1'y_2$  should be nonzero to ensure linear independence.

### Key Examples

#### Example 1: Reduction of Order (Cauchy-Euler Type)

Solve:

$$x^2y'' + xy' - y = 0,$$

given that  $y_1 = x$  is a solution.

**Solution:**

**Step 1.** Convert to standard form: Divide by  $x^2$  (for  $x \neq 0$ ):

$$y'' + \frac{1}{x}y' - \frac{1}{x^2}y = 0.$$

Here  $p(x) = \frac{1}{x}$  and  $q(x) = -\frac{1}{x^2}$ .

**Step 2.** Apply reduction of order formula:

$$y_2(x) = y_1(x) \int \frac{e^{-\int p(x) dx}}{[y_1(x)]^2} dx = x \int \frac{e^{-\int \frac{1}{x} dx}}{x^2} dx.$$

**Step 3.** Compute the integrals:

$$e^{-\int \frac{1}{x} dx} = e^{-\ln|x|} = \frac{1}{|x|} = \frac{1}{x} \quad (\text{for } x > 0).$$

Therefore:

$$y_2(x) = x \int \frac{1/x}{x^2} dx = x \int \frac{1}{x^3} dx = x \cdot \left(-\frac{1}{2x^2}\right) = -\frac{1}{2x}.$$

**Step 4.** General solution:

$$y(x) = C_1 y_1(x) + C_2 y_2(x) = C_1 x + C_2 \cdot \left(-\frac{1}{2x}\right) = C_1 x - \frac{C_2}{2x}.$$

Or, absorbing the constant:  $y(x) = C_1 x + \frac{C_2}{x}$ .

---

### Example 2: Reduction of Order (Constant Coefficient Case)

Solve:

$$y'' - 4y' + 4y = 0,$$

given that  $y_1 = e^{2x}$  is a solution.

**Solution:**

**Step 1.** Standard form:  $y'' - 4y' + 4y = 0$ , so  $p(x) = -4$ .

**Step 2.** Apply formula:

$$y_2(x) = e^{2x} \int \frac{e^{-\int (-4) dx}}{(e^{2x})^2} dx = e^{2x} \int \frac{e^{4x}}{e^{4x}} dx = e^{2x} \int 1 dx = x e^{2x}.$$

**Step 3.** General solution:

$$y(x) = C_1 e^{2x} + C_2 x e^{2x}.$$

(Note: This matches the result from the characteristic equation method with a repeated root  $r = 2$ .)

### Example 3: Reduction of Order (Variable Coefficients)

Solve:

$$xy'' - (x+1)y' + y = 0,$$

given that  $y_1 = e^x$  is a solution.

**Solution:**

**Step 1.** Convert to standard form:

$$y'' - \frac{x+1}{x}y' + \frac{1}{x}y = 0, \quad x \neq 0.$$

So  $p(x) = -\frac{x+1}{x} = -1 - \frac{1}{x}$ .

**Step 2.** Compute the integrating factor:

$$e^{-\int p(x) dx} = e^{-\int (-1 - \frac{1}{x}) dx} = e^{x + \ln|x|} = e^x \cdot |x| = xe^x \quad (\text{for } x > 0).$$

**Step 3.** Apply formula:

$$y_2(x) = e^x \int \frac{xe^x}{(e^x)^2} dx = e^x \int \frac{xe^x}{e^{2x}} dx = e^x \int xe^{-x} dx.$$

**Step 4.** Integrate by parts:

$$\int xe^{-x} dx = -xe^{-x} - e^{-x} = -e^{-x}(x+1).$$

**Step 5.** Final result:

$$y_2(x) = e^x \cdot (-e^{-x}(x+1)) = -(x+1).$$

**Step 6.** General solution:

$$y(x) = C_1 e^x + C_2(x+1).$$

---

### Quick Recipe

1. **Identify the standard form:** Write the ODE as  $y'' + p(x)y' + q(x)y = 0$ .
2. **Verify the known solution:** Check that  $y_1$  satisfies the ODE.
3. **Apply the formula:**

$$y_2(x) = y_1(x) \int \frac{e^{-\int p(x) dx}}{[y_1(x)]^2} dx.$$

4. **Simplify the integral:** Compute  $e^{-\int p(x) dx}$  and the integral.
5. **Write the general solution:**  $y(x) = C_1 y_1(x) + C_2 y_2(x)$ .

### Practice Problems

1.  $x^2 y'' - 3xy' + 4y = 0$ , given  $y_1 = x^2$ .
2.  $y'' + 2y' + y = 0$ , given  $y_1 = e^{-x}$ .
3.  $(1-x^2)y'' - 2xy' + 2y = 0$ , given  $y_1 = x$  (for  $|x| < 1$ ).
4.  $xy'' + 2y' + xy = 0$ , given  $y_1 = \frac{\sin x}{x}$ .