

On the Number of Edges of a Fan-Crossing Free Graph

November 6, 2013

Heuna Kim


Joint Work with Otfried Cheong, Sarel Har-Peled, Hyosil Kim

Motivation on Planar graphs

When do planar graphs achieve
the maximum number of edges?

Motivation on Planar graphs

When do planar graphs achieve
the maximum number of edges?





By Euler's formula, $e \leq 3n - 6$ where
 e : #edges, n : #vertices

Problem Settings


- a *topological* graph
 - vertices = points on the plane
 - edges = Jordan curves connecting the vertices

- *simple*



Definition

- A *radial (p, q) -grid*
 - a set of $p + q$ edges such that
 - p edges have a common endpoint, and
 - these p edges cross the remaining q edges.
- A *k -fan-crossing*
 - a radial $(k, 1)$ -grid
- A *fan-crossing*
 - a radial $(2, 1)$ -grid




Definition

- A *radial (p, q) -grid*
 - a set of $p + q$ edges such that
 - p edges have a common endpoint, and
 - these p edges cross the remaining q edges.
- A *k -fan-crossing*
 - a radial $(k, 1)$ -grid
- A *fan-crossing*
 - a radial $(2, 1)$ -grid

Question:

the maximum number
of edges?



Results : the Maximum Number of Edges?

($n > 5$:#vertices, $k > 3$)

- Tight bounds
 - A *fan-crossing free* graph for *all* $n \neq 6, 7, 9$: $4n - 8$ edges
 - A *fan-crossing free* graph for $n = 6, 7, 9$: $4n - 9$ edges
 - A *straight fan-crossing free* graph for *all* $n \neq 6$: $4n - 9$ edges
- Upper bounds
 - A *k-fan-crossing free* graph : $\leq 3(k - 1)(n - 2)$ edges
 - A graph G s.t $|E(G_e)| = O(n^{1+\alpha})$ for *all* $e \in E(G)$:
 $O(n^{1+\alpha})$ edges if $\alpha > 0$, and $O(n \log^2 n)$ edges if $\alpha = 0$
 - G_e : the subgraph of G containing exactly those edges that cross e
 - G_e must have a monotone graph property.

Results : the Maximum Number of Edges?

$(n > 5: \# \text{vertices}, k > 3)$

- Tight bounds
 - A *fan-crossing free* graph for *all* $n \neq 6, 7, 9$: $4n - 8$ edges
 - A *fan-crossing free* graph for $n = 6, 7, 9$: $4n - 9$ edges

Outline

- Lower bounds
- Upper bounds

Outline


- Lower bounds
- Upper bounds

Lower Bounds: Fan-crossing Free Graphs

When do fan-crossing free graphs achieve
the maximum number of edges?

Lower Bounds: Fan-crossing Free Graphs

When do fan-crossing free graphs achieve
the maximum number of edges?





ex)
13 vertices
44 edges

$$3n - 6 + \frac{2n-4}{2} = 4n - 8 \text{ edges}$$

Lower Bounds: Fan-crossing Free Graphs

When do fan-crossing free graphs achieve
the maximum number of edges?



Outline

- Lower bounds
- Upper bounds

Outline

- Lower bounds
- Upper bounds

Graph Drawing
Problems




Conversion
Tricks



Purely Combinatorial
Questions

A Toy Example of Counting Edges of G



For a face ψ of a maximal plane subgraph H of G ,

- $m(\psi)$: $\#$ edges on the boundary of ψ ; the complexity of ψ
- $a(\psi)$: $\#$ arrows that ψ possesses

$$2|E(G)| = \sum_{\psi \in \mathcal{F}_H} (m(\psi) + a(\psi)) = 8 \cdot (3 + 1) + 2 \cdot 3$$

When are we able to bound $a(\psi)$ in terms of $m(\psi)$?

Outline

- Lower bounds
- Upper bounds

Graph Drawing Problems



Conversion Tricks



Purely Combinatorial Questions


What is the maximum number of edges that can be drawn without fan crossings?

conversion to m -stars

An m -star

- What is an m -star?
- How many arrows in an m -star?
- How an m -star looks like

m -star



m -star :

A *purely combinatorial* m -gon φ
with a set of *arrows*.

- *short* arrows : of length 1
- *long* arrows : of length greater than 1

at most $f_l(m)$ long arrows,
at most $f_s(m)$ short arrows


Bound the Number of Arrows

For an m -star in a fan-crossing free graph,

$$f_s(m) \leq m - 1 \quad \text{if } m \geq 3$$


$$f_l(m) \leq \begin{cases} 0 & \text{if } m = 3, 4 \\ 2(m - 4) & \text{if } m \geq 5 \end{cases}$$

Proof for Short Arrows




$$f_s(m) \leq m - 1 \quad \text{if } m \geq 3$$

Base Cases : Long Arrows



- 3-star
 - no long arrows




- 4-star
 - no long arrows


Induction : Long Arrows

- $m \geq 5$. We delete all short arrows.
- e : a shortest long arrow, $\ell \geq 2$: length of e

an m -star





an $(m - \ell + 1)$ -star



$$f_l(m - \ell + 1) \geq f_l(m) - |\text{all arrows from } v_2 \sim v_\ell| - |\text{new short arrows}|$$

Induction : Long Arrows


- $m \geq 5$. We delete all short arrows.
- e : a shortest long arrow, $\ell \geq 2$: length of e



$$f_l(m - \ell + 1) \geq f_l(m) - |\text{all arrows from } v_2 \sim v_\ell| - |\text{new short arrows}|$$

$$f_l(m) \leq f_l(m - \ell + 1) + \ell$$

How does an m -star look like?



If a maximal plane subgraph H is connected,

$$\begin{aligned} &\text{each face } \psi : m(\psi)\text{-star} \\ &a(\psi) \leq f_l(m(\psi)) + f_s(m(\psi)) \end{aligned}$$

What if H is not connected...

Outline

- Lower bounds
- Upper bounds

Graph Drawing Problems



Conversion Tricks



Purely Combinatorial Questions

What is the maximum number of edges that can be drawn without fan crossings?



conversion to m -stars

An m -star

- What is an m -star?
- Bounds for the number of arrows
- How an m -star looks like

If a face ψ of H is not an m -star

Assume there is an arrow starting in a vertex of ξ and ending in an edge of ζ (otherwise, count them separately).



- $p(\psi) : \#(\text{boundary chains in } \psi)$
- $m(\varphi) = m(\psi) + 3(p(\psi) - 1)$
- $w(\varphi) = m(\psi) + p(\psi) - 1$
- $a(\psi) \leq f_l(m(\varphi)) + f_s(w(\varphi)) + (p(\psi) - 1)$

Outline

- Lower bounds
- Upper bounds

Graph Drawing Problems



Conversion Tricks



Purely Combinatorial Questions

What is the maximum number of edges that can be drawn without fan crossings?

conversion to m -stars

An m -star

- What is an m -star?
- Bounds for the number of arrows
- How an m -star looks like

A bound of $a(\psi)$ for ψ in general

Remind that

$$a(\psi) \leq f_l(m(\psi)) + f_s(m(\psi)) \quad \text{if } p(\psi) = 1$$

$$a(\psi) \leq f_l(m(\varphi)) + f_s(w(\varphi)) + (p(\psi) - 1) \quad \text{if } p(\psi) > 1$$

fan-crossing free :

$$a(\psi) \leq 3m(\psi) + 8p(\psi) - 17$$

k -fan-crossing free ($k \geq 3$) :

$$a(\psi) \leq 3(k-1)(m(\psi) + 2p(\psi) - 4) - (2m - 3)$$

Compute the sum of $m(\psi) + a(\psi)$

For a radial $(k, 1)$ -grid free graph G on n vertices and e edges,
 H : a maximal plane subgraph, \mathcal{F} : the set of faces in H .

$$\begin{aligned} 2e &\leq \sum_{\psi \in \mathcal{F}} m(\psi) + \sum_{\psi \in \mathcal{F}} a(\psi) \\ &\leq \begin{cases} \sum_{\psi \in \mathcal{F}} (m(\psi) + 3m(\psi) + 8p(\psi) - 16) \\ \sum_{\psi \in \mathcal{F}} \{3(k-1)(m(\psi) + 2p(\psi) - 4)\} \end{cases} \\ &\leq \begin{cases} 8n - 16 & \text{if } k = 2 \\ 6(k-1)(n-2) & \text{if } k \geq 3 \end{cases} \end{aligned}$$

using Euler's formula ($n - m + r = 1 + p$) on H with
 $\sum_{\psi \in \mathcal{F}} m(\psi) = 2m$ and $\sum_{\psi \in \mathcal{F}} (p(\psi) - 1) = p - 1$.

Questions

- What is the better bounds for k -fan-crossing free graphs?
- For a given adjacency matrix of a fan-crossing free graph, find an embedding (a drawing) of the graph without fan crossings.
- For a given adjacency matrix of a graph, test if the graph is fan-crossing free. (Testing if a graph is 1-planar is NP-hard.)
- When a graph is not a topological graph, for example, vertices are points on a torus, find the maximum number of edges of a fan-crossing free graph.

Thank you


Problem Settings

- a **monotone** *graph property* \mathcal{P}
 - preserved under edge-deletions
- A graph $G = (V, E)$ has a **derived** *graph property* \mathcal{P}^* of a monotone graph property \mathcal{P}
if G_e has \mathcal{P} for all e in E .
 - G_e : the subgraph of G containing exactly those edges that cross e
- A special case of \mathcal{P} and \mathcal{P}^*
 - \mathcal{P} : G is $K_{1,k}$ free $\Rightarrow \mathcal{P}^*$: a radial $(k, 1)$ -grid free

General bounds

Lemma 1 (Theorem 2.1 , Pach et al.) *Let G be a graph with n vertices of degree d_1, \dots, d_n and crossing number χ . Then there is a subset E of b edges of G such that removing E from G creates components of size at most $2n/3$, and*

$$b^2 \leq (1.58)^2 \left(16\chi + \sum_{i=1}^n d_i^2 \right).$$



General bounds

For a given graph G on n vertices and m edges,

G_e : the subgraph of G containing exactly those edges that cross e .

G_e has a monotone graph property.

If G_e has $O(n^{1+\alpha})$ for all e in G ,

G has $O(n^{1+\alpha})$ edges if $\alpha > 0$, and $O(n \log^2 n)$ edges if $\alpha = 0$.

Proof idea. Recursively subdivide G by removing b_0, b_1, \dots, b_p edges.

$b_i = O(\sqrt{mn} \left(\left(\frac{2}{3}\right)^i n\right)^{\alpha/2})$ if $\alpha > 0$ and $b_i = O(\sqrt{mn})$ by Lemma 1 since $\chi = O(n^{1+\alpha})$. Then, $m \leq \sum_{i=1}^p b_i$ and $p = O(\log n)$

