

# The Zeros of Elliptic Curve $L$ -Functions

Simon Spicer

University of Washington

*mlungu@uw.edu*

July 8, 2013

# Overview

- Motivated by a challenge question from Barry Mazur in the upcoming paper "How Explicit is the Explicit Formula?"
- Prove an explicit version of the explicit formula for elliptic curve  $L$ -functions, i.e. one with explicit error bounds for truncated sums over  $L$ -function zeros
- Applicable to work of Mazur, Sarnak et al
- This talk more about what to do with elliptic curve  $L$ -function zeros once you have them, as apposed to how to compute them in the first place

# Motivating Example

- Let  $\zeta(s)$  be the meromorphic continuation to  $\mathbb{C}$  of  $\sum_{n=1}^{\infty} n^{-s}$ .

# Motivating Example

- Let  $\zeta(s)$  be the meromorphic continuation to  $\mathbb{C}$  of  $\sum_{n=1}^{\infty} n^{-s}$ .
- Then  $\zeta(s)$  has a single simple pole at  $s = 1$ , and simple zeros at  $2\mathbb{Z}_{<0}$ .

# Motivating Example

- Let  $\zeta(s)$  be the meromorphic continuation to  $\mathbb{C}$  of  $\sum_{n=1}^{\infty} n^{-s}$ .
- Then  $\zeta(s)$  has a single simple pole at  $s = 1$ , and simple zeros at  $2\mathbb{Z}_{<0}$ .
- All other zeros lie in the vertical strip  $0 < \Re(s) < 1$ , symmetric about the real axis.

# Motivating Example

- Let  $\zeta(s)$  be the meromorphic continuation to  $\mathbb{C}$  of  $\sum_{n=1}^{\infty} n^{-s}$ .
- Then  $\zeta(s)$  has a single simple pole at  $s = 1$ , and simple zeros at  $2\mathbb{Z}_{<0}$ .
- All other zeros lie in the vertical strip  $0 < \Re(s) < 1$ , symmetric about the real axis.

## Conjecture (Riemann Hypothesis)

*All nontrivial zeros of  $\zeta$  are simple and lie on the line  $\Re(s) = \frac{1}{2}$ .*

# The Zeros of $\zeta$

The imaginary parts of the first few zeros of  $\zeta(s)$  in the upper half plane are

14.134725142...

21.022039639...

25.010857580...

30.424876126...

32.935061588...

37.586178159...

40.918719012...

43.327073281...

48.005150881...

49.773832478...

52.970321478...

56.446247697...


59.347044003...

60.831778525...

# The Zeros of $\zeta$

The imaginary parts of the first few zeros of  $\zeta(s)$  in the upper half plane are

14.134725142...  
21.022039639...  
25.010857580...  
30.424876126...  
32.935061588...  
37.586178159...  
40.918719012...  
43.327073281...  
48.005150881...  
49.773832478...  
52.970321478...  
56.446247697...  
59.347044003...  
60.831778525...





# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.


# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.




# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.




# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.




# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.




# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.




# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.




# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.






# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.




# The Explicit Formula for $\zeta(s)$

## The Riemann zeta function $\zeta(s)$

Consider as a function of  $x > 1$  the sum

$$S_{\zeta}(x, T) = \sum_{|\rho| < T} \frac{x^{\rho}}{\rho} = \sqrt{x} \left( \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma} \right)$$

contingent on RH, where  $\gamma$  runs over imaginary parts of nontrivial zeros.



# The Explicit Formula for $\zeta(s)$

What does this sum converge to?

# The Explicit Formula for $\zeta(s)$

What does this sum converge to?

Theorem (Riemann 1858, von Mangoldt 1905)

$$\sum_{\rho} \frac{x^{\rho}}{\rho} = \lim_{T \rightarrow \infty} S_{\zeta}(x, T) = x - \frac{1}{2} \log(1 - 1/x^2) - \log(2\pi) - \psi_{\zeta}(x)$$

where  $\psi_{\zeta}(x) = \sum'_{p^e \leq x} \log p$  is the second Chebyshev function.

# The Explicit Formula for $\zeta(s)$

What does this sum converge to?


Theorem (Riemann 1858, von Mangoldt 1905)

$$\sum_{\rho} \frac{x^{\rho}}{\rho} = \lim_{T \rightarrow \infty} S_{\zeta}(x, T) = x - \frac{1}{2} \log(1 - 1/x^2) - \log(2\pi) - \psi_{\zeta}(x)$$


where  $\psi_{\zeta}(x) = \sum'_{p^e \leq x} \log p$  is the second Chebyshev function.

This is known as (one formulation of) *the explicit formula* for  $\zeta(s)$ .

# The Explicit Formula for $\zeta(s)$



# The Explicit Formula for $\zeta(s)$



## My Goal

Prove the explicit formula for elliptic curve  $L$ -functions, with error bounds for  $S_E(x, T)$ .

# Elliptic Curves

## Definition

An elliptic curve  $E$  is a smooth projective genus 1 algebraic curve with a marked point  $\mathcal{O}$ .



# Elliptic Curves

## Definition

An elliptic curve  $E$  is a smooth projective genus 1 algebraic curve with a marked point  $\mathcal{O}$ .

## For This Talk:

$$E/\mathbb{Q}: y^2 = x^3 + Ax + B, \quad A, B \in \mathbb{Z}$$

# Elliptic Curves

## Definition

An elliptic curve  $E$  is a smooth projective genus 1 algebraic curve with a marked point  $\mathcal{O}$ .

## For This Talk:

$$E/\mathbb{Q}: y^2 = x^3 + Ax + B, \quad A, B \in \mathbb{Z}$$

## Example

$$E = 37a: y^2 = x^3 - 16x + 16$$




Figure: The Elliptic Curve 37a

# Elliptic Curves

## Definition

An elliptic curve  $E$  is a smooth projective genus 1 algebraic curve with a marked point  $\mathcal{O}$ .

## For This Talk:

$$E/\mathbb{Q}: y^2 = x^3 + Ax + B, \quad A, B \in \mathbb{Z}$$

## Example

$$E = 37a: y^2 = x^3 - 16x + 16$$




Figure: The Elliptic Curve 37a

## Theorem (Mordell 1922, Weil 1928)

$$E(\mathbb{Q}) \approx E(\mathbb{Q})_{\text{TOR}} \times \mathbb{Z}^r$$

where  $E(\mathbb{Q})_{\text{TOR}}$  is a finite abelian group, and  $r \in \mathbb{Z}_{\geq 0}$  is the algebraic rank of  $E/\mathbb{Q}$ .

## Theorem (Mordell 1922, Weil 1928)

$$E(\mathbb{Q}) \approx E(\mathbb{Q})_{\text{TOR}} \times \mathbb{Z}^r$$

where  $E(\mathbb{Q})_{\text{TOR}}$  is a finite abelian group, and  $r \in \mathbb{Z}_{\geq 0}$  is the algebraic rank of  $E/\mathbb{Q}$ .

## Example

For  $E = 37a$ , we have  $E(\mathbb{Q}) \approx \mathbb{Z}^1$ , generated by  $P = (0, 4)$ :

$n$	0	1	2	3	4	5	6
$nP$	$\mathcal{O}$	$(0, 4)$	$(4, 4)$	$(-4, -4)$	$(8, -20)$	$(1, -1)$	$(24, 116)$


$n$	7	8	9
$nP$	$(-\frac{20}{9}, \frac{172}{27})$	$(\frac{84}{25}, -\frac{52}{125})$	$(-\frac{80}{49}, -\frac{2108}{343})$

# Elliptic Curves over finite fields

## Example

$$E = 37a : y^2 = x^3 - 16x + 16$$

Consider its solutions  $(x, y)$   
modulo 101, e.g.  $(40, 7)$ :




# Elliptic Curves over finite fields

## Example

$$E = 37a : y^2 = x^3 - 16x + 16$$

Consider its solutions  $(x, y)$   
modulo 101, e.g.  $(40, 7)$ :



## Definition

- For  $p$  prime with good reduction,  $a_p(E) = a_p := p + 1 - \#E(\mathbb{F}_p)$
- For bad primes,  $a_p := 0, 1$  or  $-1$  depending on reduction type.

# Elliptic Curves over finite fields

Theorem (Hasse, 1936)

$$|a_p| \leq 2\sqrt{p} \quad \text{for all } p.$$



# Elliptic Curves over finite fields


## Theorem (Hasse, 1936)

$$|a_p| \leq 2\sqrt{p} \quad \text{for all } p.$$

## Example

For  $E = 37a$ ,

$p$	2	3	5	7	11	13	17	19	23	29	31	37
$a_p$	-2	-3	-2	-1	-5	-2	0	0	2	6	-4	-1



# The Conductor of a Curve

## Definition

The conductor of  $E$  is  $N = \prod_p p^{f_p(E)}$ , where

$$f_p(E) = \begin{cases} 0, & p \text{ good} \\ 1, & \text{mult. reduction at } p \\ 2, & \text{add. reduction at } p, \end{cases}$$

for  $p \neq 2, 3$  and possibly more for 2 and 3.

# The Conductor of a Curve

## Definition

The conductor of  $E$  is  $N = \prod_p p^{f_p(E)}$ , where

$$f_p(E) = \begin{cases} 0, & p \text{ good} \\ 1, & \text{mult. reduction at } p \\ 2, & \text{add. reduction at } p, \end{cases}$$

for  $p \neq 2, 3$  and possibly more for 2 and 3.

## Example

The conductor of  $37a$  is  $N = 37$ , hence its name.

# Elliptic Curve $L$ -Functions

## Definition

The  $L$ -function attached to  $E$  is

$$L_E(s) := \prod_{p|N} \frac{1}{1 - a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1 - a_p p^{-s} + p^{1-2s}} = \sum_{n=1}^{\infty} a_n n^{-s}$$

for  $\Re(s) > \frac{3}{2}$ .

The  $a_n$  are defined by multiplying out the Euler product.

# Elliptic Curve $L$ -Functions

## Definition

The  $L$ -function attached to  $E$  is

$$L_E(s) := \prod_{p|N} \frac{1}{1 - a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1 - a_p p^{-s} + p^{1-2s}} = \sum_{n=1}^{\infty} a_n n^{-s}$$

for  $\Re(s) > \frac{3}{2}$ .

The  $a_n$  are defined by multiplying out the Euler product.

## Definition

The *completed*  $L$ -function attached to  $E$  is

$$\Lambda_E(s) := N^{s/2} (2\pi)^{-s} \Gamma(s) L_E(s)$$

# Modularity & Analytic Continuation of $L_E(s)$

Theorem (Breuille, Conrad, Diamond, Taylor, Wiles et al, 1999,2001)

*There exists an integral newform  $f \in S_2(\Gamma_0(N))$  s.t.  $L_f(s) = L_E(s)$ .  
That is, there exists a holomorphic function  $f$  on  $\mathbb{H}$  with Fourier decomposition  $f(z) = \sum_n a_n(f) e^{2\pi i n z}$  such that  $a_n(f) = a_n(E)$ .*

# Modularity & Analytic Continuation of $L_E(s)$

Theorem (Breuille, Conrad, Diamond, Taylor, Wiles et al, 1999,2001)

*There exists an integral newform  $f \in S_2(\Gamma_0(N))$  s.t.  $L_f(s) = L_E(s)$ .  
That is, there exists a holomorphic function  $f$  on  $\mathbb{H}$  with Fourier decomposition  $f(z) = \sum_n a_n(f) e^{2\pi i n z}$  such that  $a_n(f) = a_n(E)$ .*

## Corollary

*$L_E(s)$  extends to an entire function on  $\mathbb{C}$ . Specifically,*

$$\Lambda(s) = w\Lambda(2-s),$$

*where  $w = \pm 1$ .*

# The Zeros of $L_E(s)$

Three flavors:

- A simple zero at  $0, -1, -2, -3, \dots$
- A zero of order  $r_{an}$  at  $s = 1$ ;  $r_{an}$  is called the *analytic rank* of  $E$
- Countably infinite zeros in the strip  $0 < \Re(s) < 2$ , symmetric about  $\Re(s) = 1$  and  $x$ -axis.



# The Zeros of $L_E(s)$

Three flavors:

- A simple zero at  $0, -1, -2, -3, \dots$
- A zero of order  $r_{an}$  at  $s = 1$ ;  $r_{an}$  is called the *analytic rank* of  $E$
- Countably infinite zeros in the strip  $0 < \Re(s) < 2$ , symmetric about  $\Re(s) = 1$  and  $x$ -axis.

## Conjecture (Generalized Riemann Hypothesis for Elliptic Curves)

*All nontrivial zeros of  $L_E(s)$  are simple and lie on the line  $\Re(s) = 1$ .*

# The Zeros of $L_E(s)$

Three flavors:

- A simple zero at  $0, -1, -2, -3, \dots$
- A zero of order  $r_{an}$  at  $s = 1$ ;  $r_{an}$  is called the *analytic rank* of  $E$
- Countably infinite zeros in the strip  $0 < \Re(s) < 2$ , symmetric about  $\Re(s) = 1$  and x-axis.

Conjecture (Generalized Riemann Hypothesis for Elliptic Curves)

*All nontrivial zeros of  $L_E(s)$  are simple and lie on the line  $\Re(s) = 1$ .*




Figure: The zeros of  $L_E(s)$  for  $E = 37a$

# The BSD Conjecture

## Conjecture (Birch, Swinnerton-Dyer 1960s)

- $r_{an} = r$ , i.e. the order of vanishing of  $L_E(s)$  at  $s = 1$  equals the rank of the free part of  $E(\mathbb{Q})$

# The BSD Conjecture

## Conjecture (Birch, Swinnerton-Dyer 1960s)

- $r_{an} = r$ , i.e. the order of vanishing of  $L_E(s)$  at  $s = 1$  equals the rank of the free part of  $E(\mathbb{Q})$
- The leading coefficient of  $L_E(s)$  at  $s = 1$  is

$$\frac{\Omega_E \cdot \text{Reg}_E \cdot \#\text{III}(E/\mathbb{Q}) \cdot \prod_p c_p}{(\#E_{\text{Tor}}(\mathbb{Q}))^2}$$

# The BSD Conjecture

## Conjecture (Birch, Swinnerton-Dyer 1960s)

- $r_{an} = r$ , i.e. the order of vanishing of  $L_E(s)$  at  $s = 1$  equals the rank of the free part of  $E(\mathbb{Q})$
- The leading coefficient of  $L_E(s)$  at  $s = 1$  is

$$\frac{\Omega_E \cdot \text{Reg}_E \cdot \#\text{III}(E/\mathbb{Q}) \cdot \prod_p c_p}{(\#E_{\text{Tor}}(\mathbb{Q}))^2}$$

where

- ▶  $\Omega_E$  is the real period of (an optimal model of)  $E$ ,
- ▶  $\text{Reg}_E$  is the regulator of  $E$ ,
- ▶  $\#\text{III}(E/\mathbb{Q})$  is the order of the Shafarevich-Tate group attached to  $E/\mathbb{Q}$ ,
- ▶  $\prod_p c_p$  is the product of the Tamagawa numbers of  $E$ , and
- ▶  $\#E_{\text{Tor}}(\mathbb{Q})$  is the number of rational torsion points on  $E$ .

# The Shifted Logarithmic Derivative

To state the explicit formula for elliptic curves, we will need to characterize  $\frac{L'_E}{L_E}(s+1)$ .

# The Shifted Logarithmic Derivative

To state the explicit formula for elliptic curves, we will need to characterize  $\frac{L'_E}{L_E}(s+1)$ .

## Lemma 1 (S.)

$\frac{L'_E}{L_E}(s+1) = \frac{d}{ds} \log(L_E)(s+1)$  has the Dirichlet series  $\sum_n c_n(E) n^{-s}$  which converges absolutely for  $\Re(s) > \frac{1}{2}$ , where

$$c_n(E) := \begin{cases} -\left(p^e + 1 - \#\tilde{E}(\mathbb{F}_{p^e})\right) \cdot \frac{\log(p)}{p^e}, & n = p^e \text{ a prime power,} \\ 0, & \text{otherwise} \end{cases}$$

and  $\#\tilde{E}(\mathbb{F}_{p^e})$  is the number of points on over  $\mathbb{F}_{p^e}$  on the (possibly singular) projective curve obtained by reducing  $E$  modulo  $p$ .

# The Shifted Logarithmic Derivative

Proof (Sketch).

- $L_E(s) := \prod_{p|N} \frac{1}{1-a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1-a_p p^{-s}+p^{1-2s}}$ , so



# The Shifted Logarithmic Derivative

## Proof (Sketch).

- $L_E(s) := \prod_{p|N} \frac{1}{1-a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1-a_p p^{-s}+p^{1-2s}}$ , so
$$\frac{L'_E}{L_E}(s) = - \sum_{p|N} \frac{a_p \log(p) \cdot p^{-s}}{1-a_p p^{-s}} - \sum_{p \nmid N} \frac{a_p \log(p) \cdot p^{-s} - 2p^{k-1} \log(p) \cdot p^{-2s}}{1-a_p p^{-s}+p^{k-1} \cdot p^{-2s}}$$

# The Shifted Logarithmic Derivative

## Proof (Sketch).

- $L_E(s) := \prod_{p|N} \frac{1}{1-a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1-a_p p^{-s}+p^{1-2s}}$ , so
$$\frac{L'_E}{L_E}(s) = - \sum_{p|N} \frac{a_p \log(p) \cdot p^{-s}}{1-a_p p^{-s}} - \sum_{p \nmid N} \frac{a_p \log(p) \cdot p^{-s} - 2p^{k-1} \log(p) \cdot p^{-2s}}{1-a_p p^{-s}+p^{k-1} \cdot p^{-2s}}$$
- For good  $p$ , write  $1 - a_p p^{-s} + p^{k-1} \cdot p^{-2s} = (1 - \alpha_p p^{-s})(1 - \beta_p p^{-s})$ ; invert each denominator as power series in  $p^{-s}$  and multiply out

# The Shifted Logarithmic Derivative

## Proof (Sketch).

- $L_E(s) := \prod_{p|N} \frac{1}{1-a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1-a_p p^{-s}+p^{1-2s}}$ , so  

$$\frac{L'_E}{L_E}(s) = - \sum_{p|N} \frac{a_p \log(p) \cdot p^{-s}}{1-a_p p^{-s}} - \sum_{p \nmid N} \frac{a_p \log(p) \cdot p^{-s} - 2p^{k-1} \log(p) \cdot p^{-2s}}{1-a_p p^{-s}+p^{k-1} \cdot p^{-2s}}$$
- For good  $p$ , write  $1 - a_p p^{-s} + p^{k-1} \cdot p^{-2s} = (1 - \alpha_p p^{-s})(1 - \beta_p p^{-s})$ ; invert each denominator as power series in  $p^{-s}$  and multiply out
- $\Rightarrow \frac{L'_E}{L_E}(s) = \sum_{p|N} \sum_{e \geq 1} -a_p^e \log(p) (p^e)^{-s}$   

$$+ \sum_{p \nmid N} \sum_{e \geq 1} -(\alpha_p^e + \beta_p^e) \log(p) (p^e)^{-s}$$

# The Shifted Logarithmic Derivative

## Proof (Sketch).

- $L_E(s) := \prod_{p|N} \frac{1}{1-a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1-a_p p^{-s}+p^{1-2s}}$ , so  

$$\frac{L'_E}{L_E}(s) = - \sum_{p|N} \frac{a_p \log(p) \cdot p^{-s}}{1-a_p p^{-s}} - \sum_{p \nmid N} \frac{a_p \log(p) \cdot p^{-s} - 2p^{k-1} \log(p) \cdot p^{-2s}}{1-a_p p^{-s}+p^{k-1} \cdot p^{-2s}}$$
- For good  $p$ , write  $1 - a_p p^{-s} + p^{k-1} \cdot p^{-2s} = (1 - \alpha_p p^{-s})(1 - \beta_p p^{-s})$ ; invert each denominator as power series in  $p^{-s}$  and multiply out
- $\Rightarrow \frac{L'_E}{L_E}(s) = \sum_{p|N} \sum_{e \geq 1} -a_p^e \log(p) (p^e)^{-s}$   

$$+ \sum_{p \nmid N} \sum_{e \geq 1} -(\alpha_p^e + \beta_p^e) \log(p) (p^e)^{-s}$$
- By Silverman etc.,  
 $a_p^e = p^e + 1 - \#\tilde{E}(\mathbb{F}_{p^e})$  for bad  $p$ , and  
 $\alpha_p^e + \beta_p^e = p^e + 1 - \#E(\mathbb{F}_{p^e})$  for good  $p$

# The Shifted Logarithmic Derivative

## Proof (Sketch).

- $L_E(s) := \prod_{p|N} \frac{1}{1-a_p p^{-s}} \prod_{p \nmid N} \frac{1}{1-a_p p^{-s}+p^{1-2s}}$ , so
$$\frac{L'_E}{L_E}(s) = -\sum_{p|N} \frac{a_p \log(p) \cdot p^{-s}}{1-a_p p^{-s}} - \sum_{p \nmid N} \frac{a_p \log(p) \cdot p^{-s} - 2p^{k-1} \log(p) \cdot p^{-2s}}{1-a_p p^{-s}+p^{k-1} \cdot p^{-2s}}$$
- For good  $p$ , write  $1 - a_p p^{-s} + p^{k-1} \cdot p^{-2s} = (1 - \alpha_p p^{-s})(1 - \beta_p p^{-s})$ ; invert each denominator as power series in  $p^{-s}$  and multiply out
- $\Rightarrow \frac{L'_E}{L_E}(s) = \sum_{p|N} \sum_{e \geq 1} -a_p^e \log(p) (p^e)^{-s} + \sum_{p \nmid N} \sum_{e \geq 1} -(\alpha_p^e + \beta_p^e) \log(p) (p^e)^{-s}$
- By Silverman etc.,  
 $a_p^e = p^e + 1 - \#\tilde{E}(\mathbb{F}_{p^e})$  for bad  $p$ , and  
 $\alpha_p^e + \beta_p^e = p^e + 1 - \#E(\mathbb{F}_{p^e})$  for good  $p$
- Finally, shift left by 1  $\Rightarrow (p^e)^{-(s+1)} = \frac{1}{p^e} (p^e)^{-s}$  to pick up factor of  $\frac{1}{p^e}$  in coefficients.



## Another Way to Express $\frac{L'_E}{L_E}(s+1)$

### Lemma 2 (S.)

We may express  $\frac{L'_E}{L_E}(s+1)$  as a sum over the zeros of  $L_E(s)$ . Specifically, assuming GRH then for any  $s$  not in the set of zeros of  $L_E(s+1)$ ,

$$\frac{L'_E}{L_E}(s+1) = \left[ \eta + \log \left( \frac{2\pi}{\sqrt{N}} \right) \right] - \sum_{k=1}^{\infty} \frac{s}{k(k+s)} + \sum_{\gamma} \frac{s}{s^2 + \gamma^2}$$

where  $\eta$  is the Euler-Mascheroni constant  $= 0.5772156649 \dots$

and  $\gamma$  ranges over the imaginary parts of all nontrivial zeros of  $L_E$ .

## Another Way to Express $\frac{L'_E}{L_E}(s+1)$

Proof (Sketch).

## Another Way to Express $\frac{L'_E}{L_E}(s+1)$

Proof (Sketch).

- By definition,  $L_E(s) = \left(\frac{2\pi}{\sqrt{N}}\right)^s \Gamma(s)^{-1} \Lambda_E(s)$



## Another Way to Express $\frac{L'_E}{L_E}(s+1)$

### Proof (Sketch).

- By definition,  $L_E(s) = \left(\frac{2\pi}{\sqrt{N}}\right)^s \Gamma(s)^{-1} \Lambda_E(s)$
- $\Rightarrow$  Shifted logarithmic derivative

$$\frac{L'_E}{L_E}(s+1) = \log\left(\frac{2\pi}{\sqrt{N}}\right) - \frac{\Gamma'}{\Gamma}(s+1) + \frac{\Lambda'_E}{\Lambda_E}(s+1)$$

## Another Way to Express $\frac{L'_E}{L_E}(s+1)$

### Proof (Sketch).

- By definition,  $L_E(s) = \left(\frac{2\pi}{\sqrt{N}}\right)^s \Gamma(s)^{-1} \Lambda_E(s)$
- $\Rightarrow$  Shifted logarithmic derivative
$$\frac{L'_E}{L_E}(s+1) = \log\left(\frac{2\pi}{\sqrt{N}}\right) - \frac{\Gamma'}{\Gamma}(s+1) + \frac{\Lambda'_E}{\Lambda_E}(s+1)$$
- $\frac{\Gamma'}{\Gamma}(s+1) = -\eta + \sum_{k=1}^{\infty} \frac{s}{k(k+s)}$  outside negative integers

## Another Way to Express $\frac{L'_E}{L_E}(s+1)$

### Proof (Sketch).

- By definition,  $L_E(s) = \left(\frac{2\pi}{\sqrt{N}}\right)^s \Gamma(s)^{-1} \Lambda_E(s)$
- $\Rightarrow$  Shifted logarithmic derivative
$$\frac{L'_E}{L_E}(s+1) = \log\left(\frac{2\pi}{\sqrt{N}}\right) - \frac{\Gamma'}{\Gamma}(s+1) + \frac{\Lambda'_E}{\Lambda_E}(s+1)$$
- $\frac{\Gamma'}{\Gamma}(s+1) = -\eta + \sum_{k=1}^{\infty} \frac{s}{k(k+s)}$  outside negative integers
- $\frac{\Lambda'_E}{\Lambda_E}(s+1) = \sum_{\gamma} \frac{s}{s^2 + \gamma^2}$ , obtained by logarithmically differentiating Hadamard product of  $\Lambda_E(s+1)$ .



# An Interesting Aside

## Corollary 1

If  $E/\mathbb{Q}$  has conductor  $N$  and analytic rank  $r$  then

$$N > 0.202e^{2r}$$

# An Interesting Aside

## Corollary 1

If  $E/\mathbb{Q}$  has conductor  $N$  and analytic rank  $r$  then

$$N > 0.202e^{2r}$$

Better results (S.), although nowhere close to effective yet:

$r$	$N \geq$	Smallest Known Conductor
0	3	11
1	6	37
2	16	389
3	55	5077
4	232	234446
5	1192	19047851
6	6696	5187563742

## An Interesting Aside

Contingent on GRH and BSD we have a complete description of the Taylor series of  $L_E$  about  $s = 1$ . Specifically:

## An Interesting Aside

Contingent on GRH and BSD we have a complete description of the Taylor series of  $L_E$  about  $s = 1$ . Specifically:

### Corollary 2

Let  $L_E(s+1) = s^r (a + b \cdot s + c \cdot s^2 + O(s^3))$ , where  $a$  is the leading coefficient described by BSD. Then

$$\frac{b}{a} = \eta + \log \left( \frac{2\pi}{\sqrt{N}} \right)$$
$$\frac{c}{a} = \frac{1}{2} \left[ \eta + \log \left( \frac{2\pi}{\sqrt{N}} \right) \right]^2 - \frac{\pi^2}{12} + \sum_{\gamma > 0} \gamma^{-2}$$

Recursive formulae exist for higher coefficients as well.

# The Explicit Formula for Elliptic Curves

## Definition



# The Explicit Formula for Elliptic Curves

## Definition

Let



$$S_E(x, T) := \sum_{|\gamma| < T} \frac{x^{i\gamma}}{i\gamma} = \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma}$$

where  $\gamma$  runs over imaginary parts of nontrivial zeros other than  $s = 1$

# The Explicit Formula for Elliptic Curves

## Definition

Let

•

$$S_E(x, T) := \sum_{|\gamma| < T} \frac{x^{i\gamma}}{i\gamma} = \sum_{0 < \gamma < T} \frac{2 \sin(\gamma \log x)}{\gamma}$$

where  $\gamma$  runs over imaginary parts of nontrivial zeros other than  $s = 1$

•

$$\psi_E(x) := \sum'_{n \leq x} c_n(E)$$

i.e.  $\psi_E(x)$  is the cumulative sum function of the Dirichlet coefficients of  $\frac{L'_E}{L_E}(s+1)$

(Recall  $c_n(E) = \left(p^e + 1 - \#\tilde{E}(\mathbb{F}_{p^e})\right) \cdot \frac{\log(p)}{p^e}$  for  $n = p^e$  and 0 otherwise)

# The Explicit Formula for Elliptic Curves




Figure:  $\psi_E(x)$  for  $E = 37a$

# The Explicit Formula for Elliptic Curves

## Theorem

For any any  $E/\mathbb{Q}$  with conductor  $N$  and for any  $x > 1$  the partial sum function  $S_E(x, T)$  converges as  $T \rightarrow \infty$ . Specifically,


$$\begin{aligned}\lim_{T \rightarrow \infty} S_E(x, T) &= \sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} \\ &= -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)\end{aligned}$$

where  $\eta$  is the Euler-Mascheroni constant  $= 0.5772156649 \dots$

# The Explicit Formula for Elliptic Curves

## Theorem


$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# The Explicit Formula for Elliptic Curves

## Theorem


$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# The Explicit Formula for Elliptic Curves

## Theorem


$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# The Explicit Formula for Elliptic Curves

## Theorem

$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$






# The Explicit Formula for Elliptic Curves

## Theorem


$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# The Explicit Formula for Elliptic Curves

## Theorem


$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# The Explicit Formula for Elliptic Curves

## Theorem


$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# The Explicit Formula for Elliptic Curves

## Theorem


$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# The Explicit Formula for Elliptic Curves

## Theorem

$$\sum_{\gamma > 0} \frac{2 \sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r_{an} \log x - \log(1 - x^{-1}) + \psi_E(x)$$



# Proving the Explicit Formula

How do we go about proving this?:

# Proving the Explicit Formula

How do we go about proving this?:

- Let  $\sigma > \frac{1}{2}$  and consider the integral  $\frac{-1}{2\pi i} \int_{\sigma-iT}^{\sigma+iT} \frac{L'_E}{L_E}(s+1) \frac{x^s}{s} ds$

# Proving the Explicit Formula

How do we go about proving this?:

- Let  $\sigma > \frac{1}{2}$  and consider the integral  $\frac{-1}{2\pi i} \int_{\sigma-iT}^{\sigma+iT} \frac{L'_E}{L_E}(s+1) \frac{x^s}{s} ds$
- Replace  $\frac{L'_E}{L_E}(s+1)$  with two different series representations and distribute
- Replace each integral with contour integral on  $\mathbb{C}$  plus residues
- Contour integrals  $\rightarrow 0$  as  $T \rightarrow \infty$ .



# Using the Cauchy Residue Theorem

## Example


$$\frac{1}{2\pi i} \int_{\sigma-iT}^{\sigma+iT} \frac{x^s}{s} ds = 1 + \frac{1}{2\pi i} \left( \int_{-\infty+iT}^{\sigma+iT} - \int_{-\infty-iT}^{\sigma-iT} \right) \frac{x^s}{s} ds$$



# Using the Cauchy Residue Theorem

## Example

$$\frac{1}{2\pi i} \int_{\sigma-iT}^{\sigma+iT} \frac{x^s}{s} ds = 1 + \frac{1}{2\pi i} \left( \int_{-\infty+iT}^{\sigma+iT} - \int_{-\infty-iT}^{\sigma-iT} \right) \frac{x^s}{s} ds$$



# Improving the Explicit Formula

Current proofs (e.g. Iwaniec & Kowalski, Montgomery & Vaughn) only give asymptotic arguments for proofs.

# Improving the Explicit Formula

Current proofs (e.g. Iwaniec & Kowalski, Montgomery & Vaughn) only give asymptotic arguments for proofs.

## My Work:

Rework Proofs to include explicit constants on error terms.

# Improving the Explicit Formula

Current proofs (e.g. Iwaniec & Kowalski, Montgomery & Vaughn) only give asymptotic arguments for proofs.

## My Work:

Rework Proofs to include explicit constants on error terms.

## Conjecture (S.)

Let  $\epsilon(x, T) = S_E(x, T) + \eta + \log\left(\frac{2\pi}{\sqrt{N}}\right) + r_{an} \log x + \log(1 - x^{-1}) - \psi_E(x)$ .

Then  $\exists$  a positive constant  $M$  such that

$$\epsilon(T, x) < M \cdot \frac{\log^2 T}{T} \cdot \frac{x + 1/x}{\log x} \left( 1 + \sum_{n \neq x} \left| \frac{c_n}{n \log(\frac{x}{n})} \right| \right)$$

for  $T \gg 1$ .

# The Gibbs Phenomenon

## Conjecture (S.)

$$\epsilon(T, x) < M \cdot \frac{\log^2 T}{T} \cdot \frac{x+1/x}{\log x} \left( 1 + \sum_{n \neq x} \left| \frac{c_n}{n \log(\frac{x}{n})} \right| \right) \text{ for } T \gg 1.$$




Figure: The Gibbs Phenomenon clearly visible at jump discontinuities for 37a.

# The Hard Part

Why is this hard?

- Explicit proof requires us to bound integral of  $\frac{\Lambda'_E}{\Lambda_E} (s+1) \frac{x^s}{s}$  across critical strip
- $\Rightarrow$  require explicit bounds on zero density along critical strip for EC  $L$ -functions.

## Some Neat Corollaries

Loosely,  $\{\text{nontrivial zeros of } L_E\} \sim \{a_p(E) : p \text{ prime}\}$  in an information theoretic sense. For example,



## Some Neat Corollaries

Loosely,  $\{\text{nontrivial zeros of } L_E\} \sim \{a_p(E) : p \text{ prime}\}$  in an information theoretic sense. For example,

### Corollary (S.)

$$a_p = \lim_{T \rightarrow \infty} \frac{-2\pi p}{\log p} \cdot \frac{1}{T} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$$

## Some Neat Corollaries

Loosely,  $\{\text{nontrivial zeros of } L_E\} \sim \{a_p(E) : p \text{ prime}\}$  in an information theoretic sense. For example,

### Corollary (S.)

$$a_p = \lim_{T \rightarrow \infty} \frac{-2\pi p}{\log p} \cdot \frac{1}{T} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$$




Figure:  $\frac{-2\pi p}{\log p} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$  as a function of  $T$  for various small  $p$

## Some Neat Corollaries

Loosely,  $\{\text{nontrivial zeros of } L_E\} \sim \{a_p(E) : p \text{ prime}\}$  in an information theoretic sense. For example,

### Corollary (S.)

$$a_p = \lim_{T \rightarrow \infty} \frac{-2\pi p}{\log p} \cdot \frac{1}{T} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$$




Figure:  $\frac{-2\pi p}{\log p} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$  as a function of  $T$  for various small  $p$

## Some Neat Corollaries

Loosely,  $\{\text{nontrivial zeros of } L_E\} \sim \{a_p(E) : p \text{ prime}\}$  in an information theoretic sense. For example,

### Corollary (S.)

$$a_p = \lim_{T \rightarrow \infty} \frac{-2\pi p}{\log p} \cdot \frac{1}{T} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$$




Figure:  $\frac{-2\pi p}{\log p} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$  as a function of  $T$  for various small  $p$

## Some Neat Corollaries

Loosely,  $\{\text{nontrivial zeros of } L_E\} \sim \{a_p(E) : p \text{ prime}\}$  in an information theoretic sense. For example,

### Corollary (S.)

$$a_p = \lim_{T \rightarrow \infty} \frac{-2\pi p}{\log p} \cdot \frac{1}{T} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$$




Figure:  $\frac{-2\pi p}{\log p} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$  as a function of  $T$  for various small  $p$

## Some Neat Corollaries

Loosely,  $\{\text{nontrivial zeros of } L_E\} \sim \{a_p(E) : p \text{ prime}\}$  in an information theoretic sense. For example,

### Corollary (S.)

$$a_p = \lim_{T \rightarrow \infty} \frac{-2\pi p}{\log p} \cdot \frac{1}{T} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$$




Figure:  $\frac{-2\pi p}{\log p} \sum_{0 < \gamma < T} \frac{\cos(\gamma \log p)}{\gamma}$  as a function of  $T$  for various small  $p$

# Some Neat Corollaries

## Conjecture - Alternate BSD (Sarnak, Mazur)

For any given  $E/\mathbb{Q}$ ,

$$\lim_{x \rightarrow \infty} \frac{1}{\log(x)} \sum_{p \leq x} \frac{-a_p \log(p)}{p} = r$$

# Some Neat Corollaries

## Conjecture - Alternate BSD (Sarnak, Mazur)

For any given  $E/\mathbb{Q}$ ,

$$\lim_{x \rightarrow \infty} \frac{1}{\log(x)} \sum_{p \leq x} \frac{-a_p \log(p)}{p} = r$$

## Where does this comes from?

Take explicit formula:

$$\sum_{\gamma} \frac{\sin(\gamma \log x)}{\gamma} = -\eta - \log \left( \frac{2\pi}{\sqrt{N}} \right) - r \log x - \log(1 - 1/x) + \psi_E(x)$$

Divide both sides by  $\log(x)$  and take limits\*.



# Future Work

# Future Work

- Generalize to modular  $L$ -functions of arbitrary weight + twist

# Future Work

- Generalize to modular  $L$ -functions of arbitrary weight + twist
- Computing zeros more efficiently
  - ▶ Current best algorithms are polynomial in  $N$
  - ▶ Would be great to get methods that are polynomial in  $\log(N)$ , or insensitive to  $N$  entirely

# Future Work

- Generalize to modular  $L$ -functions of arbitrary weight + twist
- Computing zeros more efficiently
  - ▶ Current best algorithms are polynomial in  $N$
  - ▶ Would be great to get methods that are polynomial in  $\log(N)$ , or insensitive to  $N$  entirely
- Bounding rank analytically in an efficient way for given curves
  - ▶ Improve work of Bober et al

# Future Work

- Generalize to modular  $L$ -functions of arbitrary weight + twist
- Computing zeros more efficiently
  - ▶ Current best algorithms are polynomial in  $N$
  - ▶ Would be great to get methods that are polynomial in  $\log(N)$ , or insensitive to  $N$  entirely
- Bounding rank analytically in an efficient way for given curves
  - ▶ Improve work of Bober et al
- Location of the first nontrivial zero
  - ▶ No theorems currently exist giving bounds on the location of the first nontrivial zero in terms of  $N$ ,  $r$  etc.

# Future Work

- Generalize to modular  $L$ -functions of arbitrary weight + twist
- Computing zeros more efficiently
  - ▶ Current best algorithms are polynomial in  $N$
  - ▶ Would be great to get methods that are polynomial in  $\log(N)$ , or insensitive to  $N$  entirely
- Bounding rank analytically in an efficient way for given curves
  - ▶ Improve work of Bober et al
- Location of the first nontrivial zero
  - ▶ No theorems currently exist giving bounds on the location of the first nontrivial zero in terms of  $N$ ,  $r$  etc.
- Computing conductor efficiently via analytic methods?

Ngiyabonga Kakhulu

Ngiyabonga Kakhulu

Hamba Kahle!