

The Bite of an Elliptic Curve

Simon Spicer

University of Washington

mlungu@uw.edu

April 21, 2015

Definition of the Bite

Let E/\mathbb{Q} be an elliptic curve with conductor N_E and L -series $L(E, s) := L_E(s)$.

Definition of the Bite

Let E/\mathbb{Q} be an elliptic curve with conductor N_E and L -series $L(E, s) := L_E(s)$.

Recall:

- Euler product definition of $L_E(s)$:

$$L_E(s) := \prod_{p|N_E} \frac{1}{1 - a_p p^{-s}} \prod_{p \nmid N_E} \frac{1}{1 - a_p p^{-s} + p^{1-2s}} = \sum_{n=1}^{\infty} a_n n^{-s},$$

converges for $\Re(s) > \frac{3}{2}$.

Definition of the Bite

Let E/\mathbb{Q} be an elliptic curve with conductor N_E and L -series $L(E, s) := L_E(s)$.

Recall:

- Euler product definition of $L_E(s)$:

$$L_E(s) := \prod_{p|N_E} \frac{1}{1 - a_p p^{-s}} \prod_{p \nmid N_E} \frac{1}{1 - a_p p^{-s} + p^{1-2s}} = \sum_{n=1}^{\infty} a_n n^{-s},$$

converges for $\Re(s) > \frac{3}{2}$.

- Modularity Theorem (Breuil, Conrad, Diamond, Taylor, Wiles):
 $L_E(s)$ admits analytic continuation to \mathbb{C} .

Definition of the Bite

Let E/\mathbb{Q} be an elliptic curve with conductor N_E and L -series $L(E, s) := L_E(s)$.

Recall:

- Euler product definition of $L_E(s)$:

$$L_E(s) := \prod_{p|N_E} \frac{1}{1 - a_p p^{-s}} \prod_{p \nmid N_E} \frac{1}{1 - a_p p^{-s} + p^{1-2s}} = \sum_{n=1}^{\infty} a_n n^{-s},$$

converges for $\Re(s) > \frac{3}{2}$.

- Modularity Theorem (Breuil, Conrad, Diamond, Taylor, Wiles): $L_E(s)$ admits analytic continuation to \mathbb{C} .
- $L_E(s)$ can be evaluated to k bits precision in $\tilde{O}(k \cdot \sqrt{N_E})$ time.

The Zeros of $L_E(s)$

Three flavors:

- A simple zero at $0, -1, -2, -3, \dots$
- A zero of order r_{an} at $s = 1$; r_{an} is called the *analytic rank* of E
- Countably infinite zeros in the strip $0 < \Re(s) < 2$, symmetric about $\Re(s) = 1$ and x-axis.

The Zeros of $L_E(s)$

Three flavors:

- A simple zero at $0, -1, -2, -3, \dots$
- A zero of order r_{an} at $s = 1$; r_{an} is called the *analytic rank* of E
- Countably infinite zeros in the strip $0 < \Re(s) < 2$, symmetric about $\Re(s) = 1$ and χ -axis.

Conjecture (Generalized Riemann Hypothesis for Elliptic Curves)

All nontrivial zeros of $L_E(s)$ are simple and lie on the line $\Re(s) = 1$.

The Zeros of $L_E(s)$

Three flavors:

- A simple zero at $0, -1, -2, -3, \dots$
- A zero of order r_{an} at $s = 1$; r_{an} is called the *analytic rank* of E
- Countably infinite zeros in the strip $0 < \Re(s) < 2$, symmetric about $\Re(s) = 1$ and x -axis.

Conjecture (Generalized Riemann Hypothesis for Elliptic Curves)

All nontrivial zeros of $L_E(s)$ are simple and lie on the line $\Re(s) = 1$.

GRH always assumed in this talk.

The Zeros of $L_E(s)$

Three flavors:

- A simple zero at $0, -1, -2, -3, \dots$
- A zero of order r_{an} at $s = 1$; r_{an} is called the *analytic rank* of E
- Countably infinite zeros in the strip $0 < \Re(s) < 2$, symmetric about $\Re(s) = 1$ and x-axis.

Conjecture (Generalized Riemann Hypothesis for Elliptic Curves)

All nontrivial zeros of $L_E(s)$ are simple and lie on the line $\Re(s) = 1$.

GRH always assumed in this talk.




Figure: The zeros of $L_E(s)$ for $E = 37a$

Definition of the Bite

Definition of the Bite

Easier to work with shifted L -function $L_E(1 + s)$.

Definition of the Bite

Easier to work with shifted L -function $L_E(1 + s)$.

$\implies L_E(1 + s)$ may or may not have zeros at the origin.

Definition of the Bite

Easier to work with shifted L -function $L_E(1+s)$.

$\implies L_E(1+s)$ may or may not have zeros at the origin.

BSD:

$$r_{al}(E) = r_{an}(E) := \text{ord}_{s=0} L_E(1+s)$$

Definition of the Bite

Easier to work with shifted L -function $L_E(1+s)$.

$\implies L_E(1+s)$ may or may not have zeros at the origin.

BSD:

$$r_{al}(E) = r_{an}(E) := \text{ord}_{s=0} L_E(1+s)$$

Definition

The *Bite* of E is

$$\beta(E) = \beta_E := \sum_{\gamma \neq 0} \frac{1}{\gamma^2}$$

where γ ranges over the imaginary parts of **noncentral** nontrivial zeros of $L_E(s)$.

Examples of the Bite





Figure: The zeros of $L_E(s)$ for $E = 37a$


Examples of the Bite



- “Add up the inverse squares of the heights of the blue dots”

Figure: The zeros of $L_E(s)$ for $E = 37a$


Examples of the Bite



- “Add up the inverse squares of the heights of the blue dots”
- Can use Rubinstein's `lcalc` package to compute zeros.

Figure: The zeros of $L_E(s)$ for $E = 37a$

Examples of the Bite



- “Add up the inverse squares of the heights of the blue dots”
- Can use Rubinstein's `lcalc` package to compute zeros.
- $E = 37a : y^2 = x^3 - 16x + 16$ has nontrivial zeros at

0.000000000...

5.00317001...

6.87039122...

8.01433081...

9.93309835...

10.7751382...

Figure: The zeros of $L_E(s)$ for $E = 37a$

Examples of the Bite




Figure: The zeros of $L_E(s)$ for $E = 37a$

- “Add up the inverse squares of the heights of the blue dots”
- Can use Rubinstein's `lcalc` package to compute zeros.
- $E = 37a : y^2 = x^3 - 16x + 16$ has nontrivial zeros at

0.000000000...

5.00317001...

6.87039122...


8.01433081...

9.93309835...


10.7751382...

• $\implies \beta_E = 0.3792\dots$

Examples of the Bite



Examples of the Bite



- Smallest bite: $E = 11a$, $\beta_E = 0.2551\dots$
- Largest bite: $E = 256944c$, $\beta_E = 3056.1912\dots$

Why is the Bite interesting?

Why is the Bite interesting?

β_E appears in higher-order coefficients of the Taylor expansion of $L_E(s)$ about the central point:

Why is the Bite interesting?

β_E appears in higher-order coefficients of the Taylor expansion of $L_E(s)$ about the central point:

Proposition

Let E/\mathbb{Q} have conductor N_E , L -function $L_E(s)$ with bite β_E and analytic rank r_E . Let $L_E(1+s)$ have Taylor expansion

$$L_E(1+s) = C \cdot s^{r_E} \cdot [1 + a \cdot s + b \cdot s^2 + O(s^3)]$$

Then

$$a = - \left[-\eta + \log \left(\frac{\sqrt{N_E}}{2\pi} \right) \right]$$
$$2b = \left[-\eta + \log \left(\frac{\sqrt{N_E}}{2\pi} \right) \right]^2 - \frac{\pi^2}{6} + \beta_E,$$

where η is the Euler-Mascheroni constant $= 0.5772\dots$

Why is the Bite interesting?

Proof.

- Define completed L -function:

$$\Lambda_E(1+s) := (N_E)^{\frac{1+s}{2}} (2\pi)^{-1-s} \Gamma(1+s) L_E(1+s)$$

Why is the Bite interesting?

Proof.

- Define completed L -function:

$$\Lambda_E(1+s) := (N_E)^{\frac{1+s}{2}} (2\pi)^{-1-s} \Gamma(1+s) L_E(1+s)$$

- Hadamard product expansion for $\Lambda_E(1+s)$:

$$\Lambda_E(1+s) = C \cdot s^r \cdot \prod_{\gamma \neq 0} \left(1 + \frac{s^2}{\gamma^2}\right)$$

$$\text{So } \Lambda_E(1+s) = C \cdot s^r \cdot (1 + \beta \cdot s^2 + O(s^4))$$

Why is the Bite interesting?

Proof.

- Define completed L -function:

$$\Lambda_E(1+s) := (N_E)^{\frac{1+s}{2}} (2\pi)^{-1-s} \Gamma(1+s) L_E(1+s)$$

- Hadamard product expansion for $\Lambda_E(1+s)$:

$$\Lambda_E(1+s) = C \cdot s^r \cdot \prod_{\gamma \neq 0} \left(1 + \frac{s^2}{\gamma^2}\right)$$

$$\text{So } \Lambda_E(1+s) = C \cdot s^r \cdot (1 + \beta \cdot s^2 + O(s^4))$$

- Gamma function has nice expansion about $s = 1$:

$$\Gamma(1+s) = 1 - \eta \cdot s + \frac{1}{12} (6\eta^2 + \pi^2) \cdot s^2 + O(s^3)$$

Why is the Bite interesting?

Proof.

- Define completed L -function:

$$\Lambda_E(1+s) := (N_E)^{\frac{1+s}{2}} (2\pi)^{-1-s} \Gamma(1+s) L_E(1+s)$$

- Hadamard product expansion for $\Lambda_E(1+s)$:

$$\Lambda_E(1+s) = C \cdot s^r \cdot \prod_{\gamma \neq 0} \left(1 + \frac{s^2}{\gamma^2}\right)$$

$$\text{So } \Lambda_E(1+s) = C \cdot s^r \cdot (1 + \beta \cdot s^2 + O(s^4))$$

- Gamma function has nice expansion about $s = 1$:

$$\Gamma(1+s) = 1 - \eta \cdot s + \frac{1}{12} (6\eta^2 + \pi^2) \cdot s^2 + O(s^3)$$

- Multiply out expansions and collect terms (tediously!)



Computing the Bite efficiently

This gives us a method to compute the bite of a curve efficiently if we know the analytic rank.

Computing the Bite efficiently

This gives us a method to compute the bite of a curve efficiently if we know the analytic rank.

Corollary

For E with conductor N and analytic rank r ,

$$\beta(E) = \frac{2}{(r+1)(r+2)} \cdot \frac{L_E^{(r+2)}(1)}{L_E^{(r)}(1)} - \left[-\eta + \log \left(\frac{\sqrt{N}}{2\pi} \right) \right]^2 + \frac{\pi^2}{6}$$

Computing the Bite efficiently

This gives us a method to compute the bite of a curve efficiently if we know the analytic rank.





Corollary

For E with conductor N and analytic rank r ,

$$\beta(E) = \frac{2}{(r+1)(r+2)} \cdot \frac{L_E^{(r+2)}(1)}{L_E^{(r)}(1)} - \left[-\eta + \log \left(\frac{\sqrt{N}}{2\pi} \right) \right]^2 + \frac{\pi^2}{6}$$

- Bites for all Cremona curves were computed using the above formula
- ± 1 week computation time on SMC.

The average size of β_E



The average size of β_E

The average size of β_E

- Clearly β_E grows on average with conductor.

The average size of β_E

- Clearly β_E grows on average with conductor.
- It would be nice to have upper and lower bounds on β_E

The average size of β_E

- Clearly β_E grows on average with conductor.
- It would be nice to have upper and lower bounds on β_E
- Lower bounds easier than upper bounds.

Zero Density

Define the zero counting function

$$M_E(t) = \#\{\gamma : |\gamma| \leq t\},$$

i.e. the number of zeros with imaginary part up to t

Zero Density

Define the zero counting function

$$M_E(t) = \#\{\gamma : |\gamma| \leq t\},$$

i.e. the number of zeros with imaginary part up to t

Proposition

$$M_E(t) = \frac{t \log t}{\pi} + \frac{t \log N_E}{2\pi} - \frac{t \log 2\pi e}{\pi} + \frac{1}{4} + O(\log t)$$

where the error term is positive as often as it negative and contributes no net bias.

Zero Density

Define the zero counting function

$$M_E(t) = \#\{\gamma : |\gamma| \leq t\},$$

i.e. the number of zeros with imaginary part up to t

Proposition

$$M_E(t) = \frac{t \log t}{\pi} + \frac{t \log N_E}{2\pi} - \frac{t \log 2\pi e}{\pi} + \frac{1}{4} + O(\log t)$$

where the error term is positive as often as it negative and contributes no net bias.

- Zero density grows with $\log N_E$

Zero Density

Define the zero counting function

$$M_E(t) = \#\{\gamma : |\gamma| \leq t\},$$

i.e. the number of zeros with imaginary part up to t

Proposition


$$M_E(t) = \frac{t \log t}{\pi} + \frac{t \log N_E}{2\pi} - \frac{t \log 2\pi e}{\pi} + \frac{1}{4} + O(\log t)$$

where the error term is positive as often as it negative and contributes no net bias.



- Zero density grows with $\log N_E$
- Expect β_E to grow with $\log N_E$ at least

Zero Density Example

Zero Density Example



Zero Density Example



Lower Bounds on β_E

- $$M_E(t) = \frac{t \log t}{\pi} + \frac{t \log N_E}{2\pi} - \frac{t \log 2\pi e}{\pi} + \frac{1}{4} + O(\log t)$$

Lower Bounds on β_E

- $M_E(t) = \frac{t \log t}{\pi} + \frac{t \log N_E}{2\pi} - \frac{t \log 2\pi e}{\pi} + \frac{1}{4} + O(\log t)$
- Average zero density grows with $\log N_E$

Lower Bounds on β_E

- $M_E(t) = \frac{t \log t}{\pi} + \frac{t \log N_E}{2\pi} - \frac{t \log 2\pi e}{\pi} + \frac{1}{4} + O(\log t)$
- Average zero density grows with $\log N_E$
- Expect β_E to grow proportional to $\log N_E$ at least

Lower Bounds on β_E

- $M_E(t) = \frac{t \log t}{\pi} + \frac{t \log N_E}{2\pi} - \frac{t \log 2\pi e}{\pi} + \frac{1}{4} + O(\log t)$
- Average zero density grows with $\log N_E$
- Expect β_E to grow proportional to $\log N_E$ at least

Theorem (S.)

For all $\epsilon > 0$ there is a constant $K(\epsilon) > 0$ such that for all elliptic curves E , β_E obeys

$$\beta_E > \frac{1}{1 + \epsilon} \log N_E - K(\epsilon).$$

A Lemma Regarding β_E

Lemma (1)

For any $\sigma > \frac{1}{2}$, the bite β_E and analytic rank r_E of a curve E obey

$$\sigma \cdot \beta_E + \frac{r_E}{\sigma} > \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma}(1 + \sigma) - \log(2\pi) - 2\frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right)$$

A Lemma Regarding β_E

Lemma (1)

For any $\sigma > \frac{1}{2}$, the bite β_E and analytic rank r_E of a curve E obey

$$\sigma \cdot \beta_E + \frac{r_E}{\sigma} > \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma}(1 + \sigma) - \log(2\pi) - 2\frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right)$$

Proof.

- Take logarithmic derivative of $\Lambda_E(1+s)$ to get

$$\frac{\Lambda'_E}{\Lambda_E}(1+s) = \sum_{\gamma} \frac{s}{s^2 + \gamma^2} = \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma}(1+s) - \log(2\pi) + \frac{L'_E}{L_E}(1+s)$$

A Lemma Regarding β_E

Lemma (1)

For any $\sigma > \frac{1}{2}$, the bite β_E and analytic rank r_E of a curve E obey

$$\sigma \cdot \beta_E + \frac{r_E}{\sigma} > \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma}(1 + \sigma) - \log(2\pi) - 2\frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right)$$

Proof.

- Take logarithmic derivative of $\Lambda_E(1+s)$ to get

$$\frac{\Lambda'_E}{\Lambda_E}(1+s) = \sum_{\gamma} \frac{s}{s^2 + \gamma^2} = \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma}(1+s) - \log(2\pi) + \frac{L'_E}{L_E}(1+s)$$

- $\left| \frac{L'_E}{L_E}(1+s) \right| < -2\frac{\zeta'}{\zeta} \left(\frac{1}{2} + \Re(s) \right)$ from Hasse bound on Dirichlet coefficients of $\frac{\Lambda'_E}{\Lambda_E}(1+s)$



A Lemma Regarding β_E

Lemma (1)

$$\sigma \cdot \beta_E + \frac{r_E}{\sigma} > \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma} (1 + \sigma) - \log(2\pi) - 2 \frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right) \text{ for } \sigma > \frac{1}{2}$$

Proof.

- $\frac{\Lambda'_E}{\Lambda_E} (1 + s) = \sum_{\gamma} \frac{s}{s^2 + \gamma^2} = \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma} (1 + s) - \log(2\pi) + \frac{L'_E}{L_E} (1 + s)$
- $\left| \frac{L'_E}{L_E} (1 + s) \right| < -2 \frac{\zeta'}{\zeta} \left(\frac{1}{2} + \Re(s) \right)$
- At $s = \sigma$,

$$\sum_{\gamma} \frac{\sigma}{\sigma^2 + \gamma^2} = \frac{1}{\sigma} \left(r_E + \sum_{\gamma \neq 0} \frac{1}{1 + \frac{\gamma^2}{\sigma^2}} \right) < \frac{1}{\sigma} (r_E + \sigma^2 \cdot \beta_E)$$

A Lemma Regarding β_E

Lemma (1)

$$\sigma \cdot \beta_E + \frac{r_E}{\sigma} > \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma} (1 + \sigma) - \log(2\pi) - 2\frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma\right) \text{ for } \sigma > \frac{1}{2}$$

Proof.

- $\frac{\Lambda'_E}{\Lambda_E} (1 + s) = \sum_{\gamma} \frac{s}{s^2 + \gamma^2} = \frac{1}{2} \log N_E + \frac{\Gamma'}{\Gamma} (1 + s) - \log(2\pi) + \frac{L'_E}{L_E} (1 + s)$
- $\left| \frac{L'_E}{L_E} (1 + s) \right| < -2\frac{\zeta'}{\zeta} \left(\frac{1}{2} + \Re(s)\right)$
- At $s = \sigma$,

$$\sum_{\gamma} \frac{\sigma}{\sigma^2 + \gamma^2} = \frac{1}{\sigma} \left(r_E + \sum_{\gamma \neq 0} \frac{1}{1 + \frac{\gamma^2}{\sigma^2}} \right) < \frac{1}{\sigma} (r_E + \sigma^2 \cdot \beta_E)$$

- Combine inequalities and simplify



Bounding Analytic Rank in terms of Conductor

Lemma (2)

r_E grows more slowly than $\log N_E$, i.e. for any $\epsilon > 0$ there exists $K(\epsilon)$ s.t. for any E ,

$$r_E < \epsilon \cdot \log N_E + K(\epsilon)$$

Bounding Analytic Rank in terms of Conductor

Lemma (2)

r_E grows more slowly than $\log N_E$, i.e. for any $\epsilon > 0$ there exists $K(\epsilon)$ s.t. for any E ,

$$r_E < \epsilon \cdot \log N_E + K(\epsilon)$$

Two sentence proof sketch

From previous talk: For any $\Delta > 0$,

$$\sum_{\gamma} \text{sinc}^2(\Delta\gamma) = \sum_{\gamma} \left(\frac{\sin(\Delta\gamma)}{\Delta\gamma} \right)^2 = \frac{1}{\pi\Delta} \log N_E + S(E, \Delta)$$

where $S(E, \Delta)$ is a finite sum with global bound that grows with Δ but is independent of E .

Bounding Analytic Rank in terms of Conductor

Lemma (2)

r_E grows more slowly than $\log N_E$, i.e. for any $\epsilon > 0$ there exists $K(\epsilon)$ s.t. for any E ,

$$r_E < \epsilon \cdot \log N_E + K(\epsilon)$$

Two sentence proof sketch

From previous talk: For any $\Delta > 0$,

$$\sum_{\gamma} \operatorname{sinc}^2(\Delta\gamma) = \sum_{\gamma} \left(\frac{\sin(\Delta\gamma)}{\Delta\gamma} \right)^2 = \frac{1}{\pi\Delta} \log N_E + S(E, \Delta)$$

where $S(E, \Delta)$ is a finite sum with global bound that grows with Δ but is independent of E .

And $\sum_{\gamma} \operatorname{sinc}^2(\Delta\gamma) > r_E$ always. So let $\Delta \rightarrow \infty$.

Lower Bounds on β_E

Theorem (S.)

$$\beta_E > \frac{1}{1+\epsilon} \log N_E - K(\epsilon) \quad \text{for any } \epsilon > 0$$

Lower Bounds on β_E

Theorem (S.)

$$\beta_E > \frac{1}{1+\epsilon} \log N_E - K(\epsilon) \quad \text{for any } \epsilon > 0$$

Proof.

- From Lemma 1, for any $\sigma > \frac{1}{2}$

$$\beta_E > \frac{1}{2\sigma} \log N_E - \frac{r_E}{\sigma^2} + \frac{1}{\sigma} \left[\frac{\Gamma'}{\Gamma} (1 + \sigma) - \log(2\pi) + 2 \frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right) \right]$$

Lower Bounds on β_E

Theorem (S.)

$$\beta_E > \frac{1}{1+\epsilon} \log N_E - K(\epsilon) \quad \text{for any } \epsilon > 0$$

Proof.

- From Lemma 1, for any $\sigma > \frac{1}{2}$

$$\beta_E > \frac{1}{2\sigma} \log N_E - \frac{r_E}{\sigma^2} + \frac{1}{\sigma} \left[\frac{\Gamma'}{\Gamma} (1 + \sigma) - \log(2\pi) + 2 \frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right) \right]$$

- Choose $\sigma = \frac{1}{2} + \frac{\epsilon}{2}$

Lower Bounds on β_E

Theorem (S.)

$$\beta_E > \frac{1}{1+\epsilon} \log N_E - K(\epsilon) \quad \text{for any } \epsilon > 0$$

Proof.

- From Lemma 1, for any $\sigma > \frac{1}{2}$

$$\beta_E > \frac{1}{2\sigma} \log N_E - \frac{r_E}{\sigma^2} + \frac{1}{\sigma} \left[\frac{\Gamma'}{\Gamma} (1 + \sigma) - \log(2\pi) + 2 \frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right) \right]$$

- Choose $\sigma = \frac{1}{2} + \frac{\epsilon}{2}$
- Bound r_E by $\epsilon^2 \log N_E + K'(\epsilon^2)$

Lower Bounds on β_E

Theorem (S.)

$$\beta_E > \frac{1}{1+\epsilon} \log N_E - K(\epsilon) \quad \text{for any } \epsilon > 0$$

Proof.

- From Lemma 1, for any $\sigma > \frac{1}{2}$

$$\beta_E > \frac{1}{2\sigma} \log N_E - \frac{r_E}{\sigma^2} + \frac{1}{\sigma} \left[\frac{\Gamma'}{\Gamma} (1 + \sigma) - \log(2\pi) + 2 \frac{\zeta'}{\zeta} \left(\frac{1}{2} + \sigma \right) \right]$$

- Choose $\sigma = \frac{1}{2} + \frac{\epsilon}{2}$
- Bound r_E by $\epsilon^2 \log N_E + K'(\epsilon^2)$
- Combine inequalities and let $\epsilon \rightarrow 0$



Issues with bounding β_E from below

- Can we show lower bound β_E grows faster than $\log N_E$?

Issues with bounding β_E from below

- Can we show lower bound β_E grows faster than $\log N_E$?
- If zero density $\sim \log N_E$, then we expect $\beta_E = \sum \gamma^{-2} \sim (\log N_E)^2$.

Issues with bounding β_E from below

- Can we show lower bound β_E grows faster than $\log N_E$?
- If zero density $\sim \log N_E$, then we expect $\beta_E = \sum \gamma^{-2} \sim (\log N_E)^2$.
- Problem: Low-lying zeros not well understood

Issues with bounding β_E from below

- Can we show lower bound β_E grows faster than $\log N_E$?
- If zero density $\sim \log N_E$, then we expect $\beta_E = \sum \gamma^{-2} \sim (\log N_E)^2$.
- Problem: Low-lying zeros not well understood
 - ▶ Sarnak: lowest zero approaches constant distribution as $N_E \rightarrow \infty$
 - ▶ \implies bite grows more slowly than $(\log N_E)^2$

Upper bounds on β_E

Corollary to efficient bite formula

For E with analytic rank r_E , bite β_E and (completed L -function) leading central Taylor coefficient C_E ,

$$\beta_E \cdot C_E = \frac{\Lambda_E^{(r_E+2)}(1)}{(r_E + 2)!}$$

where $\Lambda_E(1 + s)$ is the completed L -function for E .

Upper bounds on β_E

Corollary to efficient bite formula

For E with analytic rank r_E , bite β_E and (completed L -function) leading central Taylor coefficient C_E ,

$$\beta_E \cdot C_E = \frac{\Lambda_E^{(r_E+2)}(1)}{(r_E + 2)!}$$

where $\Lambda_E(1 + s)$ is the completed L -function for E .

- \implies The bite and the leading central Taylor coefficient are intimately linked

Upper bounds on β_E

Corollary to efficient bite formula





For E with analytic rank r_E , bite β_E and (completed L -function) leading central Taylor coefficient C_E ,

$$\beta_E \cdot C_E = \frac{\Lambda_E^{(r_E+2)}(1)}{(r_E + 2)!}$$





where $\Lambda_E(1 + s)$ is the completed L -function for E .

- \implies The bite and the leading central Taylor coefficient are intimately linked
- Behavior of $\beta_E \cdot C_E$ heavily constrained

The bite times the BSD coefficient



The bite times the BSD coefficient



Issues with bounding β_E from above

- β_E and C_E cannot both simultaneously be too big or too small

Issues with bounding β_E from above

- β_E and C_E cannot both simultaneously be too big or too small
- β_E is large when $L_E(s)$ has low-lying lowest zero γ_0

Issues with bounding β_E from above

- β_E and C_E cannot both simultaneously be too big or too small
- β_E is large when $L_E(s)$ has low-lying lowest zero γ_0
-

$$\begin{aligned}\{\text{Bounding } \beta_E \text{ from above}\} &\iff \{\text{Bounding } \gamma_0 \text{ from below}\} \\ &\iff \{\text{Bounding } C_E \text{ from below}\}\end{aligned}$$

Computational equivalence of β_E and r_E

- Knowing β_E is computationally equivalent to knowing r_E :

Computational equivalence of β_E and r_E

- Knowing β_E is computationally equivalent to knowing r_E :
- $$\beta_E = \frac{2}{(r_E+1)(r_E+2)} \cdot \frac{L_E^{(r_E+2)}(1)}{L_E^{(r_E)}(1)} - \left[-\eta + \log \left(\frac{\sqrt{N_E}}{2\pi} \right) \right]^2 + \frac{\pi^2}{6}$$

Computational equivalence of β_E and r_E

- Knowing β_E is computationally equivalent to knowing r_E :
- $$\beta_E = \frac{2}{(r_E+1)(r_E+2)} \cdot \frac{L_E^{(r_E+2)}(1)}{L_E^{(r_E)}(1)} - \left[-\eta + \log \left(\frac{\sqrt{N_E}}{2\pi} \right) \right]^2 + \frac{\pi^2}{6}$$

Proposition

The analytic rank of E is the largest integer less than

$$\frac{1}{\sqrt{\beta_E}} \left[\left(-\eta + \log \left(\frac{\sqrt{N_E}}{2\pi} \right) \right) + \frac{1}{2\sqrt{\beta_E}} \left(\frac{\pi^2}{6} - Li_2 \left(e^{-2\sqrt{\beta_E}} \right) \right) \right. \\ \left. + \sum_{\log n < 2\sqrt{\beta_E}} c_n \cdot \left(1 - \frac{\log n}{2\sqrt{\beta_E}} \right) \right] \quad \text{where}$$

- $\eta = 0.5772\dots$ is the Euler-Mascheroni constant
- $Li_2(s)$ is the dilogarithm function on \mathbb{C}
- $c_n = c_n(E)$ is the n th Dirichlet coefficient of $\frac{L'_E}{L_E}(1+s)$

Another reason β_E is interesting

- The bite controls convergence rates for explicit formula-type sums.

Another reason β_E is interesting

- The bite controls convergence rates for explicit formula-type sums.
- Mazur: BSD + GRH implies

$$r_{al}(E) = \lim_{X \rightarrow \infty} \frac{1}{\log X} \left[\sum_{p < X} \frac{-a_p(E) \log p}{p} \right]$$

Another reason β_E is interesting

- The bite controls convergence rates for explicit formula-type sums.
- Mazur: BSD + GRH implies

$$r_{al}(E) = \lim_{X \rightarrow \infty} \frac{1}{\log X} \left[\sum_{p < X} \frac{-a_p(E) \log p}{p} \right]$$

- Convergence rate of this sum should scale with $e^{\sqrt{\beta_E}}$

Open Questions

Open Questions

- Can we come up with better lower and upper bounds on β_E ?

Open Questions

- Can we come up with better lower and upper bounds on β_E ?
- Can we say anything about the mean value of β_E ?

Open Questions

- Can we come up with better lower and upper bounds on β_E ?
- Can we say anything about the mean value of β_E ?
- Is there a way to compute β_E without knowing r_E or C_E ?

Open Questions

- Can we come up with better lower and upper bounds on β_E ?
- Can we say anything about the mean value of β_E ?
- Is there a way to compute β_E without knowing r_E or C_E ?
- Is there any arithmetic significance to β_E ?

Thank You