

CHAPTER 2: 2ND ORDER DE'S§2.0: INTRO

Boyce 3.1

Definition 2.0.1: • A 2nd-order ODE is an equation of the form

$$\frac{d^2 y}{dt^2} = f(t, y, \frac{dy}{dt})$$

Remark: If $f(t, y, \frac{dy}{dt})$ is of the form

$$f(t, y, \frac{dy}{dt}) = g(t) - p(t) \frac{dy}{dt} - q(t) \cdot y,$$

then the DE is said to be linear;

in this case we can write the linear ODE in standard form:

$$\frac{d^2 y}{dt^2} + p(t) \frac{dy}{dt} + q(t) \cdot y = g(t).$$

Occasionally we'll see linear 2nd-order DEs in the form

$$P(t) \frac{d^2 y}{dt^2} + Q(t) \frac{dy}{dt} + R(t) \cdot y = G(t)$$

- If the 2nd-order ODE is not writeable in one of the above 2 forms, it is called nonlinear.

In general, ^{solving} nonlinear 2nd order ODEs is very hard, or impossible to do analytically, and requiring doing so requires a more advanced set of techniques beyond the scope of this course.

In MATH 307 we will restrict ourselves to solving linear 2nd-order ODEs; these, however, comprise a large percentage of the 2nd-order DEs we see in the real world.

§2.1: Homogeneous Equations
with constant coefficients

Definition 2.1.1 A 2nd-order linear DE $P(t)y'' + Q(t)y' + R(t)y = G(t)$ is called homogeneous if $G(t) = 0$ for all t .

It is called nonhomogeneous if $G(t) \neq 0$.

We'll consider non-homogeneous equations in a later section.

The simplest homogeneous equations we can consider are when $P(t)$, $Q(t)$ & $R(t)$ are all constant, i.e.

$$ay'' + by' + cy = 0$$

for constants a, b, c ($a \neq 0$); these DEs are always solvable in terms of elementary functions of t .

Example 3.1.2 Note 3.1.2

In general, when solving a 2nd-order DE we'll end up integrating twice: the result is that we introduce 2 independent free constants in the general solution. We therefore need a set of two conditions to specify a particular solution to a 2nd-order DE.

Definition 3.1.3 An initial value problem for a 2nd-order DE is a 2nd-order differential equation along with a pair of conditions, usually of the form

$$y(t_0) = y_0 \text{ and } y'(t_0) = y'_0$$

Sometimes we'll also see ICs in the form

$$y(t_0) = y_0 \text{ and } y(t_1) = y_1$$

Example 3.1.4 Solve the IVP

$$y'' - y = 0, \quad y(0) = 2, \quad y'(0) = -1.$$

This is a linear homogeneous 2nd-order ODE with constant coefficients. We have $y'' = y$; 2 functions that satisfy this DE immediately spring to mind:

$$y = e^t \text{ and } y = e^{-t}$$

Note, however, that any multiple of e^t also satisfies the DE, as does any multiple of e^{-t} .

Now note that we can add e^t to e^{-t} and get another solution: $y = e^t + e^{-t}$ obeys the DE.

In fact, any multiple of e^t can be added to any multiple of e^{-t} , and we still have a solution to $y'' - y = 0$.

Thus a general solution to $y'' - y = 0$ is $y = c_1 e^t + c_2 e^{-t}$

Now we can apply ICs:

$$y(0) = 2 \Rightarrow 2 = c_1 + c_2$$

$$\text{And } y'(t) = c_1 e^t - c_2 e^{-t}, \text{ so } y'(0) = -1 \\ = -1 = c_1 - c_2$$

Solving simultaneously for c_1 & c_2 yields $c_1 = \frac{1}{2}, c_2 = \frac{3}{2}$

So we get

$$y = \frac{1}{2} e^t + \frac{3}{2} e^{-t}$$

□

How do we solve the general case $ay'' + by' + cy = 0$?

Tactic 2.1.5: "Guess" a solution $y = e^{rt}$ for some r .
Then $y' = r e^{rt}$ & $y'' = r^2 e^{rt}$,
So $ay'' + by' + cy = ar^2 e^{rt} + br e^{rt} + c e^{rt} = (ar^2 + br + c) e^{rt}$

Now for this to $= 0$, since e^{rt} is never zero, we must have $ar^2 + br + c = 0$

Definition 2.1.6 The equation $ar^2 + br + c = 0$ is called the characteristic equation of the DE

So if r is a number such that $ar^2 + br + c = 0$, then $y = e^{rt}$ will satisfy the DE.

Note 2.1.7 $ar^2 + br + c = 0$ is a quadratic equation, which may have 2 real but different roots, 2 real & equal roots, or 2 complex but different roots

pro

The behaviour of the solutions to the DE turn out to depend heavily on what type of roots the characteristic polynomial have.

Today we'll consider the case where $ar^2+br+c=0$ has real & different roots; the other cases we consider in later sections.

So let r_1 & r_2 be the real roots to $ar^2+br+c=0$, with $r_1 \neq r_2$. Note that $y = e^{r_1 t}$ and $y = e^{r_2 t}$ are both solutions to $ay''+by'+cy=0$.

In fact, so is any linear combination of $e^{r_1 t}$ & $e^{r_2 t}$, i.e. $y = c_1 e^{r_1 t} + c_2 e^{r_2 t}$ is a general solution to the DE. Note that we can show that this is the full general solution to the DE.

Exercise:

If you have ICs $y(t_0) = y_0$, $y'(t_0) = y'_0$, show that the unique solution to the IVP is

$$y = c_1 e^{r_1 t} + c_2 e^{r_2 t}, \text{ where } c_1 = \frac{y'_0 r_1 - y_0 r_2}{r_1 - r_2} e^{-r_1 t_0}, c_2 = \frac{y_0 r_1 - y'_0 r_2}{r_1 - r_2} e^{-r_2 t_0}$$

Example 2.1.8 Find the General solution to $y''+5y'+6y=0$.

\Rightarrow Suppose $y = e^{rt}$ solves the DE

$$\text{Then } 0 = y'' + 5y' + 6y = r^2 e^{rt} + 5r e^{rt} + 6e^{rt} = (r^2 + 5r + 6)e^{rt},$$

$$\text{so } r^2 + 5r + 6 = 0$$

$$\text{Thus } r = -2 \text{ or } r = -3.$$

The general solution is thus

$$y = c_1 e^{-2t} + c_2 e^{-3t}$$

Example 2.1.9 Solve the IVP: $y''+5y'+6y=0$, $y(0)=2$, $y'(0)=3$

We have that $y = c_1 e^{-2t} + c_2 e^{-3t}$.


$$\text{Setting } y(0)=2 \Rightarrow 2 = c_1 + c_2$$

$$\text{And } y'(0)=3 \Rightarrow 3 = -2c_1 - 3c_2$$

Solving gets $c_1 = 9$, $c_2 = -7$.

Hence the solution is $y = 9e^{-2t} - 7e^{-3t}$.

Plot of the solution:



Example 2.1.10 Find the solution to $4y'' - 8y' + 3y = 0$, $y(0) = 2$
 And compute the t for which it is a max. $y'(0) = \frac{1}{2}$.

$$\text{get } 4r^2 - 8r + 3 = 0$$

$$\Rightarrow r = \frac{3}{2} \text{ or } r = \frac{1}{2}$$

$$\text{So } y = c_1 e^{\frac{3}{2}t} + c_2 e^{\frac{1}{2}t}$$


$$\text{Ics: } y(0) = 2 \Rightarrow c_1 + c_2 = 2$$

$$y'(0) = \frac{1}{2} \Rightarrow \frac{3}{2}c_1 + \frac{1}{2}c_2 = \frac{1}{2}$$

$$\Rightarrow c_1 = -\frac{1}{2}, c_2 = \frac{5}{2}$$

$$\text{So } y = -\frac{1}{2}e^{-\frac{3}{2}t} + \frac{5}{2}e^{\frac{1}{2}t}$$

Plot



$$\text{Max occurs when } y' = 0 \Rightarrow -\frac{3}{4}e^{\frac{3}{2}t} + \frac{5}{4}e^{\frac{1}{2}t} = 0$$

$$\Rightarrow 5e^{\frac{1}{2}t} = 3e^{\frac{3}{2}t}$$

$$\Rightarrow e^t = \frac{5}{3} \Rightarrow t = \ln\left(\frac{5}{3}\right)$$

$$\text{So max } y \text{ is } -\frac{1}{2}\left(\frac{5}{3}\right)^{\frac{3}{2}} + \frac{5}{2}\left(\frac{5}{3}\right)^{\frac{1}{2}}$$

Summary: When considering the DE $ay'' + by' + cy = 0$,
a, b, c constant, the ^{general} solution can have the following behaviors:

- If $ar^2 + br + c$ has both positive real roots,
the solution $\rightarrow \pm \infty$ as $t \rightarrow \infty$
- If " " has both negative real roots,
the solution $\rightarrow 0$ as $t \rightarrow \infty$
- If " " has one positive, one negative root,
the in general solution $\rightarrow \pm \infty$ as $t \rightarrow \infty$; however, some
specific solutions may $\rightarrow 0$. (as $t \rightarrow \infty$)
- If one root of characteristic equation $= 0$, the
solution \rightarrow a constant
(If other root is positive, the solution $\rightarrow \pm \infty$ in general)