

Relations between Inflammation, access to care and Diabetes in two representative populations of China and Mexico.

Dominik Grätz¹, Rachel Miller-Moudgil¹, Amber Somarriba¹, Brittany Spinner¹, & Tian Walker¹

¹ University of Oregon

Author Note

List of group members ordered by alphabet.

Abstract

Background. Background goes here. *Methods.* Methods go here. *Results.* Results here.

Conclusions. Conclusions here.

Keywords: Diabetes, access to care, inflammation, health, Mexico, China

Word count: X (this cannot easily be done automatically, we can also just leave it out)

Relations between Inflammation, access to care and Diabetes in two representative populations of China and Mexico.

Introduction

Enter text/code here. I make a reference here (Deshpande, Harris-Hayes, & Schootman, 2008). **This is just copied from the outline document; needs to be rewritten/references automated!**

About the data:

We will be using two datasets from the Wave 1 Study on Global Ageing (SAGE) from the World Health Organization (WHO). We will be looking at inflammation (measured through C-reactive protein; crp) and hemoglobin A1c (average blood glucose level over the last three months) to better understand the relationship between inflammation and diabetes in Mexico and China. Observations of the potential association between diabetes and inflammation were initially made over 100 years ago (Williamson, 1901), and we now know that inflammation is a pathogenic mediator in the progression of diabetes, and likely has a causal role in some cases as well (Tsalamandris et al., 2019). The relationship between inflammation and diabetes is multifaceted but we know that decreasing inflammation means decreasing diabetes progression and complications (Agrawal & Kant, 2014). The data is from two datasets designed exactly the same: one from Mexico and one from China. The data frame for China has 13,367 observations from 1,489 variables. In the dataset from Mexico there are 2,315 observations of 1690 variables. Everything else is the same as the dataset from China. Almost all of the variables are numbers or characters, and the data was originally formatted in SPSS (statistical package for the social sciences). There were more people who did the survey than did the biomarker collection (source of CRP and HbA1c data) so the number of data points in our study is smaller than the full dataset. There is demographic information along with many health-related validated scales. We have listed the variables that we will be looking at below

Why China and Mexico?

China and Mexico are interesting places to study the relationship between diabetes and inflammation because they are both classified as low to middle income countries (LMIC's) where 80% of the global burden of diabetes falls. Additionally, diabetes is a leading cause of death in Mexico and closely linked to the obesity epidemic. China experienced a famine that likely created an epigenetic environment that makes people prone to diabetes. Obesity is a relatively new phenomenon in China and so the pathophysiology of disease development may be different than other places where obesity has been a major issue for decades.

Methods

Results

Enter text/code here. Let's do all the coding here!

```
##
## Call:
## lm(formula = crp ~ hba1c + medication, data = data_dia[data_dia$country ==
##      "China", ])
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.7485 -0.9752 -0.3596  0.7661  3.2601
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   1.38413    0.20967   6.602 1.34e-10 ***
## hba1c          0.03325    0.03060   1.087  0.2778
## medication2 -0.30649    0.16159  -1.897  0.0586 .
```

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
##
```

```
## Residual standard error: 1.189 on 388 degrees of freedom
```

```
## (103 observations deleted due to missingness)
```

```
## Multiple R-squared:  0.01352,    Adjusted R-squared:  0.008432
```

```
## F-statistic: 2.658 on 2 and 388 DF,  p-value: 0.07136
```





Table 1
This is where the caption goes

predictor	b	95% CI	t(388)	p
Intercept	1.38	[0.97, 1.80]	6.60	< .001
Hba1c	0.03	[−0.03, 0.09]	1.09	.278
Medication	−0.31	[−0.62, 0.01]	−1.90	.059

Note. Model fit: $F(2, 388) = 2.66, p = 0.07, R^2 = 0.01$



```
##  
## Call:  
## lm(formula = crp ~ hba1c, data = mexico)
```

##

Residuals:

##	Min	1Q	Median	3Q	Max
##	-2.2072	-1.1830	-0.6649	0.5873	7.9933

##

Coefficients:

##		Estimate	Std. Error	t value	Pr(> t)
##	(Intercept)	1.38948	0.16425	8.459	< 2e-16 ***
##	hba1c	0.09642	0.02564	3.761	0.000173 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

##

Residual standard error: 1.808 on 2402 degrees of freedom

(3044 observations deleted due to missingness)

Multiple R-squared: 0.005854, Adjusted R-squared: 0.00544

F-statistic: 14.14 on 1 and 2402 DF, p-value: 0.0001735

##

Call:

lm(formula = crp ~ hba1c, data = china)

##

Residuals:

##	Min	1Q	Median	3Q	Max
##	-3.0866	-1.2185	-0.8918	0.3402	8.4252

##

Coefficients:

##		Estimate	Std. Error	t value	Pr(> t)
##	(Intercept)	1.13766	0.08207	13.862	< 2e-16 ***

```

## hba1c          0.10663      0.01379    7.733 1.18e-14 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 2.117 on 7547 degrees of freedom
## (7501 observations deleted due to missingness)
## Multiple R-squared:  0.007862,    Adjusted R-squared:  0.00773
## F-statistic:  59.8 on 1 and 7547 DF,  p-value: 1.185e-14
##
## Call:
## lm(formula = crp ~ hba1c + q_mean, data = data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.0621 -0.7793 -0.4305  0.4763  3.7343
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  0.698830   0.055714  12.543 < 2e-16 ***
## hba1c        0.064692   0.007052   9.174 < 2e-16 ***
## q_mean       0.164575   0.025041   6.572 5.27e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.099 on 7893 degrees of freedom
## (1128 observations deleted due to missingness)
## Multiple R-squared:  0.01633,    Adjusted R-squared:  0.01608

```



```
## F-statistic: 65.5 on 2 and 7893 DF, p-value: < 2.2e-16

##

## Call:
## lm(formula = crp ~ hba1c + q_mean, data = mexico)
##

## Residuals:

##      Min       1Q   Median       3Q      Max
## -2.1795 -1.2074 -0.6225  0.6301  7.9003

##

## Coefficients:

##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  1.20415     0.24679   4.879 1.18e-06 ***
## hba1c         0.06934     0.03297   2.103  0.03561 *
## q_mean        0.14385     0.05186   2.774  0.00561 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

##

## Residual standard error: 1.794 on 1444 degrees of freedom
## (4001 observations deleted due to missingness)
## Multiple R-squared:  0.008975, Adjusted R-squared:  0.007603
## F-statistic: 6.539 on 2 and 1444 DF, p-value: 0.001489


##

## Call:
## lm(formula = crp ~ hba1c + q_mean, data = china)
##

## Residuals:
```

```
##      Min      1Q  Median      3Q      Max
## -2.9119 -1.1872 -0.8699  0.3433  8.5226
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  0.70968    0.13590   5.222 1.84e-07 ***
## hba1c        0.09596    0.01624   5.911 3.63e-09 ***
## q_mean       0.23115    0.04519   5.115 3.25e-07 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 2.105 on 4961 degrees of freedom
## (10086 observations deleted due to missingness)
## Multiple R-squared:  0.01179,    Adjusted R-squared:  0.01139
## F-statistic: 29.59 on 2 and 4961 DF,  p-value: 1.676e-13
##
## Call:
## lm(formula = crp ~ hba1c + q_mean, data = mex_sub)
##
## Residuals:
##      Min      1Q  Median      3Q      Max
## -2.3312 -1.1832 -0.6210  0.6121  7.9189
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -0.23976    0.75435  -0.318  0.75067
## hba1c        0.31176    0.13164   2.368  0.01805 *
```

```
## q_mean      0.18209    0.06013    3.028    0.00252 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.797 on 1089 degrees of freedom
## (733 observations deleted due to missingness)
## Multiple R-squared:  0.01345,    Adjusted R-squared:  0.01164
## F-statistic: 7.426 on 2 and 1089 DF,  p-value: 0.0006263
```



The descriptive statistics for our sample look as follows:

Table 2

Descriptive statistics.

	China	Mexico
N	7196	1828
Sex		
male	3466 (48.20 %)	39.50 (39.50 %)
female	3723 (51.70 %)	60.40 (60.40 %)
unknown	7 (0.10 %)	1 (0.10 %)
Age	63 (SD = 9.30)	67.70 (SD = 9)
Diabetes		
diagnosed	494 (6.90 %)	374 (20.50 %)
undiagnosed	466 (6.50 %)	182 (10 %)

Discussion

Enter text/code here.

References

- Deshpande, A. D., Harris-Hayes, M., & Schootman, M. (2008). Epidemiology of diabetes and diabetes-related complications. *Physical Therapy*, 88(11), 1254–1264.

Figure captions

Table captions

Table 1. This is where the caption goes

Table 2. Descriptive statistics.