Homework 3 – Regression

1 Question 1

Below, I simulate a rather quite annoying set of regression data.

```
set.seed(0510)
n <- 250
dat.q1 <- tibble(x1 = rnorm(n, mean = 50, sd = 10),
                 x2 = factor(sample(c("A", "B"), n, replace = TRUE)),
                 x3 = factor(sample(c("Control", "Treat"), n, replace = TRUE)),
                 x4 = factor(sample(c("Low", "High"), n, replace = TRUE), levels=c("Low", "High")),
                 x5 = rnorm(n, mean = 0, sd = 1),
                 x6 = factor(sample(c("Group1", "Group2"), n, replace = TRUE)),
                 x7 = rnorm(n, mean = 10, sd = 2),
                 x8 = rnorm(n, mean = 10, sd = 2)) |>
        mutate(y = 5 +                          # Intercept
                   # Quantiative x1
                   2 * x1 +
                   # Categorical x2
                   (x2 == "B") * 10 +
                   # Categorical x Categorical (x3*x4)
                   (x3 == "Treat") * 5 + (x4 == "High") * 2 +
                   (x3 == "Treat" & x4 == "High") * 15 +
                   # Quantitative x Categorical  (x5*x6)
                   3 * x5 + (x6 == "Group2") * 4 +
                   (x5 * (x6 == "Group2")) * 8 +
                   # Quantitative x Quantitative (x7*x8)
                   1.5 * x7 + 0.5 * x8 +
                   0.8 * (x7 * x8) +
                   # Error
                   rnorm(n, mean = 0, sd = 15))
```

1.1 Part a

Fit the regression model.

$$
\begin{aligned}
Y_i = &\beta_0 + \beta_1 x_{1i} + \beta_2 I(x_{2i} = \text{B}) + \beta_3 I(x_{3i} = \text{Treat}) + \beta_4 I(x_{4i} = \text{High}) + \\
&\beta_5 x_{5i} + \beta_6 I(x_{6i} = \text{Group2}) + \beta_7 x_{7i} + \beta_8 x_{8i} + \\
&\beta_9 (I(x_{3i} = \text{Treat}) \times I(x_{4i} = \text{High})) + && \text{(Categorical} \times \text{Categorical)} \\
&\beta_{10} (x_{5i} \times I(x_{6i} = \text{Group2})) + && \text{(Continuous} \times \text{Categorical)} \\
&\beta_{11} (x_{7i} \times x_{8i}) + \epsilon_i && \text{(Continuous} \times \text{Continuous)}
\end{aligned}
$$

Solution: First, we fit the regression model using `lm()`.


```
mod.q1 <- lm(y ~ x1 + x2 + x3*x4 + x5*x6 + x7*x8, data=dat.q1)
```

1.2 Part b

Conduct a residual and outlier analysis.

```
plotResiduals(mod.q1)
```


Figure 1: Residual analysis for the model in Question 1.



In Figure 1, the residuals fall close to the line in the QQ plot and the histogram is approximately Gaussian. The fitted vs. residual plot shows an even band around zero with no pattern. Thus, the residuals appear Gaussian with mean zero and common variance.

```
plotInfluence(mod.q1)
```

Figure 2: Outlier and influence analysis for the model in Question 1.



In Figure 2, a few observations have high leverage and a few have large studentized residuals. However, no observation has both, and all Cook's distances are far below the cutoffs. Thus, no observation has undue influence on the fit, and we do not remove any data.

1.3 Part c

Interpret the intercept, the coefficients on x_1 and x_2.

- **Intercept** ($\hat{\beta}_0 = 8.90$): The expected value of Y when $x_1 = x_5 = x_7 = x_8 = 0$ and the categorical predictors are at their baseline levels ($x_2 =$ A, $x_3 =$ Control, $x_4 =$ Low, $x_6 =$ Group1). This is not meaningful, since x_1, x_7, and x_8 are never near zero in the data; large standard error (26.94).
- x_1 ($\hat{\beta}_1 = 2.14$): Holding all other predictors constant, a one-unit increase in x_1 is associated with a 2.14 increase in the expected value of Y.
- x_2 ($\hat{\beta}_2 = 10.03$): Holding all other predictors constant, the expected value of Y is 10.03 higher when $x_2 =$ B than when $x_2 =$ A.

1.4 Part d

Untangle and interpret the $x_3 * x_4$ interaction using algebra. Use Table 1 and Figure 3 for context.

```
library(emmeans)
emm.q1d <- emmeans(mod.q1, ~x3*x4) |> # Don't average across levels: use |x2*x6
            data.frame() |>
```


```
            rename("Mean"  = "emmean",
                   "Lower" = "lower.CL",
                   "Upper" = "upper.CL")
```

Table 1: Estimated marginal means for the model in Question 1.

x3	x4	Mean	SE	df	Lower	Upper
Control	Low	212.9676	1.7871	238	209.4470	216.4882
Treat	Low	219.4792	1.9123	238	215.7120	223.2464
Control	High	216.4894	1.7916	238	212.9601	220.0188
Treat	High	238.1930	1.7631	238	234.7198	241.6662

```
emm.q1d <- emm.q1d |>
  mutate(Observation = paste("(x3=", x3, ", ", "x4=", x4, ")", sep=""))
ggplot(emm.q1d) +
  geom_point(aes(x=Mean, y=Observation))+
  geom_errorbar(aes(y=Observation,
                    xmin=Lower, xmax=Upper),
                width = 0.25)+
  # facet_wrap(~x2*x6)+
  theme_bw()+
  xlab("Estimated Mean of Y")
```

Figure 3: Estimated marginal means for the model in Question 1.



Holding the other predictors fixed, let C denote the contribution of all other terms. Then

$$\begin{aligned} E(Y \mid \text{Control, Low}) &= C \\ E(Y \mid \text{Treat, Low}) &= C + \beta_3 \\ E(Y \mid \text{Control, High}) &= C + \beta_4 \\ E(Y \mid \text{Treat, High}) &= C + \beta_3 + \beta_4 + \beta_9 \end{aligned}$$

So the effect of Treat (vs. Control) is β_3 when $x_4 = \text{Low}$ and $\beta_3 + \beta_9$ when $x_4 = \text{High}$. That is, β_9 is the difference in the treatment effect between the High and Low groups.

- $\hat{\beta}_3 = 6.51$: When $x_4 = \text{Low}$, Treat is associated with a 6.51 higher expected Y than Control.
- $\hat{\beta}_4 = 3.52$: When $x_3 = \text{Control}$, High is associated with a 3.52 higher expected Y than Low.
- $\hat{\beta}_9 = 15.19$: The treatment effect is 15.19 larger when $x_4 = \text{High}$ $(6.51 + 15.19 = 21.70)$ than when $x_4 = \text{Low}$ (6.51).

These match Table 1: $219.48 - 212.97 = 6.51$, $216.49 - 212.97 = 3.52$, and $238.19 - 216.49 = 21.70$. In Figure 3, the (Treat, High) group sits well above the other three, with a confidence interval that does not overlap theirs, while the other three groups are similar. This reflects the interaction: the treatment effect is much larger when $x_4 = \text{High}$.

1.5 Part e

Untangle and interpret the $x_5 * x_6$ interaction using algebra. Use Table 2 and Figure 4 for context.

```
emm.q1e <- emtrends(mod.q1, var="x5", ~x5|x6) |>
            data.frame() |>
            rename("Slope"  = "x5.trend",
                   "Lower" = "lower.CL",
                   "Upper" = "upper.CL") |>
            dplyr::select(x6, Slope, SE, df, Lower, Upper)
```

Table 2: Estimated marginal trends for the model in Question 1.


x6	Slope	SE	df	Lower	Upper
Group1	2.4835	1.2942	238	-0.0660	5.0330
Group2	9.4309	1.3367	238	6.7977	12.0642

In Figure 4, the slopes are the same and just the intercepts vary depending on x_2, x_3, and x_4.

```
emm.q1e <- emmeans(mod.q1, ~x5*x6,
                   at = list(x5 = seq(-3, 3, length.out=500))) |> # Don't average across levels: use |x2*x3*x4
  data.frame() |>
            rename("Mean"  = "emmean",
                   "Lower" = "lower.CL",
                   "Upper" = "upper.CL")

ggplot(emm.q1e) +
  geom_line(aes(x=x5, y=Mean, color=x6))+
  geom_ribbon(aes(x=x5, ymin=Lower, ymax=Upper, fill=x6), alpha=0.25)+
  # facet_wrap(~x2*x3*x4)+
  theme_bw()
```

Figure 4: Estimated marginal trends for the model in Question 1.



Holding the other predictors fixed, let C denote the contribution of all other terms. Then

$$E(Y \mid x_5, \text{Group1}) = C + \beta_5 x_5$$
$$E(Y \mid x_5, \text{Group2}) = (C + \beta_6) + (\beta_5 + \beta_{10}) x_5$$

So the slope of x_5 is β_5 in Group1 and $\beta_5 + \beta_{10}$ in Group2. That is, β_{10} is the difference in the slope of x_5 between the two groups.

- $\widehat{\beta}_5 = 2.48$: In Group1, a one-unit increase in x_5 is associated with a 2.48 increase in expected Y.
- $\widehat{\beta}_6 = 4.98$: When $x_5 = 0$ (its mean), Group2 has a 4.98 higher expected Y than Group1.
- $\widehat{\beta}_{10} = 6.95$: The slope of x_5 is 6.95 larger in Group2 $(2.48 + 6.95 = 9.43)$ than in Group1 (2.48).

These match Table 2. Note the 95% confidence interval for the Group1 slope, $(-0.07, 5.03)$, includes zero, while the interval for the Group2 slope, $(6.80, 12.06)$, does not. In Figure 4, the Group2 line is much steeper, and the lines cross near $x_5 = -4.98/6.95 \approx -0.72$. So the difference between groups depends on x_5.

1.6 Part f

Untangle and interpret the $x_7 * x_8$ interaction using algebra. Use Table 3 and Table 4 for context.

```
emm.q1f.x7 <- emtrends(mod.q1, var="x7", ~x7|x8,
                   at = list(x8 = mean(dat.q1$x8) + c(-1, 0 ,1)*sd(dat.q1$x8))) |>
            data.frame() |>
            rename("Slope"  = "x7.trend",
                   "Lower" = "lower.CL",
                   "Upper" = "upper.CL") |>
            dplyr::select(x8, Slope, SE, df, Lower, Upper)
```

Table 3: Estimated marginal trends for the model in Question 1.

x8	Slope	SE	df	Lower	Upper
8.1309	7.9230	0.6751	238	6.5932	9.2529
10.0582	9.7129	0.5015	238	8.7250	10.7008
11.9854	11.5027	0.7396	238	10.0458	12.9597

```
emm.q1f.x8 <- emtrends(mod.q1, var="x8", ~x8|x7,
                   at = list(x7 = mean(dat.q1$x7) + c(-1, 0 ,1)*sd(dat.q1$x7))) |>
            data.frame() |>
            rename("Slope"  = "x8.trend",
                   "Lower" = "lower.CL",
                   "Upper" = "upper.CL") |>
            dplyr::select(x7, Slope, SE, df, Lower, Upper)
```

Table 4: Estimated marginal trends for the model in Question 1.

x7	Slope	SE	df	Lower	Upper
8.3824	6.8569	0.6069	238	5.6613	8.0525
10.2208	8.5643	0.4744	238	7.6297	9.4989
12.0592	10.2716	0.7325	238	8.8286	11.7146

A Johnson Neyman analysis using the `interactions` package for `R` (Long 2024), shows that the slope of x_7 is statistically discernible for all observed values of x_8 (Figure 5), and vice versa (Figure 6).

```
library(interactions)
johnson_neyman(mod.q1, pred = x7, modx = x8, control.fdr = T)
```


```
JOHNSON-NEYMAN INTERVAL

When x8 is OUTSIDE the interval [-13.08, 3.35], the slope of x7 is p < .05.

Note: The range of observed values of x8 is [4.69, 15.07]

Interval calculated using false discovery rate adjusted t = 1.97
```

Figure 5: Johnson Neyman interval for the model in Question 1.




```
johnson_neyman(mod.q1, pred = x8, modx = x7, control.fdr = T)
```


```
JOHNSON-NEYMAN INTERVAL

When x7 is OUTSIDE the interval [-9.96, 4.25], the slope of x8 is p < .05.

Note: The range of observed values of x7 is [4.89, 14.60]

Interval calculated using false discovery rate adjusted t = 1.97
```

Figure 6: Johnson Neyman interval for the model in Question 1.



$$\frac{\partial E(Y)}{\partial x_7} = \beta_7 + \beta_{11} x_8$$
$$\frac{\partial E(Y)}{\partial x_8} = \beta_8 + \beta_{11} x_7$$

So the slope of x_7 depends on x_8, and vice versa. The estimated slopes are $0.37 + 0.93x_8$ for x_7 and $-0.93 + 0.93x_7$ for x_8.

- $\hat{\beta}_7 = 0.37$ and $\hat{\beta}_8 = -0.93$ are the slopes when the other variable equals zero. These are not meaningful, since x_7 and x_8 are never near zero in the data.
- $\hat{\beta}_{11} = 0.93$: For each one-unit increase in x_8, the slope of x_7 increases by 0.93 (and vice versa).

Table 3 shows the slope of x_7 increases from 7.92 to 9.71 to 11.50 as x_8 goes from one SD below its mean, to its mean, to one SD above. Similarly, Table 4 shows the slope of x_8 increases from 6.86 to 8.56 to 10.27. All of these confidence intervals exclude zero.

1.7 Part g

Standardize the continuous data and refit the regression model. Redo part f. What do you notice about the coefficient estimates and why does this make the untangling process easier?

Solution: First, we fit the regression model using `lm()`.

```
# Standardize Data
dat.z <- dat.q1 |>
  mutate(across(c(x1, x5, x7, x8), ~ as.vector(scale(.))))
# Refit Model
mod.q1z <- lm(y ~ x1 + x2 + x3*x4 + x5*x6 + x7*x8, data=dat.z)
```

```
emm.q1g.z7 <- emtrends(mod.q1z, var="x7", ~x7|x8,
                       at = list(x8 = c(-1, 0 ,1))) |>
          data.frame() |>
          rename("Slope"  = "x7.trend",
                 "Lower" = "lower.CL",
                 "Upper" = "upper.CL") |>
          dplyr::select(x8, Slope, SE, df, Lower, Upper)
```

Table 5: Estimated marginal trends for the model in Question 1 when quantitative variables are standardized.

x8	Slope	SE	df	Lower	Upper
-1	14.57	1.24	238	12.12	17.01
0	17.86	0.92	238	16.04	19.67
1	21.15	1.36	238	18.47	23.83

```
emm.q1g.z8 <- emtrends(mod.q1z, var="x8", ~x8|x7,
                       at = list(x7 = c(-1, 0 ,1)))|>
              data.frame() |>
              rename("Slope"  = "x8.trend",
                     "Lower" = "lower.CL",
                     "Upper" = "upper.CL") |>
              dplyr::select(x7, Slope, SE, df, Lower, Upper)
```

Table 6: Estimated marginal trends for the model in Question 1 when quantitative variables are standardized.

x7	Slope	SE	df	Lower	Upper
-1	13.22	1.17	238	10.91	15.52
0	16.51	0.91	238	14.70	18.31
1	19.80	1.41	238	17.01	22.58

After standardizing, the slopes are

$$\frac{\partial E(Y)}{\partial z_7} = \beta_7 + \beta_{11} z_8$$
$$\frac{\partial E(Y)}{\partial z_8} = \beta_8 + \beta_{11} z_7$$

where a one-unit change is now a one-SD change. Since $z_8 = 0$ corresponds to the mean of x_8, $\hat{\beta}_7 = 17.86$ is the slope of x_7 (per SD) when x_8 is at its mean, and $\hat{\beta}_8 = 16.51$ is the slope of x_8 (per SD) when x_7 is at its mean. The interaction $\hat{\beta}_{11} = 3.29$ is how much each slope changes for a one-SD increase in the other variable (e.g., 17.86 ± 3.29 gives 14.57 and 21.15 in Table 5).

Compared to part f, the main effects for x_7 and x_8 are now large, precisely estimated, and statistically discernible. This is because the original $\hat{\beta}_7$ and $\hat{\beta}_8$ described slopes at $x_8 = 0$ and $x_7 = 0$, which are far outside the data. Standardizing does not change the model fit; it moves the reference point to the mean, so the coefficients can be interpreted directly without plugging in values of the other variable.

2 Question 2

Cui and Mao (2025) investigate how freezing and cryogenic temperatures affect the strength, flexibility, stiffness, and internal breaking behavior of 3D-printed CF/PEEK composites.

Carbon fiber reinforced PEEK (CF/PEEK) is a high-performance composite used in 3D printing for aerospace, deep-sea, polar research. The researchers aimed to assess whether and how extremely cold environments impact this composite to understand how 3D-printed CF/PEEK performs in freezing conditions. This work would be important to understand its limits and develop preventative methods to protect against structural failures.

The researchers created fifteen 3D printed dog-bone-shaped test specimens and placed them in a chamber cooled with liquid nitrogen to 25°C, −25°C, −75°C, −125°C, and −175°C. To test strength of the specimen, a machine pulled both ends of the specimen until it snapped.

In the table below, the researchers record the average tensile strength for three specimens from each temperature level.

Temperature	Mean Tensile Strength
25°C	81.93 MPa
−25°C	76.63 MPa
−75°C	63.23 MPa
−125°C	32.13 MPa
−175°C	12.83 MPa

```r
dat.q2 <- tibble(temp     = c(25, -25, -75, -125, -175),
                 strength = c(81.93, 76.63, 63.23, 32.13, 12.83))
x <- dat.q2$temp
y <- dat.q2$strength
n <- length(y)
```

2.1 Part a

Compute the estimates we derived using calculus in lecture 16:

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}_1$$
$$\hat{\beta}_1 = \frac{\sum_{i=1}^{n}(x_{1i} - \bar{x}_1)(y_i - \bar{y})}{\sum_{i=1}^{n}(x_{1i} - \bar{x}_1)^2}.$$

Solution:

```r
b1.calc <- sum((x - mean(x)) * (y - mean(y))) / sum((x - mean(x))^2)
b0.calc <- mean(y) - b1.calc * mean(x)
c(b0.calc, b1.calc)
```

```
[1] 80.7550  0.3654
```

Using the formulas, $\hat{\beta}_0 = 80.755$ and $\hat{\beta}_1 = 0.3654$.

2.2 Part b

Compute the estimates we derived using linear algebra in lecture 16:

$$\hat{\beta} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y}.$$

Hint: Run `?matrix()` in `R` for a quick reminder about how to create a matrix.

Solution:

```r
X <- matrix(c(rep(1, n), x), ncol = 2)   # design matrix: column of 1s and x
Y <- matrix(y, ncol = 1)
beta.mat <- solve(t(X) %*% X) %*% t(X) %*% Y
beta.mat
```

```
       [,1]
[1,] 80.7550
[2,]  0.3654
```

Using linear algebra, $\hat{\beta} = (80.755, 0.3654)'$, matching part a.

2.3 Part c

Compute the maximum likelihood estimates using `optim()` in `R` as described in lecture 16:

$$\max_{\beta_0,\beta_1,\sigma} \mathcal{L}(\theta = \{\beta_0, \beta_1, \sigma\}).$$

Solution:

```r
neg.llik <- function(par, x, y) {
  b0    <- par[1]
  b1    <- par[2]
  sigma <- exp(par[3])                    # optimize log(sigma) so sigma > 0
  -sum(dnorm(y, mean = b0 + b1 * x, sd = sigma, log = TRUE))
}

mle <- optim(par = c(mean(y), 0, log(sd(y))),  # starting values
             fn = neg.llik, x = x, y = y)

c(b0 = mle$par[1], b1 = mle$par[2], sigma = exp(mle$par[3]))
```

```
       b0         b1      sigma
80.7533227  0.3653883  6.4900984
```

The MLEs are $\hat{\beta}_0 \approx 80.753$, $\hat{\beta}_1 \approx 0.3654$, and $\hat{\sigma} \approx 6.49$.

2.4 Part d

Fit the model using `lm()` and plot the resulting model. Compare to the results from parts (a) through (c).

Solution:

```r
mod.q2 <- lm(strength ~ temp, data = dat.q2)
summary(mod.q2)
```

```
Call:
lm(formula = strength ~ temp, data = dat.q2)


Residuals:
    1     2     3     4     5
-7.96  5.01  9.88 -2.95 -3.98

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) 80.75500    5.46239  14.784 0.000671 ***
temp         0.36540    0.05299   6.895 0.006250 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 8.379 on 3 degrees of freedom
Multiple R-squared:  0.9406,    Adjusted R-squared:  0.9209
F-statistic: 47.54 on 1 and 3 DF,  p-value: 0.00625
```

```r
ggplot(dat.q2, aes(x = temp, y = strength)) +
  geom_point() +
  geom_smooth(method = "lm", se = FALSE) +
  theme_bw() +
  xlab("Temperature (°C)") +
  ylab("Mean Tensile Strength (MPa)")
```

Figure 7: Mean tensile strength versus temperature with the fitted regression line.



All four approaches give really similar estimates: $\hat{\beta}_0 = 80.755$ and $\hat{\beta}_1 = 0.3654$. That is, at 0°C the expected mean tensile strength is 80.76 MPa, and for each 1°C increase in temperature, mean tensile strength is expected to increase by 0.37 MPa. The only difference is the estimate of σ: the MLE divides by n ($\hat{\sigma} = 6.49$), while `lm()` divides by $n - p = 3$ to give the unbiased estimate ($s = 8.38$).

3 Question 3

Bracht et al. (2016) evaluate MFAP4 as a biomarker for severity of fibrosis in hepatitic C patients.

```r
dat.mfap4 <- read_csv("Bracht2016.csv")
```

3.1 Part a

Fit the model predicting `MFAP4` using `Fibrosis.Stage`. Write out the resulting estimate model and interpret the coefficients.

```r
mod.q3a <- lm(MFAP4 ~ Fibrosis.Stage, data = dat.mfap4)
summary(mod.q3a)
```

```
Call:
lm(formula = MFAP4 ~ Fibrosis.Stage, data = dat.mfap4)

Residuals:
    Min      1Q  Median      3Q     Max
-20.318  -5.518  -2.651   2.070  81.816

Coefficients:
                     Estimate Std. Error t value Pr(>|t|)
(Intercept)             9.230      1.102   8.378 4.73e-16 ***
Fibrosis.StageStage 1   1.301      1.372   0.948   0.3433
Fibrosis.StageStage 2   3.154      1.444   2.184   0.0294 *
Fibrosis.StageStage 3  11.742      1.724   6.812 2.59e-11 ***
Fibrosis.StageStage 4  15.388      1.724   8.928  < 2e-16 ***
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 10.85 on 537 degrees of freedom
Multiple R-squared:  0.1965,    Adjusted R-squared:  0.1905
F-statistic: 32.84 on 4 and 537 DF,  p-value: < 2.2e-16
```

The estimated model is

$$\widehat{\text{MFAP4}} = 9.23 + 1.30\, I(\text{Stage } 1) + 3.15\, I(\text{Stage } 2) + 11.74\, I(\text{Stage } 3) + 15.39\, I(\text{Stage } 4).$$

- $\hat{\beta}_0 = 9.23$: The expected MFAP4 level for patients with Stage 0 fibrosis.
- $\hat{\beta}_1 = 1.30$: Patients with Stage 1 fibrosis have an expected MFAP4 level 1.30 higher than Stage 0.
- $\hat{\beta}_2 = 3.15$: Stage 2 is 3.15 higher than Stage 0.
- $\hat{\beta}_3 = 11.74$: Stage 3 is 11.74 higher than Stage 0.
- $\hat{\beta}_4 = 15.39$: Stage 4 is 15.39 higher than Stage 0.


So expected MFAP4 increases with fibrosis stage, with a large jump from Stage 2 to Stage 3.

3.2 Part b

Conduct a residual analysis on the model in part a to demonstrate how to identify when you need a log transformation.

```r
plotResiduals(mod.q3a)
```

Figure 8: Residual analysis for the model in part a.



In Figure 8, the QQ plot curves upward in the right tail, and the histogram has a long right tail compared to the Gaussian curve. The fitted vs. residual plot also shows the spread of the residuals increasing with the fitted values.A log transformation of the response is needed.

3.3 Part c

Fit the model predicting `log(MFAP4)` using `Fibrosis.Stage`. Use the natural log, and not the $\log_2(\cdot)$ recorded in the data file. Write out the resulting estimate model and interpret the coefficients.

```r
mod.q3c <- lm(log(MFAP4) ~ Fibrosis.Stage, data = dat.mfap4)
summary(mod.q3c)
```

```
Call:
lm(formula = log(MFAP4) ~ Fibrosis.Stage, data = dat.mfap4)

Residuals:
     Min       1Q   Median       3Q      Max
-1.55668 -0.41861 -0.04282  0.35845  2.21571

Coefficients:
                      Estimate Std. Error t value Pr(>|t|)
(Intercept)            2.02572    0.06178  32.789  < 2e-16 ***
Fibrosis.StageStage 1  0.13336    0.07694   1.733 0.083637 .
Fibrosis.StageStage 2  0.30399    0.08099   3.753 0.000193 ***
Fibrosis.StageStage 3  0.77408    0.09666   8.009 7.24e-15 ***
Fibrosis.StageStage 4  0.98957    0.09666  10.238  < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.6085 on 537 degrees of freedom
Multiple R-squared:  0.2312,    Adjusted R-squared:  0.2255
F-statistic: 40.38 on 4 and 537 DF,  p-value: < 2.2e-16
```

```r
exp(coef(mod.q3c))
```

```
          (Intercept) Fibrosis.StageStage 1 Fibrosis.StageStage 2
            7.581571             1.142659             1.355252
Fibrosis.StageStage 3 Fibrosis.StageStage 4
            2.168599             2.690090
```

The estimated model is

$$\begin{aligned}\widehat{\log(\text{MFAP4})} = 2.03 &+ 0.13\, I(\text{Stage } 1) + 0.30\, I(\text{Stage } 2) \\ &+ 0.77\, I(\text{Stage } 3) + 0.99\, I(\text{Stage } 4).\end{aligned}$$


- $e^{\widehat{\beta}_0} = e^{2.03} = 7.58$: The median MFAP4 level for Stage 0 patients is 7.58.
- $e^{\widehat{\beta}_1} = 1.14$: The median MFAP4 for Stage 1 is 1.14 times that of Stage 0 (14% higher).
- $e^{\widehat{\beta}_2} = 1.36$: Stage 2 is 1.36 times Stage 0 (36% higher).
- $e^{\widehat{\beta}_3} = 2.17$: Stage 3 is 2.17 times Stage 0 (117% higher).
- $e^{\widehat{\beta}_4} = 2.69$: Stage 4 is 2.69 times Stage 0 (169% higher).

3.4 Part d

Conduct a residual analysis on the model in part c to demonstrate that the conditions for using regression are met.

```r
plotResiduals(mod.q3c)
```

Figure 9: Residual analysis for the model in part c.



In Figure 9, after the log transformation the residuals fall close to the line in the QQ plot, with only slight deviation in the right tail, and the histogram closely matches the Gaussian curve. The fitted vs. residual plot shows bands of similar width centered at zero across all stages. Thus, the conditions for regression appear to be met.

3.5 Part e

Fit the model predicting `log(MFAP4)` using `Fibrosis.Stage`, `Age`, and `Gender`. Use the natural log, and not the $\log_2(\cdot)$ recorded in the data file. Write out the resulting estimate model and interpret the coefficients.

```
mod.q3e <- lm(log(MFAP4) ~ Fibrosis.Stage + Age + Gender, data = dat.mfap4)
summary(mod.q3e)
```

```
Call:
lm(formula = log(MFAP4) ~ Fibrosis.Stage + Age + Gender, data = dat.mfap4)

Residuals:
     Min       1Q   Median       3Q      Max
-1.55799 -0.40010 -0.03682  0.35968  2.28903

Coefficients:
                      Estimate Std. Error t value Pr(>|t|)
(Intercept)           1.518667   0.115808  13.114  < 2e-16 ***
Fibrosis.StageStage 1 0.126858   0.073594   1.724  0.08533 .
Fibrosis.StageStage 2 0.212539   0.079095   2.687  0.00743 **
Fibrosis.StageStage 3 0.598866   0.096575   6.201 1.12e-09 ***
Fibrosis.StageStage 4 0.828547   0.097208   8.523  < 2e-16 ***
Age                   0.013110   0.002123   6.175 1.31e-09 ***
Gendermale           -0.143405   0.050807  -2.823  0.00494 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.5812 on 535 degrees of freedom
Multiple R-squared:  0.3012,    Adjusted R-squared:  0.2934
F-statistic: 38.44 on 6 and 535 DF,  p-value: < 2.2e-16
```

```
exp(coef(mod.q3e))
```

```
         (Intercept) Fibrosis.StageStage 1 Fibrosis.StageStage 2
           4.5661359             1.1352559             1.2368149
Fibrosis.StageStage 3 Fibrosis.StageStage 4                   Age
           1.8200537             2.2899898             1.0131960
          Gendermale
           0.8664034
```

The estimated model is

$$\log(\widehat{\text{MFAP4}}) = 1.52 + 0.13\, I(\text{Stage 1}) + 0.21\, I(\text{Stage 2}) + 0.60\, I(\text{Stage 3}) + 0.83\, I(\text{Stage 4}) + 0.013\,\text{Age} - 0.14\, I(\text{Male}).$$

Exponentiating the coefficients, holding the other predictors constant:

- $e^{\widehat{\beta}_0} = 4.57$: The median MFAP4 for a female Stage 0 patient of age 0. This is not meaningful, since no patients are near age 0.
- $e^{\widehat{\beta}_1} = 1.14$, $e^{\widehat{\beta}_2} = 1.24$, $e^{\widehat{\beta}_3} = 1.82$, $e^{\widehat{\beta}_4} = 2.29$: Compared to Stage 0 patients of the same age and gender, the median MFAP4 is 14%, 24%, 82%, and 129% higher for Stages 1, 2, 3, and 4, respectively.
- $e^{\widehat{\beta}_5} = 1.013$: Each additional year of age is associated with a 1.3% higher median MFAP4.
- $e^{\widehat{\beta}_6} = 0.87$: Males have a median MFAP4 about 13% lower than females.

Compared to part c, the stage effects are smaller after adjusting for age (e.g., Stage 4 drops from 2.69 to 2.29 times Stage 0). This suggests part of the stage effect in part c was due to patients with more advanced fibrosis tending to be older.

4 Question 4

```r
# Simulate R regressions, storing the coefficient estimates
sim.coefs <- function(error.fun, R = 1000, n = 30) {
  results <- tibble(b0 = rep(NA, R), b1 = rep(NA, R))
  for (i in 1:R) {
    x <- runif(n, min = 0, max = 10)
    e <- error.fun(n)
    y <- 2 + 1.5 * x + e
    mod <- lm(y ~ x)
    results$b0[i] <- coef(mod)[1]
    results$b1[i] <- coef(mod)[2]
  }
  results
}

# Plot the sampling distributions with the true values marked
plot.coefs <- function(results) {
  p0 <- ggplot(results, aes(x = b0)) +
    geom_histogram(aes(y = after_stat(density)), bins = 30,
                   fill = "lightgrey", color = "black") +
    geom_vline(xintercept = 2, color = "red", linewidth = 1) +
    theme_bw() + xlab(bquote(hat(beta)[0]))
  p1 <- ggplot(results, aes(x = b1)) +
    geom_histogram(aes(y = after_stat(density)), bins = 30,
                   fill = "lightgrey", color = "black") +
    geom_vline(xintercept = 1.5, color = "red", linewidth = 1) +
    theme_bw() + xlab(bquote(hat(beta)[1]))
  p0 | p1
}
```

4.1 Part a

Simulate data to match regression assumptions in a loop that runs $R = 1000$ times.


$$\begin{aligned} x_i &\sim \text{Uniform}(a = 0, b = 10) \\ \epsilon_i &\sim \text{Gaussian}(\mu = 0, \sigma = 1) \\ y_i &= 2 + 1.5x_i + \epsilon_i \\ n &= 30 \end{aligned}$$

In each iteration, fit a regression line to the data and store the coefficient estimates. Plot the resulting estimates. What do you notice?

```r
set.seed(354)
res.a <- sim.coefs(function(n) rnorm(n, mean = 0, sd = 1))
plot.coefs(res.a)
summary(res.a)
```

```
       b0              b1       
 Min.   :0.8418   Min.   :1.297  
 1st Qu.:1.7508   1st Qu.:1.457  
 Median :1.9986   Median :1.502  
 Mean   :1.9992   Mean   :1.500  
 3rd Qu.:2.2443   3rd Qu.:1.544  
 Max.   :3.1626   Max.   :1.735  
```

Figure 10: Sampling distributions of the coefficient estimates with Gaussian errors. Red lines mark the true values.



Both sampling distributions are bell-shaped and centered at the true values: the mean of $\hat{\beta}_0$ is 1.999 (true value 2) and the mean of $\hat{\beta}_1$ is 1.500 (true value 1.5). When the regression conditions are met, the OLS estimators are unbiased and approximately Gaussian.

4.2 Part b

Simulate data to match regression assumptions in a loop that runs $R = 1000$ times.

$$
\begin{aligned}
x_i &\sim \text{Uniform}(a = 0, b = 10) \\
\epsilon_i &\sim \text{Shifted Exponential}(\lambda = 1, \mu = 0) \\
y_i &= 2 + 1.5x_i + \epsilon_i \\
n &= 30
\end{aligned}
$$

In each iteration, fit a regression line to the data and store the coefficient estimates. Plot the resulting estimates. What do you notice?


Hint: Use the `tolerance::r2exp()` function with `rate=1` and `shift=-1`.

```
set.seed(354)
res.b <- sim.coefs(function(n) tolerance::r2exp(n, rate = 1, shift = -1))
plot.coefs(res.b)
summary(res.b)
```

```
       b0              b1       
 Min.   :0.9936   Min.   :1.296  
 1st Qu.:1.7279   1st Qu.:1.457  
 Median :1.9847   Median :1.501  
 Mean   :2.0081   Mean   :1.500  
 3rd Qu.:2.2414   3rd Qu.:1.543  
```

```
 Max.   :3.3185   Max.   :1.679
```

Figure 11: Sampling distributions of the coefficient estimates with shifted exponential errors. Red lines mark the true values.



Even though the errors are skewed, both sampling distributions are still centered at the true values. The distribution of $\hat{\beta}_1$ is approximately Gaussian, while $\hat{\beta}_0$ shows slight right skew. So OLS remains unbiased without Gaussian errors, and since the estimators are weighted averages of the data, the Central Limit Theorem makes their sampling distributions close to Gaussian even with $n = 30$.

4.3 Part c

Simulate data to match regression assumptions in a loop that runs $R = 1000$ times.


$$
\begin{aligned}
x_i &\sim \text{Uniform}(a = 0, b = 10) \\
\epsilon_i &\sim \text{Cauchy}(m = 0, s = 1) \\
y_i &= 2 + 1.5x_i + \epsilon_i \\
n &= 30
\end{aligned}
$$

In each iteration, fit a regression line to the data and store the coefficient estimates. Plot the resulting estimates. What do you notice?

```r
set.seed(354)
res.c <- sim.coefs(function(n) rcauchy(n, location = 0, scale = 1))
plot.coefs(res.c)
summary(res.c)
```

```
       b0                    b1
 Min.   :-1237.4315   Min.   :-69.527
 1st Qu.:    0.5564   1st Qu.:  1.177
 Median :    2.1740   Median :  1.490
 Mean   :    0.4713   Mean   :  1.624
 3rd Qu.:    3.9374   3rd Qu.:  1.761
 Max.   :  247.2638   Max.   :150.129
```

Figure 12: Sampling distributions of the coefficient estimates with Cauchy errors. Red lines mark the true values.



With Cauchy errors, the sampling distributions have extremely heavy tails: $\hat{\beta}_0$ ranges from -1237 to 247 and $\hat{\beta}_1$ from -69.5 to 150.1. While the medians are near the true values (2.17 and 1.49), the means are not (0.47 and 1.62), and a few extreme estimates dominate the histograms. Since the Cauchy distribution has no mean or variance, the Central Limit Theorem does not apply, and OLS estimates can be wildly wrong. Regression requires errors with finite variance.

References

Bracht, Thilo, Christian Mölleken, Maike Ahrens, et al. 2016. "Evaluation of the Biomarker Candidate MFAP4 for Non-Invasive Assessment of Hepatic Fibrosis in Hepatitis C Patients." *Journal of Translational Medicine* 14 (1): 201.

Cui, Mengyang, and Jian Mao. 2025. "Effect of Low Temperature on Mechanical Properties and Fracture Behavior of 3D Printed Short Carbon Fiber Reinforced Polyether Ether Ketone Composites (CF/PEEK)." *Scientific Reports* 15 (1): 36244.

Long, Jacob A. 2024. *Interactions: Comprehensive, User-Friendly Toolkit for Probing Interactions*. https://doi.org/10.32614/CRAN.package.interactions.