



NTNU

Norwegian University of Science and Technology

The Convex Bundle Method on Hadamard Manifolds

Hajg Jasa

joint work with

Ronny Bergmann and Roland Herzog

Manifolds and Geometric Integration Colloquia, Ilsetra (Øyer)

March 1st, 2023

Notation

- ▶ Smooth Riemannian manifold \mathcal{M}
- ▶ Tangent space $\mathcal{T}_p\mathcal{M}$ at the point $p \in \mathcal{M}$
- ▶ Inner product $\langle \cdot, \cdot \rangle_p: \mathcal{T}_p\mathcal{M} \times \mathcal{T}_p\mathcal{M} \rightarrow \mathbb{R}$
- ▶ Exponential map $\exp_p X_p = \gamma_{pq}(1) = q$
- ▶ Logarithmic map $\log_p q = \exp_p^{-1} q = X_p$
- ▶ Parallel transport $P_{q \leftarrow p}: \mathcal{T}_p\mathcal{M} \rightarrow \mathcal{T}_q\mathcal{M}$

The Problem

Consider the following minimization problem

$$\arg \min_{p \in \mathcal{M}} f(p)$$

where

The Problem

Consider the following minimization problem

$$\arg \min_{p \in \mathcal{M}} f(p)$$

where

- ▶ \mathcal{M} is a Hadamard manifold:

The Problem

Consider the following minimization problem

$$\arg \min_{p \in \mathcal{M}} f(p)$$

where

- ▶ \mathcal{M} is a Hadamard manifold:
 - ▶ complete
 - ▶ simply connected
 - ▶ non-positive sectional curvature everywhere

The Problem

Consider the following minimization problem

$$\arg \min_{p \in \mathcal{M}} f(p)$$

where

- ▶ \mathcal{M} is a Hadamard manifold:
 - ▶ complete
 - ▶ simply connected
 - ▶ non-positive sectional curvature everywhere
- ▶ $f: \mathcal{M} \rightarrow \mathbb{R}$ is

The Problem

Consider the following minimization problem

$$\arg \min_{p \in \mathcal{M}} f(p)$$

where

- ▶ \mathcal{M} is a Hadamard manifold:
 - ▶ complete
 - ▶ simply connected
 - ▶ non-positive sectional curvature everywhere
- ▶ $f: \mathcal{M} \rightarrow \mathbb{R}$ is
 - ▶ geodesically convex: $f \circ \gamma: [0, 1] \rightarrow \mathbb{R}$ is convex in the usual sense
 - ▶ lower semi-continuous: $\liminf_{q \rightarrow p} f(q) \geq f(p)$

The Problem

Consider the following minimization problem

$$\arg \min_{p \in \mathcal{M}} f(p)$$

where

- ▶ \mathcal{M} is a Hadamard manifold:
 - ▶ complete
 - ▶ simply connected
 - ▶ non-positive sectional curvature everywhere
- ▶ $f: \mathcal{M} \rightarrow \mathbb{R}$ is
 - ▶ geodesically convex: $f \circ \gamma: [0, 1] \rightarrow \mathbb{R}$ is convex in the usual sense
 - ▶ lower semi-continuous: $\liminf_{q \rightarrow p} f(q) \geq f(p)$

Goal. Solve this non-smooth optimization problem with the bundle method

The Subdifferentials

For a convex function, the subdifferential is defined as

$$\partial f(x) = \left\{ s \in \mathbb{R}^n \mid f(y) \geq f(x) + (s)^T (y - x) \text{ for all } y \in \mathbb{R}^n \right\}$$

and it is a non-empty, closed and convex subset.

Let $\varepsilon > 0$. The ε -subdifferential is

$$\partial_\varepsilon f(x) = \left\{ s \in \mathbb{R}^n \mid f(y) \geq f(x) + (s)^T (y - x) - \varepsilon \text{ for all } y \in \mathbb{R}^n \right\}$$

and

$$\partial f(x) \subseteq \partial_\varepsilon f(x)$$

The Subdifferentials

For a geodesically convex function, the subdifferential is defined as

$$\partial f(p) = \{X_p \in \mathcal{T}_p \mathcal{M} \mid f(q) \geq f(p) + \langle X_p, \log_p q \rangle \text{ for all } q \in \mathcal{U} \subseteq \mathcal{M}\}$$

and it is a non-empty, closed and convex subset.

Let $\varepsilon > 0$. The ε -subdifferential is

$$\partial_\varepsilon f(p) = \{X_p \in \mathcal{T}_p \mathcal{M} \mid f(q) \geq f(p) + \langle X_p, \log_p q \rangle - \varepsilon \text{ for all } q \in \mathcal{U} \subseteq \mathcal{M}\}$$

and

$$\partial f(p) \subseteq \partial_\varepsilon f(p)$$

The Bundle Method

Bundle methods are about descent as well as stability.

The Bundle Method

Bundle methods are about descent as well as stability.

Core characteristics of a general bundle algorithm (Bonnans et al. 2006):

The Bundle Method

Bundle methods are about descent as well as stability.

Core characteristics of a general bundle algorithm (Bonnans et al. 2006):

- ▶ keeps memory of the last “best” points

The Bundle Method

Bundle methods are about descent as well as stability.

Core characteristics of a general bundle algorithm (Bonnans et al. 2006):

- ▶ keeps memory of the last “best” points
- ▶ solves a stabilization subproblem

The Bundle Method

Bundle methods are about descent as well as stability.

Core characteristics of a general bundle algorithm (Bonnans et al. 2006):

- ▶ keeps memory of the last “best” points
- ▶ solves a stabilization subproblem
 - ▶ we employ a dual approach with the ε –subdifferential as in Geiger and Kanzow 2002

The Bundle Method

Bundle methods are about descent as well as stability.

Core characteristics of a general bundle algorithm (Bonnans et al. 2006):

- ▶ keeps memory of the last “best” points
- ▶ solves a stabilization subproblem
 - ▶ we employ a dual approach with the ε –subdifferential as in Geiger and Kanzow 2002
- ▶ generates sequences of *candidate* points and *stability centers*

Approximating the ε -subdifferential

Let $\varepsilon > 0$ and $k \in \mathbb{N}$. Given $x_1, \dots, x_k \in \mathbb{R}^n$, and $s^j \in \partial f(x_j)$, then

Approximating the ε -subdifferential

Let $\varepsilon > 0$ and $k \in \mathbb{N}$. Given $x_1, \dots, x_k \in \mathbb{R}^n$, and $s^j \in \partial f(x_j)$, then
 $s^j \in \partial_\varepsilon f(x_k)$

Approximating the ε -subdifferential

Let $\varepsilon > 0$ and $k \in \mathbb{N}$. Given $x_1, \dots, x_k \in \mathbb{R}^n$, and $s^j \in \partial f(x_j)$, then

$$s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq e_j^k = f(x_k) - f(x_j) - (s^j)^T (x_k - x_j) \quad \text{for all } j \in \{1, \dots, k\}$$

Approximating the ε -subdifferential

Let $\varepsilon > 0$ and $k \in \mathbb{N}$. Given $x_1, \dots, x_k \in \mathbb{R}^n$, and $s^j \in \partial f(x_j)$, then

$$s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq e_j^k = f(x_k) - f(x_j) - (s^j)^T (x_k - x_j) \quad \text{for all } j \in \{1, \dots, k\}$$

- **Main challenge on manifolds:** given $p_1, \dots, p_k \in \mathcal{M}$, and $X_{p_j} \in \partial f(p_j)$, then

Approximating the ε -subdifferential

Let $\varepsilon > 0$ and $k \in \mathbb{N}$. Given $x_1, \dots, x_k \in \mathbb{R}^n$, and $s^j \in \partial f(x_j)$, then

$$s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq e_j^k = f(x_k) - f(x_j) - (s^j)^T (x_k - x_j) \quad \text{for all } j \in \{1, \dots, k\}$$

- **Main challenge on manifolds:** given $p_1, \dots, p_k \in \mathcal{M}$, and $X_{p_j} \in \partial f(p_j)$, then

$$P_{p_k \leftarrow p_j} X_{p_j} \in \partial_\varepsilon f(p_k)?$$

Approximating the ε -subdifferential

Let $\varepsilon > 0$ and $k \in \mathbb{N}$. Given $x_1, \dots, x_k \in \mathbb{R}^n$, and $s^j \in \partial f(x_j)$, then

$$s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq e_j^k = f(x_k) - f(x_j) - (s^j)^T (x_k - x_j) \quad \text{for all } j \in \{1, \dots, k\}$$

- **Main challenge on manifolds:** given $p_1, \dots, p_k \in \mathcal{M}$, and $X_{p_j} \in \partial f(p_j)$, then

$$P_{p_k \leftarrow p_j} X_{p_j} \in \partial_\varepsilon f(p_k)?$$

- For all $\sigma > 0$ and all $p_j, p_{j+1} \in \mathcal{M}$, there exists $\delta > 0$ such that

$$\|P_{p_{j+1} \leftarrow p_j} X_{p_j} - P_{p_{j+1} \leftarrow p_{j+2}} P_{p_{j+2} \leftarrow p_j} X_{p_j}\| < \sigma \|X_{p_j}\|$$

for all $p_{j+2} \in B_\delta(p_j) \cup B_\delta(p_{j+1})$ and all $X_{p_j} \in \mathcal{T}_{p_j} \mathcal{M}$ (Azagra and Ferrera 2005, p. 169).

Approximating the ε -subdifferential

Let $\varepsilon > 0$ and $k \in \mathbb{N}$. Given $x_1, \dots, x_k \in \mathbb{R}^n$, and $s^j \in \partial f(x_j)$, then

$$s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq e_j^k = f(x_k) - f(x_j) - (s^j)^T (x_k - x_j) \quad \text{for all } j \in \{1, \dots, k\}$$

- **Main challenge on manifolds:** given $p_1, \dots, p_k \in \mathcal{M}$, and $X_{p_j} \in \partial f(p_j)$, then

$$P_{p_k \leftarrow p_j} X_{p_j} \in \partial_\varepsilon f(p_k)?$$

- For all $\sigma > 0$ and all $p_j, p_{j+1} \in \mathcal{M}$, there exists $\delta > 0$ such that

$$\|P_{p_{j+1} \leftarrow p_j} X_{p_j} - P_{p_{j+1} \leftarrow p_{j+2}} P_{p_{j+2} \leftarrow p_j} X_{p_j}\| < \sigma \|X_{p_j}\|$$

for all $p_{j+2} \in B_\delta(p_j) \cup B_\delta(p_{j+1})$ and all $X_{p_j} \in \mathcal{T}_{p_j} \mathcal{M}$ (Azagra and Ferrera 2005, p. 169).

- Enlarge the ε -subdifferential by adding $2\sigma \|X_{p_j}\|$:

$$c_j^k = f(p_k) - f(p_j) - \langle X_{p_j}, \log_{p_j} p_k \rangle + 2\sigma \|X_{p_j}\|$$

The Euclidean Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients.

The Euclidean Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Euclidean case:

$$s^j \in \partial f(x_j) \Rightarrow \sum_{j=1}^k \lambda_j s^j \in \partial_\varepsilon f(x_k)$$

The Euclidean Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Euclidean case:

$$s^j \in \partial f(x_j) \Rightarrow \sum_{j=1}^k \lambda_j s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j e_j^k$$

The Euclidean Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Euclidean case:

$$s^j \in \partial f(x_j) \Rightarrow \sum_{j=1}^k \lambda_j s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j e_j^k$$

Then the search direction is

$$d^k = - \sum_{j=1}^k \lambda_j s^j$$

The Euclidean Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Euclidean case:

$$s^j \in \partial f(x_j) \Rightarrow \sum_{j=1}^k \lambda_j s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j e_j^k$$

Then the search direction is

$$d^k = - \sum_{j=1}^k \lambda_j s^j$$

where the coefficients λ_j are solution to

The Euclidean Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Euclidean case:

$$s^j \in \partial f(x_j) \Rightarrow \sum_{j=1}^k \lambda_j s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j e_j^k$$

Then the search direction is

$$d^k = - \sum_{j=1}^k \lambda_j s^j$$

where the coefficients λ_j are solution to

$$\begin{aligned} \arg \min_{\lambda \in \mathbb{R}^{|J_k|}} \quad & \frac{1}{2} \left\| \sum_{j \in J_k} \lambda_j s^j \right\|^2 + \sum_{j \in J_k} \lambda_j e_j^k \\ \text{s.t.} \quad & \sum_{j \in J_k} \lambda_j = 1, \quad \lambda_j \geq 0 \quad \text{for all } j \in J_k \end{aligned}$$

The Hadamard Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Hadamard case:

$$s^j \in \partial f(x_j) \Rightarrow \sum_{j=1}^k \lambda_j s^j \in \partial_\varepsilon f(x_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j e_j^k$$

Then the search direction is

$$d^k = - \sum_{j=1}^k \lambda_j s^j$$

where the coefficients λ_j are solution to

$$\begin{aligned} \arg \min_{\lambda \in \mathbb{R}^{|J_k|}} \quad & \frac{1}{2} \left\| \sum_{j \in J_k} \lambda_j s^j \right\|^2 + \sum_{j \in J_k} \lambda_j e_j^k \\ \text{s.t.} \quad & \sum_{j \in J_k} \lambda_j = 1, \quad \lambda_j \geq 0 \quad \text{for all } j \in J_k \end{aligned}$$

The Hadamard Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Hadamard case:

$$X_{q_j} \in \partial f(q_j) \Rightarrow \sum_{j=1}^k \lambda_j P_{p_k \leftarrow q_j} X_{q_j} \in \partial_\varepsilon f(p_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j c_j^k$$

Then the search direction is

$$d^k = - \sum_{j=1}^k \lambda_j s^j$$

where the coefficients λ_j are solution to

$$\begin{aligned} \arg \min_{\lambda \in \mathbb{R}^{|J_k|}} \quad & \frac{1}{2} \left\| \sum_{j \in J_k} \lambda_j s^j \right\|^2 + \sum_{j \in J_k} \lambda_j e_j^k \\ \text{s.t.} \quad & \sum_{j \in J_k} \lambda_j = 1, \quad \lambda_j \geq 0 \quad \text{for all } j \in J_k \end{aligned}$$

The Hadamard Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Hadamard case:

$$X_{q_j} \in \partial f(q_j) \Rightarrow \sum_{j=1}^k \lambda_j P_{p_k \leftarrow q_j} X_{q_j} \in \partial_\varepsilon f(p_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j c_j^k$$

Then the search direction is

$$d^k = - \sum_{j=1}^k \lambda_j P_{p_k \leftarrow q_j} X_{q_j}$$

where the coefficients λ_j are solution to

$$\begin{aligned} \arg \min_{\lambda \in \mathbb{R}^{|J_k|}} \quad & \frac{1}{2} \left\| \sum_{j \in J_k} \lambda_j s_j^k \right\|^2 + \sum_{j \in J_k} \lambda_j e_j^k \\ \text{s.t.} \quad & \sum_{j \in J_k} \lambda_j = 1, \quad \lambda_j \geq 0 \quad \text{for all } j \in J_k \end{aligned}$$

The Hadamard Subproblem

Let $k \in \mathbb{N}$ and $j \in \{1, \dots, k\} = J_k$ and let λ_j be convex coefficients. In the Hadamard case:

$$X_{q_j} \in \partial f(q_j) \Rightarrow \sum_{j=1}^k \lambda_j P_{p_k \leftarrow q_j} X_{q_j} \in \partial_\varepsilon f(p_k) \quad \text{if} \quad \varepsilon \geq \sum_{j=1}^k \lambda_j c_j^k$$

Then the search direction is

$$d^k = - \sum_{j=1}^k \lambda_j P_{p_k \leftarrow q_j} X_{q_j}$$

where the coefficients λ_j are solution to

$$\begin{aligned} \arg \min_{\lambda \in \mathbb{R}^{|J_k|}} \quad & \frac{1}{2} \left\| \sum_{j \in J_k} \lambda_j P_{p_k \leftarrow q_j} X_{q_j} \right\|^2 + \sum_{j \in J_k} \lambda_j c_j^k \\ \text{s.t.} \quad & \sum_{j \in J_k} \lambda_j = 1, \quad \lambda_j \geq 0 \quad \text{for all } j \in J_k \end{aligned}$$

The Euclidean Bundle Method

Data: $x_1 = y_1 \in \mathbb{R}^n$, $g^0 = s^1 \in \partial f(x_1)$, $m \in (0, 1)$, $\varepsilon^0 = e_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k s^j, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k e_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $y_{k+1} = x_k + d^k$ and compute $s^{k+1} \in \partial f(y_{k+1})$
- 3 If $f(y_{k+1}) \leq f(x_k) + m\xi^k$, then set $x_{k+1} = y_{k+1}$, else set $x_{k+1} = x_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$e_j^{k+1} = f(x_{k+1}) - f(y_j) - (s^j)^T (x_{k+1} - y_j) \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{x_k\}_{k \in \mathbb{N}}$, $\{y_k\}_{k \in \mathbb{N}}$

The Convex Hadamard Bundle Method

Data: $p_1 = q_1 \in \mathcal{M}$, $g^0 = x_1 \in \partial f(p_1)$, $m \in (0, 1)$, $\varepsilon^0 = c_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k s^j, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k e_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $y_{k+1} = x_k + d^k$ and compute $s^{k+1} \in \partial f(y_{k+1})$
- 3 If $f(y_{k+1}) \leq f(x_k) + m\xi^k$, then set $x_{k+1} = y_{k+1}$, else set $x_{k+1} = x_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$e_j^{k+1} = f(x_{k+1}) - f(y_j) - (s^j)^T (x_{k+1} - y_j) \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{x_k\}_{k \in \mathbb{N}}$, $\{y_k\}_{k \in \mathbb{N}}$

The Convex Hadamard Bundle Method

Data: $p_1 = q_1 \in \mathcal{M}$, $g^0 = x_1 \in \partial f(p_1)$, $m \in (0, 1)$, $\varepsilon^0 = c_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k P_{p_k \leftarrow q_j} x_{q_j}, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k c_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $y_{k+1} = x_k + d^k$ and compute $s^{k+1} \in \partial f(y_{k+1})$
- 3 If $f(y_{k+1}) \leq f(x_k) + m\xi^k$, then set $x_{k+1} = y_{k+1}$, else set $x_{k+1} = x_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$e_j^{k+1} = f(x_{k+1}) - f(y_j) - (s^j)^T (x_{k+1} - y_j) \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{x_k\}_{k \in \mathbb{N}}$, $\{y_k\}_{k \in \mathbb{N}}$

The Convex Hadamard Bundle Method

Data: $p_1 = q_1 \in \mathcal{M}$, $g^0 = X_1 \in \partial f(p_1)$, $m \in (0, 1)$, $\varepsilon^0 = c_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k P_{p_k \leftarrow q_j} X_{q_j}, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k c_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $q_{k+1} = \exp_{p_k}(d^k)$ and compute $X_{q_{k+1}} \in \partial f(q_{k+1})$
- 3 If $f(y_{k+1}) \leq f(x_k) + m\xi^k$, then set $x_{k+1} = y_{k+1}$, else set $x_{k+1} = x_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$e_j^{k+1} = f(x_{k+1}) - f(y_j) - (s^j)^T (x_{k+1} - y_j) \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{x_k\}_{k \in \mathbb{N}}$, $\{y_k\}_{k \in \mathbb{N}}$

The Convex Hadamard Bundle Method

Data: $p_1 = q_1 \in \mathcal{M}$, $g^0 = X_1 \in \partial f(p_1)$, $m \in (0, 1)$, $\varepsilon^0 = c_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k P_{p_k \leftarrow q_j} X_{q_j}, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k c_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $q_{k+1} = \exp_{p_k}(d^k)$ and compute $X_{q_{k+1}} \in \partial f(q_{k+1})$
- 3 If $f(q_{k+1}) \leq f(p_k) + m\xi^k$, then set $p_{k+1} = q_{k+1}$, else set $p_{k+1} = p_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$e_j^{k+1} = f(x_{k+1}) - f(y_j) - (s^j)^T (x_{k+1} - y_j) \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{x_k\}_{k \in \mathbb{N}}$, $\{y_k\}_{k \in \mathbb{N}}$

The Convex Hadamard Bundle Method

Data: $p_1 = q_1 \in \mathcal{M}$, $g^0 = X_1 \in \partial f(p_1)$, $m \in (0, 1)$, $\varepsilon^0 = c_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k P_{p_k \leftarrow q_j} X_{q_j}, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k c_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $q_{k+1} = \exp_{p_k}(d^k)$ and compute $X_{q_{k+1}} \in \partial f(q_{k+1})$
- 3 If $f(q_{k+1}) \leq f(p_k) + m\xi^k$, then set $p_{k+1} = q_{k+1}$, else set $p_{k+1} = p_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$c_j^{k+1} = f(p_{k+1}) - f(q_j) - \langle X_{q_j}, \log_{q_j} p_{k+1} \rangle \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{x_k\}_{k \in \mathbb{N}}$, $\{y_k\}_{k \in \mathbb{N}}$

The Convex Hadamard Bundle Method

Data: $p_1 = q_1 \in \mathcal{M}$, $g^0 = X_1 \in \partial f(p_1)$, $m \in (0, 1)$, $\varepsilon^0 = c_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k P_{p_k \leftarrow q_j} X_{q_j}, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k c_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $q_{k+1} = \exp_{p_k}(d^k)$ and compute $X_{q_{k+1}} \in \partial f(q_{k+1})$
- 3 If $f(q_{k+1}) \leq f(p_k) + m\xi^k$, then set $p_{k+1} = q_{k+1}$, else set $p_{k+1} = p_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$c_j^{k+1} = f(p_{k+1}) - f(q_j) - \langle X_{q_j}, \log_{q_j} p_{k+1} \rangle + 2\|\log_{q_j} p_{k+1}\| \|X_{q_j}\| \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{x_k\}_{k \in \mathbb{N}}$, $\{y_k\}_{k \in \mathbb{N}}$

The Convex Hadamard Bundle Method

Data: $p_1 = q_1 \in \mathcal{M}$, $g^0 = X_1 \in \partial f(p_1)$, $m \in (0, 1)$, $\varepsilon^0 = c_1^1 = 0$, $k = 1$,
 $J_k = \{1\}$

- 1 Compute a solution $\lambda^k \in \mathbb{R}^{|J_k|}$ of the stabilizing subproblem and set

$$g^k = \sum_{j \in J_k} \lambda_j^k P_{p_k \leftarrow q_j} X_{q_j}, \quad \varepsilon^k = \sum_{j \in J_k} \lambda_j^k c_j^k, \quad d^k = -g^k, \quad \xi^k = -\|g^k\|^2 - \varepsilon^k$$

while not converged do

- 2 Set $q_{k+1} = \exp_{p_k}(d^k)$ and compute $X_{q_{k+1}} \in \partial f(q_{k+1})$
- 3 If $f(q_{k+1}) \leq f(p_k) + m\xi^k$, then set $p_{k+1} = q_{k+1}$, else set $p_{k+1} = p_k$
- 4 Update the index set to J_{k+1} , set $k \leftarrow k + 1$, and set

$$c_j^{k+1} = f(p_{k+1}) - f(q_j) - \langle X_{q_j}, \log_{q_j} p_{k+1} \rangle + 2\|\log_{q_j} p_{k+1}\| \|X_{q_j}\| \quad \text{for all } j \in J_{k+1}$$

- 5 **end**

Result: $\{p_k\}_{k \in \mathbb{N}}$, $\{q_k\}_{k \in \mathbb{N}}$

Convergence

- ▶ In the Euclidean case we have Geiger and Kanzow 2002, Theorem 6.80

Theorem

Let the solution set $S = \{x_ \in \mathbb{R}^n \mid f(x_*) = \inf_{x \in \mathbb{R}^n} f(x)\}$ of the minimization problem be nonempty. Then every sequence $\{x_k\}$ generated by the bundle method algorithm converges to a minimum of f .*

Convergence

- ▶ In the Euclidean case we have Geiger and Kanzow 2002, Theorem 6.80

Theorem

Let the solution set $S = \{x_ \in \mathbb{R}^n \mid f(x_*) = \inf_{x \in \mathbb{R}^n} f(x)\}$ of the minimization problem be nonempty. Then every sequence $\{x_k\}$ generated by the bundle method algorithm converges to a minimum of f .*

- ▶ In the Hadamard case, we obtain an analogous result

Convergence

- ▶ In the Euclidean case we have Geiger and Kanzow 2002, Theorem 6.80

Theorem

Let the solution set $S = \{x_ \in \mathbb{R}^n \mid f(x_*) = \inf_{x \in \mathbb{R}^n} f(x)\}$ of the minimization problem be nonempty. Then every sequence $\{x_k\}$ generated by the bundle method algorithm converges to a minimum of f .*

- ▶ In the Hadamard case, we obtain an analogous result
 - ▶ achieved by enlarging the ε -subdifferential

Convergence

- ▶ In the Euclidean case we have Geiger and Kanzow 2002, Theorem 6.80

Theorem

Let the solution set $S = \{x_ \in \mathbb{R}^n \mid f(x_*) = \inf_{x \in \mathbb{R}^n} f(x)\}$ of the minimization problem be nonempty. Then every sequence $\{x_k\}$ generated by the bundle method algorithm converges to a minimum of f .*

- ▶ In the Hadamard case, we obtain an analogous result
 - ▶ achieved by enlarging the ε -subdifferential
- ▶ Numerically, given $\text{tol} > 0$, we employ the following stopping criterion

$$-\xi^k \leq \text{tol}$$

Implementation

The algorithm is implemented¹ in Julia using `Manopt.jl` (Bergmann 2022) which uses manifolds from `Manifolds.jl` (Axen et al. 2021). A solver call just looks like

```
p_star = bundle_method(M, f, ∂f, p0)
```

where

- ▶ M is a Hadamard manifold
- ▶ f is the objective function
- ▶ ∂f is a subgradient of the objective function
- ▶ p_0 is an initial point on the manifold
- ▶ the parameter for the descent test is set at a default $m = 0.0125$

Numerical Example: \mathcal{H}^4 and $\mathcal{P}(3)$

Let $\mathcal{M}_1 = \mathcal{H}^4$ be the four-dimensional hyperbolic space, and let $\mathcal{M}_2 = \mathcal{P}(3)$ be the space of 3×3 symmetric positive definite matrices. Let $q_1, \dots, q_n \in \mathcal{M}_j$ be $n = 100$ random data points for $j \in \{1, 2\}$.

Numerical Example: \mathcal{H}^4 and $\mathcal{P}(3)$

Let $\mathcal{M}_1 = \mathcal{H}^4$ be the four-dimensional hyperbolic space, and let $\mathcal{M}_2 = \mathcal{P}(3)$ be the space of 3×3 symmetric positive definite matrices. Let $q_1, \dots, q_n \in \mathcal{M}_j$ be $n = 100$ random data points for $j \in \{1, 2\}$. Let $f_{ij}: \mathcal{M}_j \rightarrow \mathbb{R}$ be defined by

$$f_{ij}(p) = \sum_{k=1}^n \text{dist}_{\mathcal{M}_j}^i(p, q_k)$$

Numerical Example: \mathcal{H}^4 and $\mathcal{P}(3)$

Let $\mathcal{M}_1 = \mathcal{H}^4$ be the four-dimensional hyperbolic space, and let $\mathcal{M}_2 = \mathcal{P}(3)$ be the space of 3×3 symmetric positive definite matrices. Let $q_1, \dots, q_n \in \mathcal{M}_j$ be $n = 100$ random data points for $j \in \{1, 2\}$. Let $f_{ij}: \mathcal{M}_j \rightarrow \mathbb{R}$ be defined by

$$f_{ij}(p) = \sum_{k=1}^n \text{dist}_{\mathcal{M}_j}^i(p, q_k)$$

and let $g_j: \mathcal{M}_j \rightarrow \mathbb{R}$ be defined by

$$g_j(p) = \max_{i \in \{1, 2\}} \{f_{ij}(p)\}$$

for $i, j \in \{1, 2\}$.

Numerical Example: \mathcal{H}^4 and $\mathcal{P}(3)$

Let $\mathcal{M}_1 = \mathcal{H}^4$ be the four-dimensional hyperbolic space, and let $\mathcal{M}_2 = \mathcal{P}(3)$ be the space of 3×3 symmetric positive definite matrices. Let $q_1, \dots, q_n \in \mathcal{M}_j$ be $n = 100$ random data points for $j \in \{1, 2\}$. Let $f_{ij}: \mathcal{M}_j \rightarrow \mathbb{R}$ be defined by

$$f_{ij}(p) = \sum_{k=1}^n \text{dist}_{\mathcal{M}_j}^i(p, q_k)$$

and let $g_j: \mathcal{M}_j \rightarrow \mathbb{R}$ be defined by


$$g_j(p) = \max_{i \in \{1, 2\}} \{f_{ij}(p)\}$$

for $i, j \in \{1, 2\}$. We want to solve


$$\arg \min_{p \in \mathcal{M}_j} g_j(p)$$

for $j \in \{1, 2\}$.


Numerical Example: \mathcal{H}^4




Numerical Example: \mathcal{H}^4




Numerical Example: \mathcal{H}^4




Numerical Example: \mathcal{H}^4




Numerical Example: $\mathcal{P}(3)$




Numerical Example: $\mathcal{P}(3)$



Numerical Example: $\mathcal{P}(3)$



Numerical Example: $\mathcal{P}(3)$



Summary and Future Work






What we did:

- ▶ presented the bundle method for geodesically convex functions on Hadamard manifolds
- ▶ touched upon convergence
- ▶ showed two numerical examples

To do:

- ▶ further investigate convergence
- ▶ generic Riemannian manifolds

Bibliography

-  Axen, Seth D. et al. (2021). *Manifolds.jl: An Extensible Julia Framework for Data Analysis on Manifolds*. [arXiv: 2106.08777](#).
-  Azagra, Daniel and Juan Ferrera (2005). "Proximal calculus on Riemannian manifolds". In: *Mediterranean Journal of Mathematics* 2.4, pp. 437–450.
-  Bergmann, Ronny (2022). "Manopt.jl: Optimization on Manifolds in Julia". In: *Journal of Open Source Software* 7.70, p. 3866. DOI: [10.21105/joss.03866](#).
-  Bonnans, J.-F. et al. (2006). *Numerical optimization: theoretical and practical aspects*. Springer-Verlag.
-  Geiger, C. and C. Kanzow (2002). *Theorie und Numerik restringierter Optimierungsaufgaben*. New York: Springer. DOI: [10.1007/978-3-642-56004-0](#).