

Peccei-Quinn mechanism and the quality problem

Based on 1703.01112

Hajime Fukuda (Kavli IPMU, U. Tokyo) with M. Ibe, M. Suzuki, T. T. Yanagida
March 24, 2017

1. Strong CP Problem and the Peccei-Quinn symmetry
2. Big Challenge in the PQ symmetry - Quality problem
3. Our study
 - 3.1 “Gauged” axion

Strong CP Problem and the Solutions

Strong CP Problem

- Yang-Mills theory in 4D has θ vacua, which can be written in Lagrangian: $\Delta\mathcal{L} \sim \theta F\tilde{F}$ (see Seiberg 10)
- Non-zero θ breaks CP symmetry: $|\theta_{QCD}| \lesssim 10^{-10}$
- θ_{QCD} consists of two parts, θ_{YM} and $\arg(\det(Y_u Y_d))$
 - We expect the latter is around $J \sim 10^{-5}$, at least
 - No reason to cancel
- Why is θ_{QCD} so small? - *Strong CP Problem*
- Anthropic principle cannot be used see Ubbaldi 08, for instance

Peccei-Quinn Mechanism

- In a nutshell, “if a symmetry were anomalous in QCD, θ would be unphysical”
- The anomaly never disappear (e.g. Anomaly Matching)
- That anomaly must have low-energy consequence!
 - Massless *colored* Fermion - massless up
 - (For historical reason, this is usually not referred as PQ)
 - Inconsistent with lattice?
 - Pseudo NGB with $\Delta\mathcal{L} \sim \frac{a}{f_a} G\tilde{G}$
 - Peccei and Quinn 1977, Weinberg 1978, Wilczek 1978

- The pseudo NGB is called axion
- $U(1)_{PQ}$ is not exact - the axion, a , has mass
- From the chiral condensation,

$$m_a^2 \sim \frac{m_q \Lambda_{QCD}^3}{f_a^2} \sim \left(1 \text{ meV} \frac{10^9 \text{ GeV}}{f_a} \right)^2$$

- f_a : axion decay constant, $\sim \langle \phi \rangle$

Big Challenge in the PQ symmetry

- Quality problem

Big Obstacle for the Peccei-Quinn Mechanism

- $U(1)_{PQ}$ must be an **extremely** good “symmetry”

$$\Delta\mathcal{L} = c \frac{\phi^5}{M_{\text{Pl}}} \Rightarrow \mathcal{L}_a = -\frac{m_a^2}{2} a^2 + c \frac{f_a^4}{M_{\text{Pl}}} a$$

Since $\theta_{\text{eff}} = \langle a \rangle / f_a$, $c \lesssim 10^{-60}$ for $\theta \lesssim 10^{-10}$!

- *Quality problem*
- So, where does the $U(1)_{PQ}$ come?
 - $U(1)_{PQ}$ is *by definition* not a symmetry
 - Even an exact global symmetry should be broken in Quantum Gravity
 - No anthropic argument!
 - Accidental or/and “anthropic-izing” solution

How Can We Solve It? (using field theory)

- Imposing discrete (gauge) symmetries
 - Discrete symmetries may emerge from gauge symmetry
 - e.g. Abelian Higgs with charge +2
 - For $\Delta\mathcal{L} \sim O(1)\frac{\phi^n}{M_{\text{Pl}}^{n-4}}, n \gtrsim 11!$
- Imposing continuous gauge symmetries
- Heavy QCD Axion

Our Study

- We show two different models as examples
 - “Gauged” axion *HF, Ibe, Suzuki, Yanagida 1703.01112*
 - Heavy axion *Chiang, HF, Harigaya, Ibe, Yanagida 1504.06084, 1602.07909, 1702.00227*

“Gauged” Axion

“Gauged” Axion

- Needless to say, $U(1)_{PQ}$ cannot be gauged
- However, gauge symmetry may protect $U(1)_{PQ}$ *Georgi, Hall, Wise 81*

Gauged $U(1)$ Symmetries and Anomalies

- Chiral $U(1)$ symmetries generally have anomalies
- How about $U(1)_Y$?
 - There, anomalies from *lepton* and *quark* sectors are cancelled!
 - Conversely, “ $U(1)_Y$ ” in each sector are anomalous
 - Of course, due to Higgs, both sector are connected
 - However, what if we have 2HDM? - Weinberg-Wilczek model
 - $B - L$, B and L ? R -parity?
- Generally, multiple “ $U(1)$ ” are assembled into one gauged $U(1)$
- Hence, if a gauged $U(1)$ exists, anomalous $U(1)$ s may well exist

| | $SU(3)_{QCD}$ | $U(1)_g$ | $(U(1)_{PQ})$ |
|----------------------|---------------|----------|---------------|
| ϕ_1 | 1 | 1 | 1 |
| $\psi_{L1,10}$ | 3 | -1 | -1 |
| $\bar{\psi}_{R1,10}$ | $\bar{3}$ | 0 | 0 |
| ϕ_2 | 1 | -10 | 0 |
| ψ_{L2} | 3 | 10 | 0 |
| $\bar{\psi}_{R2}$ | $\bar{3}$ | 0 | 0 |

With interaction: $\Delta\mathcal{L} \sim \phi_1 \bar{\psi}_{R1,10} \psi_{L1,10} + \phi_2 \bar{\psi}_{R2} \psi_{L2}$

- Breaking Operator: $\mathcal{L}_{PQ} = \frac{1}{M_{PL}^7} \phi_1^{10} \phi_2^\dagger + \text{H.c.}$

How to Build Models?

- Building models are really easy
- Introducing two PQ sector, gauging the diagonal group
 - Always, one linear combination of two PQ currents are anomaly-free in QCD
- The problem is how we can suppress cross-terms
- How “realistic” can the model be? - Future work
- What is the cosmological evolution? - Future work

Summary

Summary

- The Peccei-Quinn mechanism is probably the most successful solution to the Strong CP problem
- However, its quality is puzzling
- We have proposed two different way to realize good quality
 - “Gauged” axion
 - Gauge symmetry may protect PQ
 - How realistic does it look? - Future work
 - Cosmological evolution - Future work

Heavy Axion

Heavy Axion


- As I mentioned, m_a is written in terms of f_a , $m_a \sim \sqrt{m_q \Lambda^3} / f_a$
- Okay, let us forget it temporarily
- With PQ-breaking operator $\Delta\mathcal{L} = c\phi^5/M_{\text{Pl}}$,

$$\mathcal{L}_a \sim -\frac{m_a^2}{2}a^2 + c\frac{f_a^4}{M_{\text{Pl}}}a$$

- Thus, if m_a were bigger, we could ignore $\Delta\mathcal{L}$!

How to Make it Heavier?

- Another gauge theory with the same θ is needed



What is the Other Gauge Theory?

- Two way is known
- Use of larger group $SU(N + 3) \supset SU(3)_{QCD} \times SU(N)$ *Dimopoulos 79*
 - Recently Shifman *et al.* 16, used this
 - However, we believe this is almost dead
- Use of a copy of the standard model *Rubakov 94*
 - This is only the possibility

- The axion mass is now $m_a \sim \sqrt{m'_q \Lambda'^3} / f_a$
- If the axion is heavier than $O(1-10) GeV$, we have no constraint for the axion
 - Cosmology like BBN,...
 - Meson Invisible Decay
 - Astrophysics like SN1978A,...
- We may make ν' and Λ' large using \mathbb{Z}_2 breaking
- Since we have introduced a copy, many particles are stable
 - We have found that one of them may be the dark matter
 - How to detect it? - Future work