

How should we tackle the strong CP problem?

Based on 1703.01112 1504.06084 1602.07909 1702.00227

Hajime Fukuda (Kavli IPMU, U. Tokyo)

July 18, 2017

1. Strong CP Problem and the solutions
2. Challenge for the origin of Peccei-Quinn symmetry
3. Our study
 - 3.1 “Gauged” axion
 - 3.2 Heavy axion

Strong CP Problem and the Solutions

Strong CP Problem

- Yang-Mills theory in 4D has θ vacua, which can be written in Lagrangian: $\Delta\mathcal{L} \sim \theta F\tilde{F}$ (see Seiberg 10)
- Non-zero θ breaks CP symmetry: $|\theta_{QCD}| \lesssim 10^{-10}$
- θ_{QCD} consists of two parts, θ_{YM} and $\arg(\det(Y_u Y_d))$
 - We expect the latter is around $J \sim 10^{-5}$, at least
 - No reason to cancel
- Why is θ_{QCD} so small? - *Strong CP Problem*
- Anthropic principle cannot be used see Ubbaldi 08, for instance

- As far as I know, there are roughly two ways
- Spontaneous CP Breaking (e.g. Nelson-Barr Mechanism)
 - Complicated and not so easy *Dine and Draper. 15 for detail*
- Use of anomaly

- Chiral (non-vectorial) U(1) symmetries in 4D generally have anomalies

$$\partial J \sim F\tilde{F}$$

- The chiral rotation only moves θ - θ becomes unphysical!
- This is why we do not have “Weak CP Problem”...

$$\partial J_L \sim W\tilde{W} + F\tilde{F} \quad \therefore \bar{e}_R \rightarrow e^{-i\theta_w/3}\bar{e}_R, L_L \rightarrow e^{i\theta_w/3}L_L, \theta_w \rightarrow 0$$

Peccei-Quinn Mechanism

- In a nutshell, “if a symmetry were anomalous in QCD, θ would be unphysical”
- The anomaly never disappear (e.g. Anomaly Matching)
- That anomaly must have low-energy consequence!
 - Massless *colored* Fermion
 - (For historical reason, this is usually not referred as PQ)
 - Pseudo NGB with $\Delta\mathcal{L} \sim \frac{a}{f_a} G\tilde{G}$
 - Peccei and Quinn 1977, Weinberg 1978, Wilczek 1978

With Massless Colored Fermion

- No additional massless Fermion is allowed in SM
- May *up* be massless? - So called “*massless up*” solution
- Inconsistent with chiral perturbation?
 - Still instanton may save it? *Georgi and McArthur 81*
- (Un)fortunately, lattice people finally concluded it is inconsistent

Axion, the Main Topic!

- Among many ways, the simplest is KSVZ axion

| | $SU(3)_{QCD}$ | $U(1)_{PQ}$ |
|----------------|---------------|-------------|
| ϕ | 1 | 1 |
| ψ_L | 3 | -1 |
| $\bar{\psi}_R$ | $\bar{3}$ | 0 |

- The Lagrangian is $\Delta\mathcal{L} \sim \phi\bar{\psi}_R\psi_L - V(\phi)$
- With the “wine-bottle” $V(\phi)$, $U(1)_{PQ}$ is spontaneously broken and axion appears, with $\Delta\mathcal{L} \sim \frac{a}{f_a} G\tilde{G}$

- $U(1)_{PQ}$ is not exact - the axion, a , has mass
- From the chiral condensation,

$$m_a^2 \sim \frac{m_q \Lambda_{QCD}^3}{f_a^2} \sim \left(1 \text{ meV} \frac{10^9 \text{ GeV}}{f_a} \right)^2$$

- f_a : axion decay constant, $\sim \langle \phi \rangle$

Rich Phenomenology of Axion

- Axion couples with baryons, photons (and leptons)
- Many experiments/observations constrain axions
 - Star lifetime, axion helioscopes, reactors, *etc.*
 - $f_a(\sim \langle \phi \rangle) \gtrsim 10^9 \text{ GeV}$
- Axion coherent oscillation may be the dark matter
 - $\ddot{a}(t) + 3H\dot{a}(t) - m_a^2 a(t) = 0 \Rightarrow \rho_a(t)R(t)^3 = \text{const.}$
 - The initial amplitude of oscillation is $O(1)f_a$
 - For $\Omega h^2 \sim 0.1$, $f_a \sim 10^{12} \text{ GeV}$

Summary So Far

- The Standard Model has the Strong CP problem
- The most plausible (\simeq consistent with other observations) solution is the Peccei-Quinn mechanism and axion
- Rich phenomenology, many things to do!

Challenge for the origin of Peccei-Quinn symmetry

Big Obstacle for the Peccei-Quinn Mechanism

- $U(1)_{PQ}$ must be an **extremely** good “symmetry”

$$\Delta\mathcal{L} = c \frac{\phi^5}{M_{\text{Pl}}} \Rightarrow \mathcal{L}_a = -\frac{m_a^2}{2} a^2 + c \frac{f_a^4}{M_{\text{Pl}}} a$$

Since $\theta_{\text{eff}} = \langle a \rangle / f_a$, $c \lesssim 10^{-60}$ for $\theta \lesssim 10^{-10}$!

- *Quality problem*
- So, *where does the $U(1)_{PQ}$ come?*
 - $U(1)_{PQ}$ is *by definition* not a symmetry
 - Even an exact global symmetry should be broken in Quantum Gravity
 - No anthropic argument!
 - Accidental or/and “anthropic-izing” solution

How Can We Solve It?

- Expecting something in some unknown theory
 - Brane separation, *etc.*
- Using moduli in String Theory
 - Many moduli, many “light” mode
 - Most of them get mass
 - Some of them might be really light
 - The decay constant should be around string / Planck scale
 - Too large for DM, but anthropic arguments may be used
 - This will be confirmed / excluded by BH superradiance

How Can We Solve It? (cont.)

- Imposing discrete (gauge) symmetries
 - Discrete symmetries may emerge from gauge symmetry
 - e.g. Abelian Higgs with charge +2
 - For $\Delta\mathcal{L} \sim O(1)\frac{\phi^n}{M_{\text{Pl}}^{n-4}}, n \gtrsim 11!$
- Imposing continuous gauge symmetries
- Heavy QCD Axion

Our Study

- We show two different models as examples
 - “Gauged” axion *HF, Ibe, Suzuki, Yanagida 1703.01112*
 - Heavy axion *Chiang, HF, Harigaya, Ibe, Yanagida 1504.06084, 1602.07909, 1702.00227*

“Gauged” Axion

“Gauged” Axion

- Needless to say, $U(1)_{PQ}$ cannot be gauged
- However, gauge symmetry may protect $U(1)_{PQ}$ *Georgi, Hall, Wise 81*

Gauged U(1) Symmetries and Anomalies

- Chiral U(1) symmetries generally have anomalies
- How about $U(1)_Y$?
 - There, anomalies from *lepton* and *quark* sectors are cancelled!
 - Conversely, “ $U(1)_Y$ ” in each sector are anomalous
 - Of course, due to Higgs, both sector are connected
 - However, what if we have 2HDM? - Weinberg-Wilczek model
 - $B - L$, B and L ? R -parity?
- Generally, multiple “U(1)” are assembled into one gauged U(1)
- Hence, if a gauged U(1) exists, anomalous U(1)s may well exist

Gauged $U(1)$ and Protection of $U(1)_{PQ}$

- No operators breaking gauged $U(1)$ is allowed
- The problem is the connection between two sector
 - Depending on the charge assignment, PQ breaking operators are suppressed

| | $SU(3)_{QCD}$ | $U(1)_g$ | $(U(1)_{PQ})$ |
|----------------------|---------------|----------|---------------|
| ϕ_1 | 1 | 1 | 1 |
| $\psi_{L1,10}$ | 3 | -1 | -1 |
| $\bar{\psi}_{R1,10}$ | $\bar{3}$ | 0 | 0 |
| ϕ_2 | 1 | -10 | 0 |
| ψ_{L2} | 3 | 10 | 0 |
| $\bar{\psi}_{R2}$ | $\bar{3}$ | 0 | 0 |

With interaction: $\Delta\mathcal{L} \sim \phi_1 \bar{\psi}_{R1,10} \psi_{L1,10} + \phi_2 \bar{\psi}_{R2} \psi_{L2}$

- Breaking Operator: $\mathcal{L}_{PQ} = \frac{1}{M_{\text{PL}}^7} \phi_1^{10} \phi_2^\dagger + \text{H.c.}$

How to Build Models?

- Building models are really easy
- Introducing two PQ sector, gauging the diagonal group
 - Always, one linear combination of two PQ currents are anomaly-free in QCD
- The problem is how we can suppress cross-terms
- How “realistic” can the model be? - Future work
- What is the cosmological evolution? - Future work

Heavy Axion


- As I mentioned, m_a is written in terms of f_a , $m_a \sim \sqrt{m_q \Lambda^3} / f_a$
- Okay, let us forget it temporarily
- With PQ-breaking operator $\Delta\mathcal{L} = c\phi^5/M_{\text{Pl}}$,

$$\mathcal{L}_a \sim -\frac{m_a^2}{2}a^2 + c\frac{f_a^4}{M_{\text{Pl}}}a$$

- Thus, if m_a were bigger, we could ignore $\Delta\mathcal{L}$!

How to Make it Heavier?

- Another gauge theory with the same θ is needed



What is the Other Gauge Theory?

- Two way is known
- Use of larger group $SU(N + 3) \supset SU(3)_{QCD} \times SU(N)$ *Dimopoulos 79*
 - Recently Shifman *et al.* 16, used this
 - However, we believe this is almost dead
- Use of a copy of the standard model *Rubakov 94*
 - This is only the possibility

- The axion mass is now $m_a \sim \sqrt{m'_q \Lambda'^3} / f_a$
- If the axion is heavier than $O(1-10) GeV$, we have no constraint for the axion
 - Cosmology like BBN,...
 - Meson Invisible Decay
 - Astrophysics like SN1978A,...
- We may make ν' and Λ' large using \mathbb{Z}_2 breaking
- Since we have introduced a copy, many particles are stable
 - We have found that one of them may be the dark matter
 - How to detect it? - Future work

Summary

Summary

- The Peccei-Quinn mechanism is probably the most successful solution to the Strong CP problem
- However, its quality is puzzling
- We have proposed two different way to realize good quality
 - “Gauged” axion
 - Gauge symmetry may protect PQ
 - How realistic does it look? - Future work
 - Cosmological evolution - Future work
 - Heavy axion
 - If axion were heavy, quality becomes better
 - Many phenomenology - Future work