



# 1 ar-bgc-argo: Jupyter Notebook templates for 2 searching, downloading, and post-processing 3 biogeochemical Argo float time series

4 **Haruto Fujishima** <sup>1\*</sup> and **Hakase Hayashida** <sup>2\*</sup>




5 <sup>1</sup> Department of Geophysics, Graduate School of Science, Tohoku University, Sendai, Japan <sup>2</sup>

6 Application Laboratory, Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan 

7 Corresponding author \* These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright  
and release the work under a  
Creative Commons Attribution 4.0  
International License ([CC BY 4.0](#))


## 8 Summary

9 ar-bgc-argo is a set of Jupyter Notebook templates that transform raw profiles of biogeochem-  
10 ical Argo (BGC-Argo) floats into “analysis-ready” time series of ocean temperature, salinity,  
11 and biogeochemical variables. Users can search for floats based the dates, geographic region,  
12 and biogeochemical variables of their interests. After downloading the profile time series of  
13 a selected float, ar-bgc-argo can visualize, filter, interpolate, and save the post-processed  
14 time series as a netCDF file. In addition, ar-bgc-argo applies variable-specific treatments and  
15 derive additional oceanographic variables using empirical equations. ar-bgc-argo is designed  
16 to help expand the end users of the growing BGC-Argo float data.

## 17 Statement of need

18 Biogeochemical Argo (BGC-Argo) is a global network of autonomous profiling floats in the  
ocean that has brought dramatic advances in our understanding of ocean biogeochemistry and  
marine ecosystems in recent years ([Thierry et al., 2025](#)). The global BGC-Argo community  
aims to cover the global ocean with 1,000 floats that operate every 10 days or so, monitoring  
22 biogeochemical properties from the sea surface to 2,000 m deep ([Claustre et al., 2020](#)). These  
23 profiling floats are equipped with sensors that can measure up to six key variables: chlorophyll-a,  
24 pH, oxygen, nitrate, irradiance, and suspended particles ([Bittig et al., 2019](#)). As of December  
25 2025, 861 BGC-Argo floats are already in operation, but only 111 of these are equipped with  
26 the six full sensors (Figure 1). The collected profiles are made publicly available in the netCDF  
27 format within a day or so ([Wong et al., 2020](#)).

28 Despite the growing application of BGC-Argo data, the raw profiles include technical errors and  
29 doubtful values because of poor sensor calibration and high sensitivity to noise and artifacts.  
30 Furthermore, the data are unfiltered (containing both good- and bad-quality samples) and  
31 have inconsistent sampling depths among profiles from a given float. These issues necessitate  
32 post-processing prior to scientific analysis, which requires technical knowledge hence, becomes  
33 a time-consuming task.



**Figure 1:** Figure 1: Global coverage of operational BGC-Argo floats as of December 2025 (<https://www.ocean-ops.org/share/Argo/Maps/bgc.png>; accessed on January 16, 2026)

## Overview of ar-bgc-argo

ar-bgc-argo consists of three Jupyter Notebook templates: `search.ipynb`; `download.ipynb`; and `generate.ipynb`. To use these templates, users create a copy of the template of their interest, modify its inputs, and run it on their Jupyter environment.

### search.ipynb

`search.ipynb` searches for BGC-Argo floats from the synthetic-profile index file (`argo_synthetic-profile_index.txt`) of the Global Data Assembly Center (GDAC; [bittig2019]) and based on the user inputs, including the temporal and spatial coverages and the biogeochemical variables of interest. In addition, `search.ipynb` allows users to narrow down the float selection based on three key criteria:

- `mindays`: the minimum duration of the data record to ensure sufficient temporal coverage (e.g., at least 365 days).
- `minfreq`: the minimum sampling frequency required to capture temporal variability (e.g., at least every 14 days).
- `maxdrift`: the float's maximum drift speed (e.g., 0.05 m/s), which is particularly useful for identifying “quasi-Eulerian” floats suitable for one-dimensional modeling (e.g., [Bruggeman et al., 2024]).


The trajectories of all qualified floats based on the search criteria are drawn on a map and their temporal coverages are visualized on a time series. These visualizations provide an intuitive overview, enabling users to identify potential spatial and temporal biases in observational coverage within the study region of interest prior to data retrieval ([Hayashida et al., 2025]).

## 55 download.ipynb

56 download.ipynb retrieves the concatenated synthetic-profiles time series of a selected float  
57 ([wmoid]\_Sprof.nc, where [wmoid] is the seven-digit World Meteorological Organization Identifier or WMO ID). While download.ipynb naturally follows the selection made in search.ipynb,  
58 it can also be used independently if the WMO ID of the target float is already known. Upon  
59 execution, download.ipynb creates a directory named after the WMO ID to store the data.  
60 It then identifies the correct file path from the synthetic-profile index file and downloads the  
61 data from one of the two GDACs.  
62

## 63 generate.ipynb

64 generate.ipynb is the core component of ar-bgc-argo, designed to transform raw BGC-Argo  
65 profiles into “analysis-ready” time series suitable for immediate scientific application. The  
66 data processing workflow consists of eight steps (Figure 2). At every step, diagnostic plots  
67 are generated and enable users to visually verify the reliability of the post-processing, which  
68 prevents a “black-box” approach and ensuring the production of high-quality datasets.



**Figure 2:** Figure 2: Schematic workflow of the data-processing pipeline implemented in generate.ipynb. This example assumes a full-sensor float with an arbitrary WMO ID (1234567)

## 69 Acknowledgements

70 BGC-Argo data are freely available through one of the two Global Data Assembly Centers  
71 (GDAC), using the WMO number of the float, which is its specific identifier. We thank  
72 Hidehiro Fujio for computational assistance and Pete Strutton, Clara Vives, Kanako Sato,  
73 Yoshimi Kawai for insightful discussion on BGC-Argo floats and feedback on earlier versions of  
74 ar-bgc-argo. This work was supported by JSPS KAKENHI Grant Number JP24H02226 and  
75 the Nakajima Foundation.

## 76 References

77 Bittig, H. C., Maurer, T. L., Plant, J. N., Schmechtig, C., Wong, A. P. S., Claustre, H., Trull, T.  
78 W., Udaya Bhaskar, T. V. S., Boss, E., Dall’Olmo, G., Organelli, E., Poteau, A., Johnson,  
79 K. S., Hanstein, C., Leymarie, E., Le Reste, S., Riser, S. C., Rupan, A. R., Taillandier, V.,

- 80 ... Xing, X. (2019). A BGC-Argo Guide: Planning, Deployment, Data Handling and Usage.  
81 *Frontiers in Marine Science*, 6. <https://doi.org/10.3389/fmars.2019.0050>
- 82 Bruggeman, J., Bolding, K., Nerger, L., Teruzzi, A., Spada, S., Skákala, J., & Ciavatta,  
83 S. (2024). EAT v1.0.0: A 1D test bed for physical–biogeochemical data assimilation in  
84 natural waters. *Geoscientific Model Development*, 17, 5619–5639. [https://doi.org/10.](https://doi.org/10.5194/gmd-17-5619-2024)  
85 [5194/gmd-17-5619-2024](https://doi.org/10.5194/gmd-17-5619-2024)
- 86 Claustre, H., Johnson, K. S., & Takeshita, Y. (2020). Observing the Global Ocean with  
87 Biogeochemical-Argo. *Annual Review of Marine Science*, 12, 23–48. [https://doi.org/10.](https://doi.org/10.1146/annurev-marine-010419-010956)  
88 [1146/annurev-marine-010419-010956](https://doi.org/10.1146/annurev-marine-010419-010956)
- 89 Hayashida, H., Strutton, P. G., Hashihama, F., Horimoto-Miyazaki, N., Katano, T., &  
90 Yasui-Tamura, S. (2025). Status and prospects of biogeochemical Argo observations of  
91 chlorophyll-a in the northwest Pacific. *Journal of Oceanography*, 81, 289–299. [https:](https://doi.org/10.1007/s10872-025-00750-2)  
92 [//doi.org/10.1007/s10872-025-00750-2](https://doi.org/10.1007/s10872-025-00750-2)
- 93 Thierry, V., Claustre, H., Pasqueron de Fommervault, O., Zilberman, N., Johnson, K. S., King,  
94 B. A., Wijffels, S. E., Bhaskar, U. T. V. S., Balmaseda, M. A., Belbeoch, M., Bollard,  
95 M., Boutin, J., Boyd, P., Cancouët, R., Chai, F., Ciavatta, S., Crane, R., Cravatte, S.,  
96 Dall'Olmo, G., ... Zuo, H. (2025). Advancing ocean monitoring and knowledge for societal  
97 benefit: The urgency to expand Argo to OneArgo by 2030. *Frontiers in Marine Science*,  
98 12. <https://doi.org/10.3389/fmars.2025.1593904>
- 99 Wong, A. P. S., Wijffels, S. E., Riser, S. C., Pouliquen, S., Hosoda, S., Roemmich, D., Gilson,  
100 J., Johnson, G. C., Martini, K., Murphy, D. J., Scanderbeg, M., Bhaskar, T. V. S. U.,  
101 Buck, J. J. H., Merceur, F., Carval, T., Maze, G., Cabanes, C., André, X., Poffa, N., ...  
102 Park, H.-M. (2020). Argo Data 1999–2019: Two Million Temperature-Salinity Profiles and  
103 Subsurface Velocity Observations From a Global Array of Profiling Floats. *Frontiers in*  
104 *Marine Science*, 7, 700. <https://doi.org/10.3389/fmars.2020.00700>