



Histopathology Inflammation

3rd class

Lab 3

2026-2027

FIBMS. Pathology
Dr.Duha Talib Kareem

M.Sc. Pathology
Dr.ahmed A. Hussein

M.Sc. pathology
Dr. Mahdi Jasim

learning objectives

Recall the cardinal signs of inflammation (such as the PRISH acronym) and the .1
.fundamental definitions distinguishing acute from chronic inflammation

Explain the three major components of acute inflammation and describe the .2
.physiological differences between an exudate and a transudate

Classify the specific morphological patterns of acute inflammation—including .3
serous, fibrinous, suppurative (purulent) inflammation, and ulceration—based on
.their gross and microscopic characteristics

Differentiate between acute and chronic inflammatory responses by comparing .4
their cellular infiltrates, tissue injury patterns, and systemic manifestations in
.conditions such as acute appendicitis, chronic bronchitis, and rheumatoid arthritis

Assess the histological features of granulomatous inflammation to distinguish .5
between infectious caseating granulomas (such as in tuberculosis) and foreign-body
.granulomas



Inflammation



:Inflammation

is a response of vascularized tissues to infections and tissue damage that brings cells and molecules from the circulation to the sites where they are needed, to eliminate .the offending agents





Types of inflammation

Acute inflammation: The initial, rapid response to infections and .tissue damage

- It typically develops within minutes or hours.
- is of short duration, lasting for several hours or a few days.
- Its main characteristics:
 - edema: the exudation of fluid and plasma proteins.
 - the emigration of leukocytes, predominantly neutrophils.



:Chronic inflammation

- may follow acute inflammation or arise de novo.
- Characterized by:
 - It is of longer duration.
 - is associated with more tissue destruction.
 - the presence of lymphocytes and macrophages.
 - the proliferation of blood vessels.
 - fibrosis.


Table 3.1 Features of Acute and Chronic Inflammation

Feature	Acute	Chronic
Onset	Fast: minutes or hours	Slow: days
Cellular infiltrate	Mainly neutrophils	Monocytes/macrophages and lymphocytes
Tissue injury, fibrosis	Usually mild and self-limited	May be severe and progressive
Local and systemic signs	Prominent	Less

Signs of inflammation

English	Latin	Cause
Heat	<i>Calor</i>	Vasodilation
Redness	<i>Rubor</i>	Vasodilation
Swelling	<i>Tumor</i>	Increased vascular permeability Increased granulation tissue
Pain	<i>Dolor</i>	Physical and chemical stimulation of nociceptors
Loss of function	<i>Functio laesa</i>	Pain Reflex muscle inhibition Disruption of tissue structure Fibroplasia and metaplasia

The external manifestations of inflammation
:(cardinal signs) are



Acute inflammation

Acute inflammation

:has three major components

- .Dilation of small vessels: leading to an increase in blood flow .1
- .Increased vascular permeability: enabling plasma proteins and leukocytes to leave the circulation .2
- .Emigration of the leukocytes from the microcirculation .3
- .Accumulation in the focus of injury •
- .Activation to eliminate the offending agent •

Exudation : is the escape of fluid, proteins, and blood cells from the vascular system into – interstitial tissues or body cavities. Its presence implies that there is an increase in the permeability .small blood vessels

Transudate: • is a fluid with low protein content, • little or no cellular material • It is an ultrafiltrate – of blood plasma. • is produced as a result of vessels. osmotic or hydrostatic imbalance across .normal vascular permeability

Acute inflammation



Redness (rubor) and swelling (tumour), both characteristic features of acute inflammation

Acute appendicitis




This is the normal appearance of the appendix against the background of the cecum




acute appendicitis with yellow to tan exudate and hyperemia, including the periappendiceal fat superiorly, rather than a smooth, glistening pale tan serosal surface.


Normal appendix




Acute appendicitis



Acute appendicitis



Microscopically, acute appendicitis is marked by mucosal inflammation and necrosis. Note the remnant of mucosa at the left, and the ulceration of the entire mucosa from the extensive necrosis.



Here, the mucosa shows ulceration and undermining by an extensive neutrophilic exudate.



MORPHOLOGIC PATTERNS OF ACUTE INFLAMMATION



Serous inflammation:

- Is the outpouring of a watery, relatively protein-poor fluid ,
depending on the site of injury.
- Typically, the fluid :
 1. is not infected.
 2. does not contain large numbers of leukocytes.

Serous inflammation caused by a burn

Serous inflammation



Serous inflammation:




Second-degree burns manifest as erythema with superficial blistering of the skin.

Serous inflammation:



Contact dermatitis blisters from the poison by toxic material

- 
- 
- The skin blister resulting from a burn or viral infection:
 - a serous effusion either within or immediately beneath the epidermis of the skin .

Serous inflammation:




Fig. 3.12 Serous inflammation. Low-power view of a cross section of a skin blister showing the epidermis separated from the dermis by a focal collection of serous effusion.



Fibrinous inflammation:

- occurs as a consequence of more severe injuries, resulting in greater vascular permeability that allows large molecules (such as fibrinogen) to pass the endothelial barrier.
- Such exudates may be ended with:

1. Resolution
2. Organization & scarring:

A fibrinous exudate is characteristic of inflammation in the lining of .body cavities, such as the meninges, pericardium

Fibrinous inflammation:





Figure 2–12 Fibrinous pericarditis. **A**, Deposits of fibrin on the pericardium. **B**, A pink meshwork of fibrin exudate (*F*) overlies the pericardial surface (*P*).

Fibrinous inflammation:



The pericardial surface is dry with a coarse granular appearance caused by fibrinous exudate



In fibrinous inflammation, the site of injury would contain cells mixed with fibrous exudate, further enriching the extracted fluid with inflammatory cells and additional proteins

Fibrinous inflammation:



This is a higher-power photomicrograph demonstrating fronds of fibrin (arrows) projecting from the surface of the pericardium



Histologic examination of the pericardial sac showed a fibrinous exudate containing patchy infiltrates of neutrophils, with plump reactive mesothelial cells.



Suppurative (purulent) inflammation & abscess formation.

- These are manifested by the collection of large amounts of purulent exudate (pus) consisting of neutrophils, necrotic cells, and edema fluid.
- Certain organisms (e.g., staphylococci) are more likely to induce such localized suppuration and are therefore referred to as pyogenic (pus-forming).
- Abscesses: are localized collections of pus caused by buried in a tissue, an organ , or a confined space suppuration . They are produced by seeding of pyogenic bacteria into a tissue.

Suppurative (purulent) inflammation

Swollen appendix with congested blood vessels and covered by diffuse yellowish suppurative exudate

Diagnosis:

Acute suppurative appendicitis



Suppurative wounds inflammation caused by inadequate cleaning and improper care may lead to infection.

Suppurative (purulent) inflammation of the lung





Figure 2-13 Purulent inflammation with abscess formation. **A**, Multiple bacterial abscesses in the lung (*arrows*) in a case of bronchopneumonia. **B**, The abscess contains neutrophils and cellular debris and is surrounded by congested blood vessels.


Suppurative (purulent) inflammation of skin



Suppurative inflammation with abscess and acute inflammation was noted in deep dermis of the skin with area of ulceration




An ulcer:

- A local defect, or excavation, of the surface of an organ or tissue that is produced by necrosis of cells and sloughing (shedding) of necrotic and inflammatory tissue.
 - Ulceration can occur only when tissue necrosis & resultant inflammation exist on or near a surface.
- 

ulcer




A mouth ulcer is a small, painful ulcer that forms inside the oral cavity or on the lips. Although they are common and usually not serious, they can cause considerable discomfort




Features of peptic ulcers on OGD (A) peptic ulcer located in the gastric antrum (B) haemorrhaging gastric ulcer

Stomach - Ulcer



This is a low-power photomicrograph of a section through the transected ulcer. (1) The blue cells on the right hand side of this section are the normal gastric epithelial cells of the mucosa. (2) Note the absence of any epithelial cells within the crater of the ulcer.



A portion of the epithelium is absent, exposing the underlying submucosa.

ulcer



Venous leg ulcers are wounds or sores that can develop between knee and ankle joints. They often begin as shallow wounds with uneven edges that drain or weep a lot. Pain can vary




CHRONIC INFLAMMATION


CHRONIC INFLAMMATION

- Chronic inflammation is the cause of tissue damage in some of the most common and disabling human diseases, such as:
 1. Rheumatoid arthritis.
 2. Atherosclerosis.
 3. Tuberculosis.
 4. Chronic lung diseases.
 5. infiltration with mononuclear cells:—
 - . Macrophages
 - .Lymphocytes
 - .plasma cells


Chronic inflammation ,Rheumatoid arthritis




chronic autoimmune disease causing systemic inflammation that destroys joints and leads to permanent deformities, damages the joint lining, cartilage, and bones, resulting in painful swelling, stiffness, loss of function (deformity).

- 
- 
- It frequently begins from the outset as a chronic (chronic inflammation ab initio), insidious, and low-grade response.
 - Chronic Inflammation may complicate acute inflammation which is almost always a suppurative type of inflammation that presents as a purulent discharge (pus) as seen in abscess.

Rheumatoid arthritis



Patients with rheumatoid arthritis often have chronic synovitis with a lymphoplasmacytic infiltrate also with lymphoid aggregates.




Patients have rheumatoid arthritis with joint have outer areas of fibrosis can be quite prominent.

chronic bronchitis




chronic inflammation of the bronchi in the distal lung parenchyma has led to dilation and scarring with increased tan to white collagenous tissue.



This picture shows a bronchus with increased numbers of chronic inflammatory cells in the submucosa.

chronic inflammation in the lung




The three characteristic features of chronic inflammation (in the lung)

1-Chronic inflammatory cells infiltration

2-destruction of the normal tissue (normal alveoli are replaced by spaces lined by cuboidal cells (arrow heads)

3. Replacement by fibrosis (arrows)

chronic cervicitis



Here is chronic cervicitis in the squamous epithelium at the right, with non-inflamed non-keratinizing squamous mucosa at the left



Normal



Acute
cholecystitis




Chronic and acute inflammation
with empyema

chronic cholecystitis




Chronic Calculous Cholecystitis: The gallbladder lumen is filled with numerous mixed cholesterol stones. The wall is irregularly thickened due to inflammation and fibrosis. There is focal transmural necrosis.



The histologic hallmarks of chronic calculous cholecystitis are: mononuclear inflammatory infiltrate in the lamina propria, fibrosis, and metaplastic changes in the mucosa. This low magnification view shows inflammatory infiltrate extending through the muscularis to subserosal tissues. There is considerable thickening of the wall due to inflammation and fibrosis.



GRANULOMATOUS INFLAMMATION

- This is a distinctive pattern of chronic inflammatory reaction characterized by focal accumulations of activated macrophages, which often develop an epithelioid (epithelial-like) appearance.
- 



Granuloma

- A granuloma is a focus of chronic inflammation consisting of a microscopic aggregation of macrophages that are transformed into epithelioid cells surrounded by a collar of mononuclear leukocytes, principally lymphocytes and occasionally plasma cells.
 - Granulomas associated with certain infectious organisms (Mycobacterium tuberculosis) often contain a central zone of necrosis.
- 

- 
- The epithelioid cells have a pale pink granular cytoplasm with indistinct cell borders and a vesicular nucleus that is oval or elongate.
 - Older granulomas develop an enclosing rim of fibroblasts and connective tissue.
 - Frequently, epithelioid cells fuse to form giant cells in the periphery or in the center of granulomas.
 - They have a large mass of cytoplasm containing 20 or more small nuclei arranged either peripherally (**Langhans-type giant cell**) or haphazardly (**foreign body-type giant cell**).

Diagram of typical TB granuloma


Multinucleated GC

Epithelioid cells

Caseation

Lymphocytes


Fibroblasts



Granulomatous disease (lung)




Here are numerous confluent granulomas in upper lung fields in a case of active pulmonary tuberculosis. the granulomatous inflammation can lead to caseous necrosis




Grossly, a granuloma tends to be a focal lesion. Seen here in a hilar lymph node is a granuloma. Granulomas due to infectious agents such as mycobacteria are often described as "caseating" when they have prominent caseous necrosis.

Granuloma. lung




there are scattered granulomas in the parenchyma.




Giant cells are a "committee" of epithelioid macrophages. Seen here are two Langhans type giant cells in which the nuclei are lined up around the periphery of the cell. Additional pink epithelioid macrophages compose most of the rest of the .granuloma

pulmonary granuloma



Here is a pulmonary granuloma involving a bronchiole. Granulomatous inflammation typically consists of mixtures of cells including epithelioid macrophages, Langhans giant cells, lymphocytes, plasma cells, and fibroblasts. There may even be some neutrophils




This is a caseating granuloma. Epithelioid cells surround a central area of necrosis that appears irregular, amorphous, and pink. Grossly, areas of caseation appear cheese-like.

Foreign body granulomas

- Foreign body granulomas form when material that can not be phagocytosis by a single macrophage such as:
 1. Talc.
 2. Sutures.
 3. fibers
- Epithelioid cells and giant cells form and are apposed to the surface of the foreign body and/or actually include it.
- The foreign material can usually be identified in the center of the granuloma.

Foreign body granuloma




Two foreign body giant cells are seen, where there is a bluish strand of suture material (arrow) from a previous operation



Here is a foreign body type giant cell at the upper left of center adjacent to a segment of vegetable material aspirated into the lung (Note the thick cell walls of the plant material). Such foreign body giant cells have nuclei scattered haphazardly about the cell.

Foreign body granuloma



Foreign body giant cells were attached to the silk implants



Good luck