

ANATOMY OF FOREARM

DR. AHMED AL-BAGHDADI, *Ph.D*

THE FOREARM

- BONE
- FASCIA
- ANTERIOR COMPARTMENT
- POSTERIOR COMPARTMENT
- JOINTS
- SPECIFIC AREAS


BONES


Humerus Bone

□ Distal Region


contains some important bony landmarks such as

- ✓ **supraepicondylar ridges** (medial , Lateral)
- ✓ **epicondyles**
 - medial: 1- Gives origin to superficial flexor muscles of forearm
2- Grooved posteriorly by ulnar nerve
 - Lateral: Gives origin to superficial extensor muscles of forearm
- ✓ **trochlea.**
- ✓ **Capitulum**
- ✓ **fossae :**
 - 1) coronoid,
 - 2) radial
 - 3) olecranon






The medial epicondyle (7) is more prominent than the lateral (5).
The medial part of the trochlea (13) is more prominent than the lateral part.
The olecranon fossa (10) on the posterior surface is deeper than the radial and coronoid fossae on the anterior surface (12 and 3).



Bones Of the forearm

□ ulna

- 1) Proximal Portion of the Ulna
- 2) Shaft of the Ulna
- 3) Distal Portion of the Ulna



Bones Of the forearm

□ ulna


1) Proximal Portion of the Ulna

- ✓ articulates with the trochlea of the humerus
- ✓ Important landmarks

□ Olecranon

- Separated from skin by olecranon bursa

- Coronoid process is distal attachment of brachialis muscle
- Radial notch is lateral facet for head of radius at proximal radioulnar joint





The trochlear notch (11) faces forwards, with the radial notch (9) on the lateral side.

Bones Of the forearm

□ Ulna

3) Distal Portion of the Ulna

- much smaller in diameter
- rounded head,
- **styloid process.**
- The head articulates with the **ulnar notch** of the radius to form the distal radio-ulnar joint



Bones Of the forearm

Radius

- lies laterally and parallel to ulna,
- pivots around the ulna to produce **movement** at the proximal and distal radio-ulnar joints.
- articulates in four places:

1) Elbow Joint

formed by an articulation between the head of the radius, and the capitulum of the humerus.

2) Proximal Radioulnar Joint


between the radial head, and the radial notch of the ulna.

3) Wrist Joint

between the distal end of the radius and the carpal bones.

4) Distal Radioulnar Joint

between the ulnar notch and the head of the ulna.



Bones Of the forearm

Radius

1) proximal end of the radius

a. Head

(1) Articulates proximally with capitulum and medially with radial notch of ulna at proximal radioulnar joint

(2) Encircled by strong annular ligament except at radial notch of ulna

b. Neck


- Related to deep radial nerve as it pierces supinator muscle

c. Tuberosity

- Insertion of biceps brachii tendon, just lateral to brachial artery

2) Shaft of the Radius

- **triangular** in shape,
- three borders and three surfaces.
- In the middle of the lateral surface, there is a small roughening for the attachment of the **pronator teres** muscle






Bones Of the forearm

Radius

3) Distal Region of the Radius


- Expanded end.
- **styloid process**
- **ulnar notch**, which articulates with the head of ulna, forming the distal **radioulnar joint**.
- distal surface of the radius has two facets, for articulation with **scaphoid** and **lunate** carpal bones. This makes up the **wrist joint**



The lower end of the radius is concave anteriorly (at the lower label 1 in E), with the ulnar notch medially (G13) and the dorsal tubercle on the posterior surface (F2).


Dorsal tubercle of lister

Act as a pulley for **EPL** tendon before tendon pivots & turns obliquely to insert onto distal phalanx of thumb



a

Lister's
tubercle

**b**

Lister's
tubercle



Bones Of the hand

bones of the hand(**27**) can be divided into **three** categories:

- **Carpal bones** (Most proximal)


A set of eight irregularly shaped bones. These are located in the wrist area.

- **Metacarpals**

There are five metacarpals, each one related to a digit

- **Phalanges** (Most distal)

The bones of the fingers. Each finger has three phalanges, except for the thumb, which has two



Bones Of the hand


Carpal Bones


organised into two rows – proximal and distal.

- In the **proximal** row, the bones are (lateral to medial):
 - ✓ Scaphoid
 - ✓ Lunate
 - ✓ Triquetrum
 - ✓ **Pisiform** –
- In the **distal** row, the bones are (lateral to medial):
 - ✓ Trapezium
 - ✓ Trapezoid
 - ✓ Capitate
 - ✓ Hamate – has a projection on its palmar surface called the hook of hamate

N.B


- 1) Proximally, the scaphoid and lunate articulate with the radius to form the wrist joint.
- 2) In the distal row, all of them, articulate with proximal row of carpal bones at **midcarpal joint** and with metacarpal bones at **metacarpophalangeal joints**





Note

Distal end of ulnar Separated from carpal bones by articular disc (triangular ligament) of radiocarpal joint






Fascia

Fascia


- forearm is enclosed in a sheath of deep fascia (antebrachial fascia).
- from which fascial intermuscular septa extend to bone
- deep fascia, with interosseous membrane and intermuscular septa divide the forearm into compartments, each having its own muscles, nerves, and blood supply
- **anterior compartment** contains the structures on the front of the forearm
- **posterior compartment** contains the structure on the back of the forearm




Fascia

- Near the wrist, the deep fascia presents two localized thickenings, which retain the digital tendons in position during hand movements.


1) Flexor Retinaculum




2) Extensor Retinaculum,



Flexor Retinaculum



2) Extensor Retinaculum



ANTERIOR COMPARTMENT

FLOXOR COMPARTMENT

Muscles of the anterior compartment

Arranged In two layers:

1- A superficial layer **(5)**

2- A deep layer **(3)**

Nerves of the anterior compartment

3 nerves

- ☐ The **median nerve** passes between the 2 heads of pronator teres.
- ☐ The **ulnar nerve** passes between the 2 heads of flexor carpi ulnaris.
- ☐ **The anterior interosseous nerve** , branch from median nerve

arterial supply

- ☐ comes from **the common interosseous branch of the ulnar artery** .
- ☐ This artery divides into the **anterior and posterior interosseous** arteries.



IV. Muscles of Forearm.

A. Muscles of Front of Forearm (Flexors).

Muscle	Origin	Insertion	Action.
1- Pronator teres	a) Humeral head: from lower part of medial supracondylar ridge & C.F.O. b) Ulnar head: medial border of coronoid process.	Rough area in the middle of lateral surface of radius (Pronator tuberosity)	Flexes elbow & pronation of forearm at radioulnar joints (main action).
2- Flexor carpi radialis	C.F.O.	Base of 2 nd & 3 rd metacarpal bones.	Flexes elbow, flexion & radial deviation of the wrist. This muscle with number 5 are main flexors of wrist
3- Palmaris longus	C.F.O, may be absent. It has a long tendon which can be used as a tendon graft.	Palmar aponeurosis	Common action.
4- Flexor digitorum superficialis	a) Humero-ulnar head: C.F.O. & medial border of coronoid process. b) Radial head: anterior oblique line of radius i.e. it has 2 heads arising from 3 bones.	By 4 tendons into sides of middle phalanges of medial 4 fingers	Flexes elbow, wrist, metacarpophalangeal & proximal interphalangeal joints of medial 4 fingers
5- Flexor carpi ulnaris	a) Humoral head: C.F.O. b) Ulnar head : From medial border of olecranon & upper 2/3 of posterior border of ulna	Pisiform bone & base. of 5th metacarpal bone	Flexes elbow, flexion & ulnar deviation of the wrist (main action). One of main flexor of wrist .

IV. Muscles of Forearm.

A. Muscles of Front of Forearm (Flexors).

	Origin	Insertion	Action.
1- Flexor digitorum profundus	Upper 2/3 of anterior & medial surface of shaft of ulna & interosseous membrane.	By 4 tendons into base of terminal phalanges of medial 4 fingers.	Flexion of wrist & all joints of medial 4 fingers.
2- Flexor pollicis longus.	Upper 2/3 of anterior surface of shaft of radius & interosseous membrane .	Base of terminal phalanx of thumb.	Flexion of wrist & all joints of thumb.
3- Pronator quadratus.	Lower 1/4 of front of ulna	Lower 1/4 of front of radius	Pronation at radioulnar joints .

Superficial Group

- 1) Pronator teres(PT)
- 2) Flexor carpi radialis(FCR)
- 3) Palmaris longus(PL)
- 4) Flexor carpi ulnaris(FCU)
- 5) Flexor Digitorum Superficialis(FDS)


NOTE

1) origin\

- **ALL have common origin** from the medial epicondyle of the humerus
- Some of them have additional origin from other bone

2) **All of are supplied** by median n. except FCU

3) **All of them help in flexion of elbow joint**, moreover, each muscle has its specific action indicated by its name




-  Flexor carpi ulnaris
-  Palmaris longus
-  Flexor carpi radialis
-  Pronator teres


Deep Group



- 1) Flexor digitorum profundus(FDP)
- 2) Flexor pollicis longus(FPL)
- 3) Pronator quadratus(PQ)


NOTE

- All of are supplied by ANTERIOR INTEROSSEOUS NERVE branch from median nerve
- In addition to that **FDP** has additional supply from ulnar nerve













- A lesion of median nerve at or above elbow paralyzes pronator quadratus & pronator teres muscles, with loss of ability to pronate the forearm.
- When supinator muscle is paralyzed from a radial nerve lesion, the biceps brachii muscle can still supinate the forearm.





Radial artery



- 1- Courses under brachioradialis to distal forearm
- 2- Lies between **flexor carpi radialis** & **brachioradialis** tendons near wrist (where pulse is taken)
- 3- Winds dorsally **through anatomical snuffbox** to reach dorsum
enters palm by passing between two heads of **first dorsal interosseous muscle**
- 4- In hand, anastomose with ulnar a., by forming two arches, superficial palmar arch & deep palmar arch.



Ulnar artery

1- In middle third of forearm is joined by ulnar nerve deep to flexor carpi ulnaris, and then distally courses lateral to flexor carpi ulnaris tendon

2- Passes with ulnar nerve lateral to pisiform & superficial to flexor retinaculum in **Guyon canal** to reach hand with ulnar nerve




Common interosseous artery

- ❑ Short branch from ulnar artery
- ❑ divides into anterior & posterior

interosseous arteries

Anterior interosseous artery descends with anterior interosseous nerve to supply forearm flexor muscles.

Posterior interosseous artery enters posterior compartment & descends with posterior interosseous nerve to supply forearm extensor muscles



Median nerve

❑ In the cubital fossa

- ❖ it runs on the medial side of the brachial a.
- ❖ It descends between the 2 head of pronator teres

❑ In the forearm

- ❖ runs between **FDS& FDP**.
- ❖ Its lower part, emerges from the lateral side of **FDS**
- ❖ becomes superficial (or covered by **PL** tendon).


❑ In the hand

- It enters deep to the flexor retinaculum.

❑ Branches in forearm

- 1) All mm of anterior compartment except FCU & 1/2 FDP
- 2) Articular branches:
- 3) **Anterior interosseus nerve**: supply mm of deep group
- 4) **Palmar cutaneous** branches:

- passes superficial to the flexor retinaculum
- supplies skin of the lateral 2/3 of the palm.



Ulnar Nerve

❑ At cubital fossa

- Passes behind medial epicondyle.
- Enters anterior compartment of forearm by passing between two heads of FCU

❑ In forearm


- Runs on medial side of ulnar artery .
- In between FDP & FDS

❑ At wrist

- **Enters hand superficial to flexor retinaculum**

❑ Supply

- 1) **Muscular branches** to 1 & 1/2 mm. (**FCU & 1/2 FDP**)
- 2) **Articular branches**: to the elbow joint.
- 3) **Palmar cutaneous branch**:
 - ❖ passes superficial to the flexor retinaculum
 - ❖ supplies skin of the medial 1/3 of the palm.
- 4) **Dorsal cutaneous branch**:
 - ❖ it supplies skin of medial 1/3 of dorsal surface of hand & dorsal surface of medial 1.5 fingers (except nail beds).



POSTERIOR COMPARTMENT

EXTENSOR COMPARTMENT

B. Muscles of Back of Forearm (Extensors).

a. Superficial group

Muscle	Origin	Insertion	Action
1- Brachioradialis.	Upper 2/3 of lateral supracondylar ridge	Lateral surface of radius above the styloid process.	Initiate pronation & supination to midprone position at radioulnar joints. Flexion of elbow in the midprone position. •
2- Extensor carpi radialis longus	Lower 1/3 of lateral supracondylar ridge	Base of 2 nd. metacarpal bone	Common action + radial deviation (Abduction) of wrist.
3- Extensor carpi radialis brevis.	C.E.O.	Base of 3 rd. metacarpal bone	As No. 2.
4- Extensor digitorum.	C.E.O.	By 4 tendons into extensor expansion on dorsum of proximal phalanges of medial 4 fingers.	Common action + extension of metacarpo- phalangeal joints of medial 4 fingers.
5- Extensor digiti minimi.	C.E.O.	Extensor expansion of little finger.	Common action + Extensor of metacarpo- phalangeal joints of little finger.
6. Extensor carpi ulnaris.	C.E.O.	Base of 5 th. metacarpal bone	Common action + ulnar deviation (Adduction) of the wrist.
7. Anconeus	• Separately from lateral epicondyle.	• Lateral aspect of back of olecranon	• Assists triceps in extension of elbow.

B. Muscles of Back of Forearm (Extensors).

a. deep group

Muscles	Origin	Insertion	Action.
1- Supinator	Supinator fossa & supinator crest of ulna.	Anterior, lateral & posterior surface of upper 1/3 of radius	supination of forearm at radioulnar joints.
2- Abductor pollicis longus	Posterior surface of radius , ulna & interosseous membrane.	Base of 1st . metacarpal bone.	Abduction of thumb .
3- Extensor pollicis brevis	Posterior surface of radius & interosseous membrane.	Base of proximal phalanx of thumb.	Extension of proximal phalanx of thumb.
4- Extensor pollicis longus	Posterior surface of ulna & interosseous membrane.	Base of terminal phalanx of thumb.	Extension of all joints of thumb.
5- Extensor indicis	Posterior surface of ulna.	Extensor expansion of index finger.	Extension of proximal phalanx of index finger.

Muscles of the posterior compartment

- Arranged In two layers:

1- A superficial layer (7)


2- A deep layer (5)

Nerve of the anterior compartment

- ❑ The posterior interosseous nerve , branch from radial nerve

Arterial supply

- ❑ posterior interosseous artery
- ❑ comes from **common interosseous branch** of the ulnar artery .




Superficial Group

- 1) Brachioradialis
- 2) Extensor carpi radialis longus
- 3) Extensor carpi radialis brevis
- 4) Extensor digitorum
- 5) Extensor digiti minimi
- 6) Extensor carpi ulnaris
- 7) Anconeus

NOTE

Except for the brachioradialis and anconeus,

- **ALL have common origin** from the lateral epicondyle of the humerus
- **ALL have tendons** extend into the hand.




Deep Group


- 1) Supinator
- 2) Abductor pollicis longus
- 3) Extensor pollicis brevis
- 4) Extensor pollicis Longus
- 5) Extensor indicis

NOTE


Except for the supinator muscle, **ALL OF THEM**

- originate from the posterior surfaces of the radius, ulna, and interosseous membrane
- pass into the thumb and fingers in ratio 3:1









- Supinator**
- Abductor pollicis longus**
- Extensor pollicis longus and brevis**
- Extensor indicis**



- Brachioradialis**
- Extensor carpi radialis longus and brevis**
- Extensor digitorum**
- Extensor digit minimi**
- Extensor carpi ulnaris**
- Anconeus**








extensor expansion

- tendon of extensor digitorum expands to surround distal metacarpal head & proximal phalanx like a hood
- To which tendons of intrinsic mm. of hand (lumbricals & interossei) are inserted

Function

- ☐ serves to hold extensor tendons in place
- ☐ allow extensors, lumbricals, interossei to effect extension at proximal & distal interphalangeal joints





Posterior interosseous nerve

- Pure motor
- Origins
 - originates from the radial nerve at the radiohumeral joint li
- Course
 - Dips between heads of supinator (arcade of Frohse)
 - winds around radial neck within substance of muscle
 - reaches interosseous membrane of forearm
 - ends at dorsal wrist capsule

Motor innervation

all muscles of back of forearm except

1. Brachioradialis ,
2. extensor carpi radialis longus
3. anconeus.




Blood supply of the posterior compartment

❑ Posterior interosseous artery

- Originated from common interosseous branch of the ulnar artery
- Gives recurrent interosseous artery
- enters posterior compartment & descends with posterior interosseous nerve to supply forearm extensor muscles
- *End by joining to dorsal carpal arch of the wrist*

Anterior interosseous artery



JOINTS

Distal radio-ulnar joint

❖Type &variety :


synovial, pivot(uniaxial)

❖Articular surfaces:

- Head of ulnar medially
- Ulnar notch of lower end of radius laterally
- Triangular articular disc inferiorly

❖Capsule:

- attached around the articular surface
- Project upward (together with its lining synovial membrane) Forming small pouch between radius & ulna



Distal radio-ulnar joint

❖ Articular disc:

- thick triangular band of fibrocartilage

- ✓ apex

attach medially to styloid process of ulna

- ✓ Base

attached laterally to lower border of ulnar notch of the radius

- ✓ upper concave surface

articulate with head of ulna


- ✓ Lower convex surface

articulate with lunate triquetrum bone



- It Separate inferior radioulnar joint above from wrist joint below

❖ Movement:

Supination & pronation



- lesion of **median nerve** at or above elbow paralyzes the pronator quadratus & pronator teres muscles, with loss of ability to pronate forearm.
- When the supinator muscle is paralyzed from a radial nerve lesion, the biceps brachii muscle can still supinate the forearm.



WRIST JOINT

- **Type:**

Synovial, biaxial (ellipsoid) joint


Articular surfaces:

A. Proximal articular surface: Formed by:

1. The **inferior surface of lower end of radius**
 2. **Inferior surface of articular disc** of inferior radio-ulnar joint which lies below the head of ulna (separating it from the triquetrum bone & wrist joint).
- This proximal articular surface has **an ellipsoid outline** with a transverse diameter of about 5 cm. and an antero-posterior diameter of about 2 cm. It is slightly **concave in both directions**

B. Distal articular surface: Is formed of three carpal bones;

1. **Scaphoid and lunate** lie below the radius.
 2. **Triquetrum** lies below the articular disc below the head of ulna.
- These three bones form an oval surface which is **convex in all directions**.




WRIST JOINT

❖ Capsule and ligaments

1. The capsule is attached to the margins of the articular surfaces.
2. The lateral and medial parts of the capsule are thickened to form **radial (lateral) and ulnar (medial) collateral ligaments** of the wrist.
3. The capsule is strengthened in front and behind by **anterior and posterior (palmar and dorsal) radio-carpal ligaments** which pass obliquely downwards and medially from the lower end of the radius to the carpal bones.

They are important as they force the carpus to move with the radius as one unit during pronation and supination.



WRIST JOINT

Movements:

A. Flexion:

1. Flexor carpi radialis is a main flexor
2. Flexor carpi ulnaris is a main flexor
3. Palmaris longus
4. All the long muscles whose tendons cross in front of the wrist help in flexion.

B. Extension:

1. Extensor carpi radialis longus is a main extensor
2. Extensor carpi radialis brevis is a main extensor
3. Extensor carpi ulnaris is a main extensor
4. All the long muscles whose tendons cross behind the wrist help in extension.


C. Adduction:

1. Flexor carpi ulnaris
2. Extensor carpi ulnaris

D. Abduction:

1. Extensor carpi radialis longus
2. Extensor carpi radialis brevis
3. Flexor carpi radialis
4. Abductor pollicis longus

E. Circumduction: Combination of the above mentioned four movements.




SPECIFIC AREAS

RELATIONSHIP OF STRUCTURES ON THE FRONT OF THE WRIST

1) structures lying in front of the conventional wrist from lateral to medial side are:



1. Radial artery.
2. Tendon of flexor carpi radialis (FCR).
3. Median nerve
4. Tendon of palmaris longus.
5. Flexor digitorum superficialis.
6. Ulnar artery.
7. Ulnar nerve.
8. Tendon of flexor carpi ulnaris.



RELATIONSHIP OF STRUCTURES ON THE FRONT OF THE WRIST

2) Structures passing superficial to the flexor retinaculum (from med. To lat.)

- Ulnar n. (just lat. To pisiform bone)
- Ulnar a. (just lat. To ulnar n.)
- Palmar cutaneous br. Of ulnar n.
- Palmaris longus tendon
- Palmar cutaneous br. Of median n.



RELATIONSHIP OF STRUCTURES ON THE FRONT OF THE WRIST


3) Carpal tunnel

❖ Definition :

it is a fibro- osseous canal in front of wrist transmitting the tendons of the muscles from the forearm to the hand

❖ Boundaries

- **Posteriorly** :the bony groove formed by the carpal bones & their interosseous ligaments
- **Anteriorly** :it is closed by the flexor retinaculum of the wrist



Note:

Dense connective tissue converts the depression between the hook of hamate and the pisiform into **Guyon canal** (ulnar tunnel), where ulnar nerve may be compressed (e.g., by a ganglion cyst).

RELATIONSHIP OF STRUCTURES ON THE FRONT OF THE WRIST

3) Carpal tunnel

❖ Structures passing through it (deep to it

1-Median n. most superficial structure(immediately behind ftex. Retinaculum)

2- **4 tendons of flex. Digit. Superficialis** enclosed by their common flex. Synovial sheath (ulnar bursa)


3- **4 tendons of flex. Digit. profundus**

4-Tendon of flex. Pollicis longus within its own synovial sheath(radial bursa)

5-Recurrent br. Of deep palmar arch ascending infront of the carpal bones to joins the ant. Carpal anastomosis

❖ Clinical importance: carpal tunnel syndrome

Definition: condition in which there is a compression of the median n. as it passes through the tunnel leading to its injury



RELATIONSHIP OF STRUCTURES ON THE FRONT OF THE WRIST


4) Anatomical snuff box


Definition

Small depression on the posterolateral aspect of the wrist. It becomes more evident when the thumb is extended

Boundaries

- **Anterolaterally** :the tendons of the abd. Pollicis longus & ext. pollicis brevis
- **Posteromedially** :tendon of ext. pollicis longus
- **Roof** :formed by
 - 1-Skin
 - 2-Superficial fascia containing a) beginning of cephalic v. b) digital brs of superficial radial n.
 - 3- deep fascia
- **Floor** :formed by
 - 1-Styloid process of radius
 - 2-Scaphoid bone
 - 3-Trapezium





RELATIONSHIP OF STRUCTURES ON THE FRONT OF THE WRIST

4) Anatomical snuff box

Contents

- 1-Tendons of ext. carpi radialis longus & brevis m.
- 2-Radial a.

Clinical importance : pain & swelling in it occurs in fracture of the scaphoid bone



Clinical findings:
pain,
tenderness,
and swelling
in anatomical
snuffbox

