

Bone Fracture

Classification • Clinical Picture • First Aid

1. Classification of Fractures

According to skin:

- Closed (simple)
- Open (compound)

According to fracture line:

- Transverse, Oblique
- , Spiral,
- Comminuted

According to displacement:

- Undisplaced
- Displaced



Open fracture




Closed fracture







ransverse



Oblique



Spiral



Comminuted

2. Clinical Picture

- ❖ Pain — usually severe and worsened by movement
- ❖ Swelling and tenderness
- ❖ Bruising or discoloration
- ❖ Deformity
- ❖ Loss of function / inability to use the limb
- ❖ Abnormal mobility and possible crepitus (do not test deliberately)
- ❖ Open fracture: wound and possible exposed bone

Closed vs. Open Fractures

► Closed Fracture




► Open Fracture




3. First Aid

- ❖ Ensure safety and call emergency medical services
- ❖ Immobilize/splint in the position found
- ❖ Support the joints above and below the suspected fracture
- ❖ Do NOT straighten or realign the bone?
- ❖ For open fractures: cover the wound with a clean/sterile dressing
- ❖ Do NOT push exposed bone back inside



A



B



4. Additional First Aid

- ❖ Apply a cold pack wrapped in cloth near the injury
- ❖ Do not apply ice directly to the skin
- ❖ Keep the person warm and reassured
- ❖ Avoid food or drink if surgery/anesthesia may be required


Emergency Signs

- Severe or uncontrolled bleeding
- Difficulty breathing or loss of consciousness
- Suspected spinal or pelvic injury
- Absent pulse distal to the injury
- Cold, pale, or blue limb
- These require immediate emergency medical care


Pressure with Hands



Apply Dressing and Press



Tourniquet



Wrap

Wind

Secure

Time

Key Points

- ❖ Recognize: pain, swelling, deformity, and loss of function
- ❖ Protect: keep the fracture still and splint appropriately
- ❖ Do not manipulate or straighten the fracture
- ❖ Control bleeding and cover open wounds
- ❖ Check circulation and sensation before and after splinting
- ❖ Seek professional medical assessment