

CONCORDIA UNIVERSITY
Department of Mathematics and Statistics

| | | |
|-----------------------|-----------------------|-------------------|
| Course Mathematics | Number 204 | Section(s) All |
| Examination Final | Date December 2022 | Pages 3 |

Instructors

M. Khasraghi, S. Saputra, D. Basu, C. Pagano, M. Hindawi, S. Jessup

Course Examiner

A. Kokotov

Special Instructions: Only approved calculators are allowed !

1. [10] Let

$$A = \begin{pmatrix} 3 & 2 & 3 \\ 4 & 3 & 7 \\ 3 & 2 & 4 \end{pmatrix}$$

Solve the equation

$$AXA^{-1} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} + A \begin{bmatrix} 3 & 2 & 3 \\ 4 & 3 & 7 \\ 3 & 2 & 4 \end{bmatrix} = \begin{bmatrix} 4 & 3 & 4 \\ 4 & 3 & 4 \\ 4 & 3 & 4 \end{bmatrix}$$

for (3×3) matrix X .

2. [10] Find the determinant of (4×4) matrix X if

$$\det \left(\begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \\ 3 & 4 & 1 & 2 \\ 4 & 1 & 2 & 3 \end{pmatrix} X \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix} \right) = \det \left(\begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \right)$$

$$\det(ABC) = \det(A) \det(B) \det(C)$$

3. [10] For which a the system

$$\begin{cases} 3x_1 + 2x_2 + 3x_4 = 0 \\ 11x_1 + 4x_2 + ax_3 + 5x_4 = 0 \\ 4x_1 + 3x_2 + 7x_4 = 0 \\ 3x_1 + 2x_2 + x_3 + 4x_4 = 0 \end{cases}$$

has infinitely many solutions?

4. [10] Three lines l_1 , l_2 and l_3 pass through the origin $O = (0, 0, 0)$ and are parallel to the vectors $(0, 4, 2)$, $(-1, -3, -2)$ and $(10, 2, 4)$ respectively. Find the area of the triangle whose vertices are the points of intersection of these three lines with the plane

$$x + 2y - 3x = 1$$

5. [10] The plane α with equation

$$ax + by + 12z = 2$$

is parallel to the plane

$$x + 2y + 4x = \sqrt{2}.$$

The line l is the intersection of the plane α with the plane

$$x + y + z = 1.$$

$$\begin{array}{l} -14 - 66 + 80 \checkmark \\ 2 - 22 + 20 \checkmark \end{array}$$

Write down the parametric equation of the line l .

6. [10] A vector \vec{a} has norm 2 and is orthogonal to the vectors $\vec{b} = (7, 3, 4)$ and $\vec{c} = (-1, 1, 1)$. Find the norm of the vector $2\vec{a} - 0.5\vec{b} + \vec{c}$.

$$\vec{a} = (-1, -11, 10)$$

$$\|\vec{a}\| = \sqrt{1 + 121 + 100}$$

$$= \sqrt{122 + 100}$$

$$= \sqrt{222}$$

$$\frac{\vec{a}}{\|\vec{a}\|} = \frac{-1}{\sqrt{222}}, \frac{-11}{\sqrt{222}}, \frac{10}{\sqrt{222}}$$

$$2 \cdot \frac{\vec{a}}{\|\vec{a}\|} = \frac{-2}{\sqrt{222}}, \frac{-22}{\sqrt{222}}, \frac{20}{\sqrt{222}}$$

7. [10] For two square (2×2) matrices A and B one has the equalities

$$A^2 = \begin{pmatrix} 3 & 2 \\ 4 & 3 \end{pmatrix}; \quad B^2 = \begin{pmatrix} 2 & 4 \\ 4 & 10 \end{pmatrix};$$

$$AB = \begin{pmatrix} 2 & 4 \\ 3 & 5 \end{pmatrix}; \quad (A+B)^2 = \begin{pmatrix} 10 & 12 \\ 18 & 22 \end{pmatrix}$$

Find the matrix BA .

8. [10] Find a basis for the solution space of the homogeneous system

$$\begin{cases} x_1 + x_2 + x_3 + x_4 + x_5 = 0 \\ 2x_1 - x_2 - x_3 - x_4 + x_6 = 0 \\ 4x_1 + x_2 - x_3 + x_4 - 2x_5 + x_6 = 0 \end{cases}$$

9. [10] Find the eigenvalue of the matrix

$$A = \begin{pmatrix} 7 & 7 & 1 \\ 13 & 13 & -1 \\ 1 & -3 & 0 \end{pmatrix}$$

corresponding to an eigenvector that is orthogonal to the plane

$$x - y + 2z = -\sqrt{3}$$

10. [10] Find an invertible 2×2 matrix P such that the matrix

$$P^{-1} \begin{pmatrix} 17 & -6 \\ 35 & -12 \end{pmatrix} P$$

is diagonal.

¹The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.